



(19) World Intellectual Property Organization  
International Bureau

**(43) International Publication Date**  
**26 July 2001 (26.07.2001)**

**(10) International Publication Number**  
**WO 01/53454 A2**

**PCT**

(51) International Patent Classification <sup>7</sup> :	C12N	US	09/552,317 (CIP)
(21) International Application Number:	PCT/US00/34983	Filed on	25 April 2000 (25.04.2000)
(22) International Filing Date:	22 December 2000 (22.12.2000)	Filed on	09/598,042 (CIP)
(25) Filing Language:	English	Filed on	20 June 2000 (20.06.2000)
(26) Publication Language:	English	Filed on	09/620,312 (CIP)
(30) Priority Date:	21 January 2000 (21.01.2000)	Filed on	19 July 2000 (19.07.2000)
	00/488,775	US	09/653,450 (CIP)
		Filed on	31 August 2000 (31.08.2000)
		US	09/729,739 (CIP)
		Filed on	4 December 2000 (04.12.2000)
		US	Not furnished (CIP)
		Filed on	21 December 2000 (21.12.2000)

(71) Applicant (for all designated States except US): HYSEQ, INC. {USUS}; 670 Almanor Avenue, Sunnyvale, CA 94086 (US).

Not furnished	December 2000 (01.12.2000)	US
(72) Inventors; and		
(75) Inventor/Applicants (for US only):		
	Victoria JPUITS; 883 Portland Place, Redwood Shores, CA 94065 (US); TANG, Y., Tom (CN/US); 420 Ranwick Court, San Jose, CA 95118 (US); LIU, Chinghua (CN/US); 11255 Ranchero Way #14, San Jose, CA 95117 (US)	
(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:		
	09/488,725 (CIP)	
	Filed on 21 January 2000 (21.01.2000)	

[Continued on next page]

(54) Title: METHODS AND MATERIALS RELATING TO G PROTEIN-COUPLED RECEPTOR-LIKE POLYPEPTIDES AND POLYNYLIC ESTERS

BLASTP ALIGNMENT OF SEQ ID NO: 4, G PROTEIN-COUPLED RECEPTOR-LIKE POLYPEPTIDE (IDENTIFIED AS GPCR-LIKE) WITH HUMAN COT-40 PROTEIN SEQ ID NO: 48

187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600	601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

**(57) Abstract:** The invention provides novel polynucleotides and polypeptides encoded by such polynucleotides and mutants or variants thereof that correspond to a novel human secreted GPCR-like polypeptide. The polynucleotides comprise nucleic acid sequences isolated from cDNA libraries from human leukocyte mRNA (GIBCO Laboratories) (SEQ ID NO: 1), from human adult liver mRNA (GIBCO) (SEQ ID NO: 10); from human adult kidney mRNA (GIBCO) (SEQ ID NO: 17); from human adult brain mRNA (GIBCO) (SEQ ID NO: 26) and from human adult kidney mRNA (Invitrogen) (SEQ ID NO: 33). Other aspects of the invention include vectors for producing novel human secreted GPCR-like polypeptides, and antibodies specific for such polypeptides.

## METHODS AND MATERIALS RELATING TO

G PROTEIN-COUPLED RECEPTOR-LIKE POLYPEPTIDES AND  
POLYNUCLEOTIDES

5

## 1. TECHNICAL FIELD

The present invention provides novel polynucleotides and proteins encoded by such polynucleotides, along with uses for these polynucleotides and proteins, for example in therapeutic, diagnostic and research methods. In particular, the invention relates to novel G protein-coupled receptor-like (GPCR-like) polypeptides.

10

## 2. BACKGROUND ART

Identified polynucleotide and polypeptide sequences have numerous applications in, for example, diagnostics, forensics, gene mapping; identification of mutations responsible for genetic disorders or other traits, to assess biodiversity, and to produce many other types of data and products dependent on DNA and amino acid sequences. Proteins are known to have biological activity, for example, by virtue of their secreted nature in the case of leader sequence cloning, by virtue of their cell or tissue source in the case of PCR-based techniques, or by virtue of structural similarity to other genes of known biological activity. It is to these polypeptides and the polynucleotides encoding them that the present invention is directed. In particular, this invention is directed to novel GPCR-like polypeptides and polynucleotides.

20

Effective intercellular communication is obligatory for the successful survival of multicellular organisms. Environmental cues are normally recognized by a plethora of specific receptors present mainly on the cell membrane. Binding of the appropriate ligand activates the receptor, which initiates different signaling cascades that finally result in modification of cellular activity. Cells communicate with other cells, extracellular matrix, soluble hormones and chemokines, pheromones, toxins, viruses and bacteria using these receptors. The nature of the interactions and resulting signal transduction events define the fate of cell.

25

G protein-coupled receptors (GPCRs) constitute an evolutionarily conserved, but functionally very diverse family of such membrane receptors. All GPCR members share a common central seven transmembrane helices, termed TM-I through TM-VII; connected by three intracellular and three extracellular loops. Two conserved cysteine residues in these helices form a disulfide link that may be important for packing and stabilization of these seven TMs. The unique extracellular regions of individual GPCRs recognize specific ligands, the disulfide

30

bridge is implicated in interactions with agonists and antagonists, and the third intracellular loop interacts with G proteins that in turn activate second messengers such as cyclic adenosine monophosphate (cAMP), phospholipase C, inositol triphosphate, or ion channel proteins.

In vertebrates, the GPCR family contains more than 2000 gene members that can be subdivided into at least five subfamilies based on their ligand-binding properties. Family 1a binds small ligands including rhodopsin, odorants, and beta-adrenergic receptors and

5

interestingly the ligand-binding site is contained within the seven TM region. Family 1b binds small peptides and the binding site is located in the extracellular loop and the seven TM region, while family 1c binds large glycoproteins and the binding site is mainly located in the

10

extracellular domain with extracellular loops making some contacts. Family 2 is similar to family 1c with respect to ligand-binding but does not share any sequence similarities with family

1. Family 3 contains the  $\text{Ca}^{2+}$  sensing receptors while family 4 has pheromone receptors as its members. Finally, family 5 primarily consists of receptors involved in embryonic development.

Thus, GPCRs are involved in the recognition and transduction of messages as diverse as light,  $\text{Ca}^{2+}$ , odorants, small molecules including amino acids, nucleotides, lipids, and peptides,

15

hormones and pheromones, chemokines and complement, neurotransmitters, as well as larger proteins. GPCRs control the activity of enzymes, ion channels, and transport of vesicles via the catalysis of the GDP-GTP exchange on heterotrimeric G proteins ( $\text{G}\alpha$ - $\beta\gamma$ ) (Bockaert and Pin, (1999) EMBO J. 18, 1723-1729).

20

Olfactory GPCRs are responsible for transmission of volatile chemical signals from the environment through the olfactory neurons to the brain. Homologous receptors are also expressed in human testis and aid in sperm chemotaxis. Chemotactic GPCRs are also involved in immune response. Chemokines, platelet activating factor, and complement components all use GPCRs to transduce signals in the immune system. Regulation of GPCR activity is achieved at several levels. Apart from transcriptional and translational regulation, the GPCR family members have been shown to homo- and heterodimerize which can modulate their functions. Further, it has been shown that GPCRs can also interact with arrestins and certain PDZ domain containing proteins to transduce signals.

25

Abnormal GPCR function has been reported for various diseases including

hypothyroidism, familial precocious puberty, and congenital nephrogenic diabetes insipidus. Some of the GPCRs have been shown to function as proto-oncogenes and can be activated by mutagenesis. GPCRs are thus involved in many of the pathologies of human diseases.

30

Thus, the GPCR-like polypeptides and polynucleotides of the invention may be used in the treatment of diseases of ophthalmic, neurological, immunological, and nephritic systems.

They may also be used to treat hormonal dysfunction, cancer and other neoplasia, atherosclerosis, and diabetes.

### 3. SUMMARY OF THE INVENTION

5 This invention is based on the discovery of novel GPCR-like polypeptides, novel isolated polynucleotides encoding such polypeptides, including recombinant DNA molecules, cloned genes or degenerate variants thereof, especially naturally occurring variants such as allelic variants, antisense polynucleotide molecules, and antibodies that specifically recognize one or more epitopes present on such polypeptides, as well as hybridomas producing such 10 antibodies. Specifically, the polynucleotides of the present invention are based on GPCR-like polynucleotides isolated from cDNA libraries prepared from human leukocyte mRNA (GIBCO Laboratories) (SEQ ID NO: 1), from human adult liver mRNA (GIBCO) (SEQ ID NO: 10); from human adult kidney mRNA (GIBCO) (SEQ ID NO: 17); from human adult brain mRNA (GIBCO) (SEQ ID NO: 26) and from human adult kidney mRNA (Invitrogen) (SEQ ID NO: 33).

The compositions of the present invention additionally include vectors such as

expression vectors containing the polynucleotides of the invention, cells genetically engineered to contain such polynucleotides and cells genetically engineered to express such polynucleotides.

20 The compositions of the invention provide isolated polynucleotides that include, but are not limited to, a polynucleotide comprising the nucleotide sequence set forth in SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62; or a fragment of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62; a polynucleotide comprising the full length protein coding sequence of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 (for example, SEQ ID NO: 4, 13, 20, 29, 36, and 42); and a polynucleotide comprising the nucleotide sequence of the mature protein coding sequence of any of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62. The polynucleotides of the present invention also include, but are not limited to, a polynucleotide that hybridizes under stringent hybridization conditions to (a) the complement of any of the nucleotide sequences set forth in SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62; (b) a nucleotide sequence encoding any of SEQ ID NO: 4, 6-9, 13, 15-16, 20, 22-25, 29, 31-32, 36, 38-40, 42, 44-47, 61, or 63; a polynucleotide which is an allelic variant of any polynucleotides recited above having at least 70% polynucleotide sequence identity to the polynucleotides; a polynucleotide which encodes a species homolog (e.g. orthologs) of any of the

peptides recited above; or a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of the polypeptide comprising SEQ ID NO: 4, 13, 20, 29, 36, or 42. Preferably the polynucleotides include a polynucleotide comprising the sequence set forth in nucleotides 1-1351 of SEQ ID NO: 19 or nucleotides 271-1351 of SEQ ID NO: 19. Preferably the polynucleotides include a polynucleotide comprising the sequence set forth in nucleotides 1-1668 of SEQ ID NO: 28, nucleotides 52-1668 of SEQ ID NO: 28, or nucleotides 2845-3993 of SEQ ID NO: 28.

A collection as used in this application can be a collection of only one polynucleotide.

The collection of sequence information or unique identifying information of each sequence can be provided on a nucleic acid array. In one embodiment, segments of sequence information are provided on a nucleic acid array to detect the polynucleotide that contains the segment. The array can be designed to detect full-match or mismatch to the polynucleotide that contains the segment. The collection can also be provided in a computer-readable format.

15 This invention further provides cloning or expression vectors comprising at least a fragment of the polynucleotides set forth above and host cells or organisms transformed with these expression vectors. Useful vectors include plasmids, cosmids, lambda phage derivatives, phagemids, and the like, that are well known in the art. Accordingly, the invention also provides a vector including a polynucleotide of the invention and a host cell containing the polynucleotide. In general, the vector contains an origin of replication functional in at least one organism, convenient restriction endonuclease sites, and a selectable marker for the host cell. Vectors according to the invention include expression vectors, replication vectors, probe generation vectors, and sequencing vectors. A host cell according to the invention can be a prokaryotic or eukaryotic cell and can be a unicellular organism or part of a multicellular organism.

25 The compositions of the present invention include polypeptides comprising, but not limited to, an isolated polypeptide selected from the group comprising the amino acid sequence of SEQ ID NO: 4, 6-9, 13, 15-16, 20, 22-25, 29, 31-32, 36, 38-40, 42, 44-47, 61, or 63; or the corresponding full length or mature protein. Polypeptides of the invention also include polypeptides with biological activity that are encoded by (a) any of the polynucleotides having a nucleotide sequence set forth in SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62; or (b) polynucleotides that hybridize to the complement of the polynucleotides of (a) under stringent hybridization conditions. Biologically or immunologically active variants of any of the protein sequences listed as SEQ ID NO: 4, 6-9, 13, 15-16, 20, 22-25, 29, 31-32, 36, 38-40, 42, 44-47, 61, or 63 and substantial equivalents thereof that retain biological or immunological activity are also contemplated. Preferably the polypeptides include a polypeptide

comprising the sequence set forth in amino acid residues 1-360 of SEQ ID NO: 20. Preferably the polypeptides include a polypeptide comprising the sequence set forth in amino acid residues 1-539 of SEQ ID NO: 29 or the sequence set forth in amino acid residues 932-1314 of SEQ ID NO: 29. The polypeptides of the invention may be wholly or partially chemically synthesized but are preferably produced by recombinant means using the genetically engineered cells (e.g. host cells) of the invention.

The invention also provides compositions comprising a polypeptide of the invention.

Pharmaceutical compositions of the invention may comprise a polypeptide of the invention and an acceptable carrier, such as a hydrophilic, e.g., pharmaceutically acceptable, carrier.

The invention also relates to methods for producing a polypeptide of the invention comprising culturing host cells comprising an expression vector containing at least a fragment of a polynucleotide encoding the polypeptide of the invention in a suitable culture medium under conditions permitting expression of the desired polypeptide, and purifying the protein or peptide from the culture or from the host cells. Preferred embodiments include those in which the protein produced by such a process is a mature form of the protein.

Polynucleotides according to the invention have numerous applications in a variety of techniques known to those skilled in the art of molecular biology. These techniques include use as hybridization probes, use as oligomers, or primers, for PCR, use in an array, use in computer-readable media, use for chromosome and gene mapping, use in the recombinant production of protein, and use in generation of antisense DNA or RNA, their chemical analogs and the like. For example, when the expression of an mRNA is largely restricted to a particular cell or tissue type, polynucleotides of the invention can be used as hybridization probes to detect the presence of the particular cell or tissue mRNA in a sample using, e.g., *in situ* hybridization.

In other exemplary embodiments, the polynucleotides are used in diagnostics as expressed sequence tags for identifying expressed genes or, as well known in the art and exemplified by Vollrath et al., Science 258:52-59 (1992), as expressed sequence tags for physical mapping of the human genome.

The polypeptides according to the invention can be used in a variety of conventional procedures and methods that are currently applied to other proteins. For example, a polypeptide of the invention can be used to generate an antibody that specifically binds the polypeptide. Such antibodies, particularly monoclonal antibodies, are useful for detecting or quantitating the polypeptide in tissue. The polypeptides of the invention can also be used as molecular weight markers, and as a food supplement.

Methods are also provided for preventing, treating, or ameliorating a medical condition which comprises the step of administering to a mammalian subject a therapeutically effective amount of a composition comprising a peptide of the present invention and a pharmaceutically acceptable carrier.

Thus, the GPCR-like polypeptides and polynucleotides of the invention may be used in the treatment of diseases of ophthalmic, neurological, immunological, and nephritic systems. They may also be used to treat hormonal dysfunction, cancer and other neoplasia, atherosclerosis, and diabetes.

The methods of the invention also provide methods for the treatment of disorders as recited herein which comprise the administration of a therapeutically effective amount of a composition comprising a polynucleotide or polypeptide of the invention and a pharmaceutically acceptable carrier to a mammalian subject exhibiting symptoms or tendencies related to disorders as recited herein. In addition, the invention encompasses methods for treating diseases or disorders as recited herein comprising the step of administering a composition comprising compounds and other substances that modulate the overall activity of the target gene products and a pharmaceutically acceptable carrier. Compounds and other substances can effect such modulation either on the level of target gene/protein expression or target protein activity. Specifically, methods are provided for preventing, treating or ameliorating a medical condition, including viral diseases, which comprises administering to a mammalian subject, including but not limited to humans, a therapeutically effective amount of a composition comprising a polypeptide of the invention or a therapeutically effective amount of a composition comprising a binding partner of (e.g., antibody specifically reactive for) GPCR-like polypeptides of the invention. The mechanics of the particular condition or pathology will dictate whether the polypeptides of the invention or binding partners (or inhibitors) of these would be beneficial to the individual in need of treatment.

According to this method, polypeptides of the invention can be administered to produce an *in vitro* or *in vivo* inhibition of cellular function. A polypeptide of the invention can be administered *in vivo* alone or as an adjunct to other therapies. Conversely, protein or other active ingredients of the present invention may be included in formulations of a particular agent to minimize side effects of such an agent.

The invention further provides methods for manufacturing medicaments useful in the above-described methods.

The present invention further relates to methods for detecting the presence of the polynucleotides or polypeptides of the invention in a sample (e.g., tissue or sample). Such



methods can, for example, be utilized as part of prognostic and diagnostic evaluation of disorders as recited herein and for the identification of subjects exhibiting a predisposition to such conditions.

The invention provides a method for detecting a polypeptide of the invention in a sample comprising contacting the sample with a compound that binds to and forms a complex with the polypeptide under conditions and for a period sufficient to form the complex and detecting formation of the complex, so that if a complex is formed, the polypeptide is detected. The invention also provides kits comprising polynucleotide probes and/or monoclonal antibodies, and optionally quantitative standards, for carrying out methods of the invention. Furthermore, the invention provides methods for evaluating the efficacy of drugs, and monitoring the progress of patients, involved in clinical trials for the treatment of disorders as recited above.

The invention also provides methods for the identification of compounds that modulate (i.e., increase or decrease) the expression or activity of the polynucleotides and/or polypeptides of the invention. Such methods can be utilized, for example, for the identification of compounds that can ameliorate symptoms of disorders as recited herein. Such methods can include, but are not limited to, assays for identifying compounds and other substances that interact with (*e.g.*, bind to) the polypeptides of the invention.

The invention provides a method for identifying a compound that binds to the polypeptide of the present invention comprising contacting the compound with the polypeptide under conditions and for a time sufficient to form a polypeptide/compound complex and detecting the complex, so that if the polypeptide/compound complex is detected, a compound that binds to the polypeptide is identified.

Also provided is a method for identifying a compound that binds to the polypeptide comprising contacting the compound with the polypeptide in a cell for a time sufficient to form a polypeptide/compound complex wherein the complex drives expression of a reporter gene sequence in the cell and detecting the complex by detecting reporter gene sequence expression so that if the polypeptide/compound complex is detected a compound that binds to the polypeptide is identified.

#### 4. BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 3 (i.e. SEQ ID NO: 4) GPCR-like polypeptide and human CGI-40 protein (Lai et al, (2000) Genome Res. 10, 703-713) (SEQ ID NO: 48), indicating that the two

sequences share 76% similarity over 526 amino acid residues and 63% identity over the same 526 amino acid residues, wherein A=Alanine, C=Cysteine, D=Aspartic Acid, E= Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine. Gaps are presented as dashes.

Figure 2 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 3 (i.e. SEQ ID NO: 4) GPCR-like polypeptide and protein of clone CT748\_2 (Patent Application No. WO9824905) (SEQ ID NO: 49), indicating that the two sequences share 97% similarity over 445 amino acid residues and 96% identity over the same 445 amino acid residues, wherein A=Alanine, C=Cysteine, D=Aspartic Acid, E= Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine. Gaps are presented as dashes.

Figure 3 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 12 (i.e. SEQ ID NO: 13) GPCR-like polypeptide and human six transmembrane epithelial antigen of prostate (Hubert et al, (1999) Proc. Natl. Acad. Sci. U.S.A. 96, 14523-14528) (SEQ ID NO: 50), indicating that the two sequences share 68% similarity over 267 amino acid residues and 47% identity over the same 267 amino acid residues, wherein A=Alanine, C=Cysteine, D=Aspartic Acid, E= Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine. Gaps are presented as dashes.

Figure 4 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 12 (i.e. SEQ ID NO: 13) GPCR-like polypeptide and human STRAP-1 protein (Patent Application No. WO9962941) (SEQ ID NO: 51), indicating that the two sequences share 68% similarity over 267 amino acid residues and 47% identity over the same 267 amino acid residues, wherein A=Alanine, C=Cysteine, D=Aspartic Acid, E= Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine. Gaps are presented as dashes.

Figure 5A, 5B, and 5C show the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 19 (i.e. SEQ ID NO: 20) GPCR-like polypeptide and rat seven transmembrane receptor protein (Abe et al, (1999) J. Biol. Chem. 274, 19957-19964) (SEQ ID NO: 52), indicating that the two sequences share 81% similarity over 1354 amino

acid residues and 72 % identity over the same 1354 amino acid residues, wherein A = Alanine, C = Cysteine, D = Aspartic Acid, E = Glutamic Acid, F = Phenylalanine, G = Glycine, H = Histidine, I = Isoleucine, K = Lysine, L = Leucine, M = Methionine, N = Asparagine, P = Proline, Q = Glutamine, R = Arginine, S = Serine, T = Threonine, V = Valine, W = Tryptophan, Y = Tyrosine. Gaps are presented as dashes.

5 Figure 6A, and 6B show the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 19 (i.e. SEQ ID NO: 20) GPCR-like polypeptide and human brain-derived G protein-coupled receptor protein (Patent Application No. WO200008053) (SEQ ID NO: 53), indicating that the two sequences share 100 % similarity over 986 amino acid residues and 100 % identity over the same 986 amino acid residues, wherein A = Alanine, C = Cysteine, D = Aspartic Acid, E = Glutamic Acid, F = Phenylalanine, G = Glycine, H = Histidine, I = Isoleucine, K = Lysine, L = Leucine, M = Methionine, N = Asparagine, P = Proline, Q = Glutamine, R = Arginine, S = Serine, T = Threonine, V = Valine, W = Tryptophan, Y = Tyrosine. Gaps are presented as dashes.

15 Figure 7 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 28 (i.e. SEQ ID NO: 29) GPCR-like polypeptide and putative seven pass transmembrane protein (Spangenberg et al (1998) Genomics 48, 178-185) (SEQ ID NO: 54), indicating that the two sequences share 72 % similarity over 323 amino acid residues and 57 % identity over the same 323 amino acid residues, wherein A = Alanine, C = Cysteine, D = Aspartic Acid, E = Glutamic Acid, F = Phenylalanine, G = Glycine, H = Histidine, I = Isoleucine, K = Lysine, L = Leucine, M = Methionine, N = Asparagine, P = Proline, Q = Glutamine, R = Arginine, S = Serine, T = Threonine, V = Valine, W = Tryptophan, Y = Tyrosine. Gaps are presented as dashes.

25 Figure 8 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 28 (i.e. SEQ ID NO: 29) GPCR-like polypeptide and human h-TRAAP polypeptide #1 (Patent Application No. WO200026253) (SEQ ID NO: 55), indicating that the two sequences share 100 % similarity over 392 amino acid residues and 100 % identity over the same 392 amino acid residues, wherein A = Alanine, C = Cysteine, D = Aspartic Acid, E = Glutamic Acid, F = Phenylalanine, G = Glycine, H = Histidine, I = Isoleucine, K = Lysine, L = Leucine, M = Methionine, N = Asparagine, P = Proline, Q = Glutamine, R = Arginine, S = Serine, T = Threonine, V = Valine, W = Tryptophan, Y = Tyrosine. Gaps are presented as dashes.

30 Figure 9 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 35 (i.e. SEQ ID NO: 36) GPCR-like polypeptide and human olfactory

receptor (Rouquier et al, (1998) Nature Genet. 18 (3), 243-250) (SEQ ID NO: 56), indicating that the two sequences share 92 % similarity over 166 amino acid residues and 87 % identity over the same 166 amino acid residues, wherein A = Alanine, C = Cysteine, D = Aspartic Acid, E = Glutamic Acid, F = Phenylalanine, G = Glycine, H = Histidine, I = Isoleucine, K = Lysine, L = Leucine, M = Methionine, N = Asparagine, P = Proline, Q = Glutamine, R = Arginine, S = Serine, T = Threonine, V = Valine, W = Tryptophan, Y = Tyrosine. Gaps are presented as dashes.

5 Figure 10 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 35 (i.e. SEQ ID NO: 36) GPCR-like polypeptide and human G protein-coupled receptor GPR1 protein (Patent Application No. WO9630406) (SEQ ID NO: 57), indicating that the two sequences share 93 % similarity over 171 amino acid residues of and 92 % identity over the same 171 amino acid residues, wherein A = Alanine, C = Cysteine, D = Aspartic Acid, E = Glutamic Acid, F = Phenylalanine, G = Glycine, H = Histidine, I = Isoleucine, K = Lysine, L = Leucine, M = Methionine, N = Asparagine, P = Proline, Q = Glutamine, R = Arginine, S = Serine, T = Threonine, V = Valine, W = Tryptophan, Y = Tyrosine. Gaps are presented as dashes.

15 Figure 11 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 41 (i.e. SEQ ID NO: 42) GPCR-like polypeptide and human gene AC005587, similar to mouse olfactory receptor 13 polypeptide (SEQ ID NO: 58), indicating that the two sequences share 81 % similarity over 304 amino acid residues and 68 % identity over the same 304 amino acid residues, wherein A = Alanine, C = Cysteine, D = Aspartic Acid, E = Glutamic Acid, F = Phenylalanine, G = Glycine, H = Histidine, I = Isoleucine, K = Lysine, L = Leucine, M = Methionine, N = Asparagine, P = Proline, Q = Glutamine, R = Arginine, S = Serine, T = Threonine, V = Valine, W = Tryptophan, Y = Tyrosine. Gaps are presented as dashes.

25 Figure 12 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 41 (i.e. SEQ ID NO: 42) GPCR-like polypeptide and human G protein-coupled receptor GPR1 polypeptide (Patent Application No. WO9630406) (SEQ ID NO: 59), indicating that the two sequences share 91 % similarity over 287 amino acid residues and 90 % identity over the same 287 amino acid residues, wherein A = Alanine, C = Cysteine, D = Aspartic Acid, E = Glutamic Acid, F = Phenylalanine, G = Glycine, H = Histidine, I = Isoleucine, K = Lysine, L = Leucine, M = Methionine, N = Asparagine, P = Proline, Q = Glutamine, R = Arginine, S = Serine, T = Threonine, V = Valine, W = Tryptophan, Y = Tyrosine. Gaps are presented as dashes.

## S. DETAILED DESCRIPTION OF THE INVENTION

The GPCR-like polypeptide of SEQ ID NO: 4 is an approximately 827-amino acid transmembrane protein with a predicted molecular mass of approximately 93 kDa unglycosylated. Hyseq's sequence database searches using the Pfam models that were

5 categorized under G protein-coupled receptors and using the hmmssearch program (hmmssearch - search a sequence database with a profile HMM: HMMER 2.1.1 (Dec 1998) Washington University School of Medicine), SEQ ID NO 4 was found to be homologous to G protein-coupled receptor model sequences with an E-value of 0.011. The homologous sequence identified using Pfam hmmssearch is shown in SEQ ID NO: 6. Further analyses with protein database searches with the BLASTP algorithm (Altschul S.F. et al., *J. Mol. Evol.* 36:290-300 (1993) and Altschul S.F. et al., *J. Mol. Biol.* 21:403-10 (1990), herein incorporated by reference) indicate that SEQ ID NO: 4 is homologous to human CGI-40 protein and with protein of clone CT748\_2.

Figure 1 shows the BLASTP amino acid sequence alignment between the protein

15 encoded by SEQ ID NO: 3 (i.e. SEQ ID NO: 4) GPCR-like polypeptide and human CGI-40 protein (Lai et al. (2000) *Genome Res.* 10, 703-713) (SEQ ID NO: 48), indicating that the two sequences share 76% similarity over 526 amino acid residues and 63% identity over the same 526 amino acid residues.

Figure 2 shows the BLASTP amino acid sequence alignment between the protein

20 encoded by SEQ ID NO: 3 (i.e. SEQ ID NO: 4) GPCR-like polypeptide and protein of clone CT748\_2 (Patent Application No. WO9824905) (SEQ ID NO: 49), indicating that the two sequences share 97% similarity over 445 amino acid residues and 96% identity over the same 445 amino acid residues.

A predicted approximately nineteen-residue signal peptide is encoded from approximately residue 1 through residue 19 of SEQ ID NO: 4 (SEQ ID NO: 7). The extracellular portion is useful on its own. This can be confirmed by expression in mammalian cells and sequencing of the cleaved product. The signal peptide region was predicted using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). One of skill in the art will recognize that the actual cleavage site may be different than that predicted by the computer program.

The GPCR-like polypeptide of SEQ ID NO: 13 is an approximately 488-amino acid transmembrane protein with a predicted molecular mass of approximately 55-kDa unglycosylated. Hyseq's sequence database searches using the Pfam models that were categorized under G protein-coupled receptors and using the hmmssearch program (hmmssearch

- search a sequence database with a profile HMM: HMMER 2.1.1 (Dec 1998) Washington University School of Medicine), SEQ ID NO 13 was found to be homologous to G protein-coupled receptor model sequences with an E-value of 0.017. The homologous sequence identified using Pfam hmmssearch is shown in SEQ ID NO: 15. Protein database searches with the BLASTP algorithm (Altschul S.F. et al., *J. Mol. Evol.* 36:290-300 (1993) and Altschul

5 S.F. et al., *J. Mol. Biol.* 21:403-10 (1990), herein incorporated by reference) indicate that SEQ ID NO: 4 is homologous to six transmembrane epithelial antigen of prostate and with human STRAP-1 protein.

Figure 3 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 12 (i.e. SEQ ID NO: 13) GPCR-like polypeptide and human six transmembrane epithelial antigen of prostate (Hubert et al. (1999) *Proc. Natl. Acad. Sci. U.S.A.* 96, 14523-14528) (SEQ ID NO: 50), indicating that the two sequences share 68% similarity over 267 amino acid residues and 47% identity over the same 267 amino acid residues.

Figure 4 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 12 (i.e. SEQ ID NO: 13) GPCR-like polypeptide and human STRAP-1 protein (Patent Application No. WO9962941) (SEQ ID NO: 51), indicating that the two sequences share 68% similarity over 267 amino acid residues and 47% identity over the same 267 amino acid residues.

20 The GPCR-like polypeptide of SEQ ID NO: 20 is an approximately 1346-amino acid transmembrane protein with a predicted molecular mass of approximately 151-kDa unglycosylated. Hyseq's sequence database searches using the Pfam models that were categorized under G protein-coupled receptors and using the hmmssearch program (hmmssearch - search a sequence database with a profile HMM: HMMER 2.1.1 (Dec 1998) Washington

25 University School of Medicine), SEQ ID NO 20 was found to be homologous to G protein-coupled receptor model sequences with an E-value of 2.7e-24. The homologous sequence identified using Pfam hmmssearch is shown in SEQ ID NO: 22. Further analyses with protein database searches with the BLASTP algorithm (Altschul S.F. et al., *J. Mol. Evol.* 36:290-300 (1993) and Altschul S.F. et al., *J. Mol. Biol.* 21:403-10 (1990), herein incorporated by reference) indicate that SEQ ID NO: 20 is homologous to the rat seven transmembrane receptor and to the human brain-derived G protein-coupled receptor proteins.

Figure 5A, 5B, and 5C show the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 19 (i.e. SEQ ID NO: 20) GPCR-like polypeptide and rat seven transmembrane receptor protein (Abe et al. (1999) *J. Biol. Chem.* 274, 19957-19964)

(SEQ ID NO: 52), indicating that the two sequences share 81 % similarity over 1354 amino acid residues and 72 % identity over the same 1354 amino acid residues.

Figure 6A, and 6B show the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 19 (i.e. SEQ ID NO: 20) GPCR-like polypeptide and human brain-derived G protein-coupled receptor protein (Patent Application No. WO200008053) (SEQ ID NO: 53), indicating that the two sequences share 100 % similarity over 986 amino acid residues and 100 % identity over the same 986 amino acid residues.

A predicted approximately twenty one-residue signal peptide is encoded from approximately residue 1 through residue 21 of SEQ ID NO: 20 (SEQ ID NO: 23). The extracellular portion is useful on its own. This can be confirmed by expression in mammalian cells and sequencing of the cleaved product. The signal peptide region was predicted using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). One of skill in the art will recognize that the actual cleavage site may be different than that predicted by the computer program.

The GPCR-like polypeptide of SEQ ID NO: 29 is an approximately 1314-amino acid transmembrane protein with a predicted molecular mass of approximately 147-kDa unglycosylated. Hyseq's sequence database searches with the Pfam models that were categorized under G protein-coupled receptors using the hmsearch program (hmsearch - search a sequence database with a profile HMM: HMMER 2.1.1 (Dec 1998) Washington University School of Medicine), SEQ ID NO 29 was found to be homologous to G protein-coupled receptor model sequences with an E-value of 0.0036. The homologous sequence identified using Pfam hmsearch is shown in SEQ ID NO: 31. Further analyses with protein database searches with the BLASTP algorithm (Altschul S.F. et al., J. Mol. Evol. 36:290-300 (1993) and Altschul S.F. et al., J. Mol. Biol. 21:403-10 (1990), herein incorporated by reference) indicate that SEQ ID NO: 29 is homologous to the putative seven pass transmembrane protein and to the human h-TRAAK polypeptide #1.

Figure 7 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 28 (i.e. SEQ ID NO: 29) GPCR-like polypeptide and putative seven pass transmembrane protein (Spangenberg et al (1998) Genomics 48, 178-185) (SEQ ID NO: 54), indicating that the two sequences share 72 % similarity over 323 amino acid residues and 57 % identity over the same 323 amino acid residues.

Figure 8 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 28 (i.e. SEQ ID NO: 29) GPCR-like polypeptide and human h-TRAAK polypeptide #1 (Patent Application No. WO2000026253) (SEQ ID NO: 55), indicating that the

two sequences share 100 % similarity over 392 amino acid residues and 100 % identity over the same 392 amino acid residues.

The GPCR-like polypeptide of SEQ ID NO: 36 is an approximately 194-amino acid transmembrane protein with a predicted molecular mass of approximately 22-kDa unglycosylated. Hyseq's sequence database searches using the Pfam models that were categorized under G protein-coupled receptors and using the hmsearch program (hmsearch - search a sequence database with a profile HMM: HMMER 2.1.1 (Dec 1998) Washington University School of Medicine), SEQ ID NO 36 was found to be homologous to G protein-coupled receptor model sequences with an E-value of 1.8e-28. The homologous sequence identified using Pfam hmsearch is shown in SEQ ID NO: 38. Further analyses with protein database searches with the BLASTP algorithm (Altschul S.F. et al., J. Mol. Evol. 36:290-300 (1993) and Altschul S.F. et al., J. Mol. Biol. 21:403-10 (1990), herein incorporated by reference) indicate that SEQ ID NO: 36 is homologous to the human olfactory receptor protein and to the human G protein-coupled receptor GPR1 polypeptide.

Figure 9 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 35 (i.e. SEQ ID NO: 36) GPCR-like polypeptide and human olfactory receptor (Rouquier et al, (1998) Nature Genet. 18 (3), 243-250) (SEQ ID NO: 56), indicating that the two sequences share 92 % similarity over 166 amino acid residues and 87 % identity over the same 166 amino acid residues.

Figure 10 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 35 (i.e. SEQ ID NO: 36) GPCR-like polypeptide and human G protein-coupled receptor GPR1 protein (Patent Application No. WO9630406) (SEQ ID NO: 57), indicating that the two sequences share 93 % similarity over 171 amino acid residues of and 92 % identity over the same 171 amino acid residues.

A predicted approximately thirty five-residue signal peptide is encoded from approximately residue 1 through residue 35 of SEQ ID NO: 36 (SEQ ID NO: 39). The extracellular portion is useful on its own. This can be confirmed by expression in mammalian cells and sequencing of the cleaved product. The signal peptide region was predicted using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). One of skill in the art will recognize that the actual cleavage site may be different than that predicted by the computer program.

The GPCR-like polypeptide of SEQ ID NO: 42 is an approximately 308-amino acid transmembrane protein with a predicted molecular mass of approximately 34-kDa unglycosylated. Hyseq's sequence database searches using the Pfam models that were

categorized under G protein-coupled receptors and using the hmmssearch program (hmmssearch - search a sequence database with a profile HMM: HMMER 2.1.1 (Dec 1998) Washington University School of Medicine), SEQ ID NO 42 was found to be homologous to G protein-coupled receptor model sequences with an E-value of 1.1e-47. The homologous sequence identified using Pfam hmmssearch is shown in SEQ ID NO: 44. Further analyses with protein database searches with the BLASTP algorithm (Altschul S.F. et al., J. Mol. Evol. 36:290-300 (1993) and Altschul S.F. et al., J. Mol. Biol. 21:403-10 (1990), herein incorporated by reference) indicate that SEQ ID NO: 42 is homologous to the mouse olfactory receptor 13 polypeptide and to the human G protein-coupled receptor GPR1 polypeptide.

Figure 11 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 41 (i.e. SEQ ID NO: 42) GPCR-like polypeptide and human gene AC005587, similar to mouse olfactory receptor 13 polypeptide (SEQ ID NO: 58), indicating that the two sequences share 81 % similarity over 304 amino acid residues and 68 % identity over the same 304 amino acid residues.

Figure 12 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 41 (i.e. SEQ ID NO: 42) GPCR-like polypeptide and human G protein-coupled receptor GPR1 polypeptide (Patent Application No. WO9630406) (SEQ ID NO: 59), indicating that the two sequences share 91 % similarity over 287 amino acid residues and 90 % identity over the same 287 amino acid residues.

A predicted approximately forty two-residue signal peptide is encoded from approximately residue 1 through residue 42 of SEQ ID NO: 42 (SEQ ID NO: 45). The extracellular portion is useful on its own. This can be confirmed by expression in mammalian cells and sequencing of the cleaved product. The signal peptide region was predicted using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). One of skill in the art will recognize that the actual cleavage site may be different than that predicted by the computer program.

Thus, the GPCR-like polypeptides and polynucleotides of the invention may be used in the treatment of diseases of ophthalmic, neurological, immunological, and nephritic systems. They may also be used to treat hormonal dysfunction, cancer and other neoplasia, atherosclerosis, and diabetes.

## 5.1 DEFINITIONS

It must be noted that as used herein and in the appended claims, the singular forms "a", "an" and "the" include plural references unless the context clearly dictates otherwise.

The term "active" refers to those forms of the polypeptide that retain the biologic and/or immunologic activities of any naturally occurring polypeptide. According to the invention, the terms "biologically active" or "biological activity" refer to a protein or peptide having structural, regulatory or biochemical functions of a naturally occurring molecule.

Likewise "biologically active" or "biological activity" refers to the capability of the natural, recombinant or synthetic GPCR-like peptide, or any peptide thereof, to induce a specific biological response in appropriate animals or cells and to bind with specific antibodies. The term "GPCR-like biological activity" refers to biological activity that is similar to the biological activity of a GPCR-like polypeptide.

The term "activated cells" as used in this application are those cells which are engaged in extracellular or intracellular membrane trafficking, including the export of secretory or enzymatic molecules as part of a normal or disease process.

The terms "complementary" or "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence 5'-AGT-3' binds to the complementary sequence 3'-TCA-5'. Complementarity between two single-stranded molecules may be "partial" such that only some of the nucleic acids bind or it may be "complete" such that total complementarity exists between the single stranded molecules. The degree of complementarity between the nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands.

The term "embryonic stem cells (ES)" refers to a cell that can give rise to many differentiated cell types in an embryo or an adult, including the germ cells. The term "germ line stem cells (GSCs)" refers to stem cells derived from primordial stem cells that provide a steady and continuous source of germ cells for the production of gametes. The term "primordial germ cells (PGCs)" refers to a small population of cells set aside from other cell lineages particularly from the yolk sac, mesenteries, or gonadal ridges during embryogenesis that have the potential to differentiate into germ cells and other cells. PGCs are the source from which GSCs and ES cells are derived. The PGCs, the GSCs and the ES cells are capable of self-renewal. Thus these cells not only populate the germ line and give rise to a plurality of terminally differentiated cells that comprise the adult specialized organs, but are able to regenerate themselves.

The term "expression modulating fragment," EMF, means a series of nucleotides that modulates the expression of an operably linked ORF or another EMF.

As used herein, a sequence is said to "modulate the expression of an operably linked sequence" when the expression of the sequence is altered by the presence of the EMF. EMFs

include, but are not limited to, promoters, and promoter modulating sequences (inducible elements). One class of EMFs is nucleic acid fragments which induce the expression of an operably linked ORF in response to a specific regulatory factor or physiological event.

The terms "nucleotide sequence" or "nucleic acid" or "polynucleotide" or "oligonucleotide" are used interchangeably and refer to a heteropolymers of nucleotides or the sequence of these nucleotides. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA) or to any DNA-like or RNA-like material. In the sequences, A is adenine, G is guanine, C is cytosine, T is thymine, and N is A, G, C, or T (U). It is contemplated that where the polynucleotide is RNA, the T (thymine) in the sequence herein may be replaced with U (uracil). Generally, nucleic acid segments provided by this invention may be assembled from fragments of the genome and short oligonucleotide linkers, or from a series of oligonucleotides, or from individual nucleotides, to provide a synthetic nucleic acid which is capable of being expressed in a recombinant transcriptional unit comprising regulatory elements derived from a microbial or viral operon, or a eukaryotic gene.

The terms "oligonucleotide fragment" or a "polynucleotide fragment", "portion," or "segment" or "probe" or "primer" are used interchangeably and refer to a sequence of nucleotide residues which are at least about 5 nucleotides, more preferably at least about 7 nucleotides, more preferably at least about 9 nucleotides, more preferably at least about 11 nucleotides and most preferably at least about 17 nucleotides. The fragment is preferably less than about 500 nucleotides, preferably less than about 200 nucleotides, more preferably less than about 100 nucleotides, more preferably less than about 50 nucleotides and most preferably less than 30 nucleotides. Preferably the probe is from about 6 nucleotides to about 200 nucleotides, preferably from about 15 to about 50 nucleotides, more preferably from about 17 to 30 nucleotides and most preferably from about 20 to 25 nucleotides. Preferably the fragments can be used in polymerase chain reaction (PCR), various hybridization procedures or microarray procedures to identify or amplify identical or related parts of mRNA or DNA molecules. A fragment or segment may uniquely identify each polynucleotide sequence of the present invention. Preferably the fragment comprises a sequence substantially similar to a portion of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62.

Probes may, for example, be used to determine whether specific mRNA molecules are present in a cell or tissue or to isolate similar nucleic acid sequences from chromosomal DNA as described by Walsh et al. (Walsh, P.S. et al., 1992, PCR Methods Appl 1:241-250). They

may be labeled by nick translation, Klenow fill-in reaction, PCR, or other methods well known in the art. Probes of the present invention, their preparation and/or labeling are elaborated in Sambrook, J. et al., 1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, NY; or Ausubel, F.M. et al., 1989, Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, both of which are incorporated herein by reference in their entirety.

The nucleic acid sequences of the present invention also include the sequence information from any of the nucleic acid sequences of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62. The sequence information can be a segment of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 that uniquely identifies or represents the sequence information of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62. One such segment can be a twenty-mer nucleic acid sequence because the probability that a twenty-mer is fully matched in the human genome is 1 in 300. In the human genome, there are three billion base pairs in one set of chromosomes. Because 4<sup>20</sup> possible twenty-mers exist, there are 300 times more twenty-mers than there are base pairs in a set of human chromosomes. Using the same analysis, the probability for a seventeen-mer to be fully matched in the human genome is approximately 1 in 5. When these segments are used in arrays for expression studies, fifteen-mer segments can be used. The probability that the fifteen-mer is fully matched in the expressed sequences is also approximately one in five because expressed sequences comprise less than approximately 5% of the entire genome sequence.

Similarly, when using sequence information for detecting a single mismatch, a segment can be a twenty-five mer. The probability that the twenty-five mer would appear in a human genome with a single mismatch is calculated by multiplying the probability for a full match (1-4<sup>25</sup>) times the increased probability for mismatch at each nucleotide position (3 x 25). The probability that an eighteen mer with a single mismatch can be detected in an array for expression studies is approximately one in five. The probability that a twenty-mer with a single mismatch can be detected in a human genome is approximately one in five.

The term "open reading frame," ORF, means a series of nucleotide triplets coding for amino acids without any termination codons and is a sequence translatable into protein.

The terms "operably linked" or "operably associated" refer to functionally related nucleic acid sequences. For example, a promoter is operably associated or operably linked with a coding sequence if the promoter controls the transcription of the coding sequence.

While operably linked nucleic acid sequences can be contiguous and in the same reading frame, certain genetic elements e.g. repressor genes are not contiguously linked to the coding sequence but still control transcription/translation of the coding sequence.

The term "pluripotent" refers to the capability of a cell to differentiate into a number of differentiated cell types that are present in an adult organism. A pluripotent cell is restricted in its differentiation capability in comparison to a totipotent cell.

The terms "polypeptide" or "peptide" or "amino acid sequence" refer to an oligopeptide, peptide, polypeptide or protein sequence or fragment thereof and to naturally occurring or synthetic molecules. A polypeptide "fragment," "portion," or "segment" is a stretch of amino acid residues of at least about 5 amino acids, preferably at least about 7 amino acids, more preferably at least about 9 amino acids and most preferably at least about 17 or more amino acids. The peptide preferably is not greater than about 200 amino acids, more preferably less than 150 amino acids and most preferably less than 100 amino acids.

Preferably the peptide is from about 5 to about 200 amino acids. To be active, any

polypeptide must have sufficient length to display biological and/or immunological activity.

The term "naturally occurring polypeptide" refers to polypeptides produced by cells that have not been genetically engineered and specifically contemplates various polypeptides arising from post-translational modifications of the polypeptide including, but not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation.

The term "translated protein coding portion" means a sequence which encodes for the full length protein which may include any leader sequence or a processing sequence.

The term "mature protein coding sequence" refers to a sequence which encodes a peptide or protein without any leader/signal sequence. The "mature protein portion" refers to that portion of the protein without the leader/signal sequence. The peptide may have the leader sequences removed during processing in the cell or the protein may have been produced synthetically or using a polynucleotide only encoding for the mature protein coding sequence. It is contemplated that the mature protein portion may or may not include the initial methionine residue. The initial methionine is often removed during processing of the peptide.

The term "derivative" refers to polypeptides chemically modified by such techniques as ubiquitination, labeling (e.g., with radionuclides or various enzymes), covalent polymer attachment such as pegylation (derivatization with polyethylene glycol) and insertion or substitution by chemical synthesis of amino acids such as ornithine, which do not normally occur in human proteins.

The term "variant" (or "analog") refers to any polypeptide differing from naturally occurring polypeptides by amino acid insertions, deletions, and substitutions, created using, e.g., recombinant DNA techniques. Guidance in determining which amino acid residues may be replaced, added or deleted without abolishing activities of interest, may be found by comparing the sequence of the particular polypeptide with that of homologous peptides and minimizing the number of amino acid sequence changes made in regions of high homology (conserved regions) or by replacing amino acids with consensus sequence.

Alternatively, recombinant variants encoding these same or similar polypeptides may be synthesized or selected by making use of the "redundancy" in the genetic code. Various codon substitutions, such as the silent changes which produce various restriction sites, may be introduced to optimize cloning into a plasmid or viral vector or expression in a particular prokaryotic or eukaryotic system. Mutations in the polynucleotide sequence may be reflected in the polypeptide or domains of other peptides added to the polypeptide to modify the properties of any part of the polypeptide, to change characteristics such as ligand-binding affinities, interchain affinities, or degradation/turnover rate.

Preferably, amino acid "substitutions" are the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, i.e., conservative amino acid replacements. "Conservative" amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid. "Insertions" or "deletions" are preferably in the range of about 1 to 20 amino acids, more preferably 1 to 10 amino acids. The variation allowed may be experimentally determined by systematically making insertions, deletions, or substitutions of amino acids in a polypeptide molecule using recombinant DNA techniques and assaying the resulting recombinant variants for activity.

Alternatively, where alteration of function is desired, insertions, deletions or non-conservative alterations can be engineered to produce altered polypeptides. Such alterations can, for example, alter one or more of the biological functions or biochemical characteristics of the polypeptides of the invention. For example, such alterations may change polypeptide characteristics such as ligand-binding affinities, interchain affinities, or degradation/turnover

rate. Further, such alterations can be selected so as to generate polypeptides that are better suited for expression, scale up and the like in the host cells chosen for expression. For example, cysteine residues can be deleted or substituted with another amino acid residue in order to eliminate disulfide bridges.

5 The terms "purified" or "substantially purified" as used herein denotes that the indicated nucleic acid or polypeptide is present in the substantial absence of other biological macromolecules, e.g., polynucleotides, proteins, and the like. In one embodiment, the polynucleotide or polypeptide is purified such that it constitutes at least 95 % by weight, more preferably at least 99 % by weight, of the indicated biological macromolecules present (but water, buffers, and other small molecules, especially molecules having a molecular weight of 10 less than 1000 daltons, can be present).

The term "isolated" as used herein refers to a nucleic acid or polypeptide separated from at least one other component (e.g., nucleic acid or polypeptide) present with the nucleic acid or polypeptide in its natural source. In one embodiment, the nucleic acid or polypeptide is found in the presence of (if anything) only a solvent, buffer, ion, or other components 15 normally present in a solution of the same. The terms "isolated" and "purified" do not encompass nucleic acids or polypeptides present in their natural source.

The term "recombinant," when used herein to refer to a polypeptide or protein, means that a polypeptide or protein is derived from recombinant (e.g., microbial, insect, or mammalian) expression systems. "Microbial" refers to recombinant polypeptides or proteins made in bacterial or fungal (e.g., yeast) expression systems. As a product, "recombinant microbial" defines a polypeptide or protein essentially free of native endogenous substances and unaccompanied by associated native glycosylation. Polypeptides or proteins expressed in most bacterial cultures, e.g., *E. coli*, will be free of glycosylation modifications; polypeptides 25 or proteins expressed in yeast will have a glycosylation pattern in general different from those expressed in mammalian cells.

The term "recombinant expression vehicle or vector" refers to a plasmid or phage or virus or vector, for expressing a polypeptide from a DNA (RNA) sequence. An expression vehicle can comprise a transcriptional unit comprising an assembly of (1) a genetic element or elements having a regulatory role in gene expression, for example, promoters or enhancers, (2) a structural or coding sequence which is transcribed into mRNA and translated into protein, and (3) appropriate transcription initiation and termination sequences. Structural units intended for use in yeast or eukaryotic expression systems preferably include a leader sequence enabling extracellular secretion of translated protein by a host cell. Alternatively, where recombinant

protein is expressed without a leader or transport sequence, it may include an amino terminal methionine residue. This residue may or may not be subsequently cleaved from the expressed recombinant protein to provide a final product.

The term "recombinant expression system" means host cells which have stably integrated a recombinant transcriptional unit into chromosomal DNA or carry the recombinant transcriptional unit extrachromosomally. Recombinant expression systems as defined herein will express heterologous polypeptides or proteins upon induction of the regulatory elements linked to the DNA segment or synthetic gene to be expressed. This term also means host cells which have stably integrated a recombinant genetic element or elements having a regulatory role in gene expression, for example, promoters or enhancers. Recombinant expression systems as defined herein will express polypeptides or proteins endogenous to the cell upon induction of the regulatory elements linked to the endogenous DNA segment or gene to be expressed. The cells can be prokaryotic or eukaryotic.

The term "secreted" includes a protein that is transported across or through a membrane, including transport as a result of signal sequences in its amino acid sequence when it is expressed in a suitable host cell. "Secreted" proteins include without limitation proteins secreted wholly (e.g., soluble proteins) or partially (e.g., receptors) from the cell in which they are expressed. "Secreted" proteins also include without limitation proteins that are transported across the membrane of the endoplasmic reticulum. "Secreted" proteins are also intended to include proteins containing non-typical signal sequences (e.g. Interleukin-1 Beta, see Krasney, P.A. and Young, P.R. (1992) Cytokine 4(2):134 -143) and factors released from damaged cells (e.g. Interleukin-1 Receptor Antagonist, see Arend, W.P. et. al. (1998) Annu. Rev. Immunol. 16:27-55)

Where desired, an expression vector may be designed to contain a "signal or leader sequence" which will direct the polypeptide through the membrane of a cell. Such a sequence may be naturally present on the polypeptides of the present invention or provided from heterologous protein sources by recombinant DNA techniques.

The term "stringent" is used to refer to conditions that are commonly understood in the art as stringent. Stringent conditions can include highly stringent conditions (i.e., hybridization to filter-bound DNA in 0.5 M NaHPO<sub>4</sub>, 7 % sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1X SSC/0.1 % SDS at 68°C), and moderately stringent conditions (i.e., washing in 0.2X SSC/0.1 % SDS at 42°C). Other exemplary hybridization conditions are described herein in the examples.



In instances of hybridization of deoxyoligonucleotides, additional exemplary stringent hybridization conditions include washing in 6X SSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligonucleotides), 48°C (for 17-base oligonucleotides), 55°C (for 20-base oligonucleotides), and 60°C (for 23-base oligonucleotides).

As used herein, "substantially equivalent" can refer both to nucleotide and amino acid sequences, for example a mutant sequence, that varies from a reference sequence by one or more substitutions, deletions, or additions, the net effect of which does not result in an adverse functional dissimilarity between the reference and subject sequences. Typically, such a substantially equivalent sequence varies from one of those listed herein by no more than about 35% (*i.e.*, the number of individual residue substitutions, additions, and/or deletions in a substantially equivalent sequence, as compared to the corresponding reference sequence, divided by the total number of residues in the substantially equivalent sequence is about 0.35 or less). Such a sequence is said to have 65% sequence identity to the listed sequence. In one embodiment, a substantially equivalent, *e.g.*, mutant, sequence of the invention varies from a listed sequence by no more than 30% (70% sequence identity); in a variation of this embodiment, by no more than 25% (75% sequence identity); and in a further variation of this embodiment, by no more than 20% (80% sequence identity) and in a further variation of this embodiment, by no more than 10% (90% sequence identity) and in a further variation of this embodiment, by no more than 5% (95% sequence identity). Substantially equivalent, *e.g.*, mutant, amino acid sequences according to the invention preferably have at least 80% sequence identity with a listed amino acid sequence, more preferably at least 90% sequence identity, most preferably at least 95% identity. Substantially equivalent nucleotide sequence of the invention can have lower percent sequence identities, taking into account, for example, the redundancy or degeneracy of the genetic code. Preferably, nucleotide sequence has at least about 65% identity, more preferably at least about 75% identity, and most preferably at least about 95% identity. For the purposes of the present invention, sequences having substantially equivalent biological activity and substantially equivalent expression characteristics are considered substantially equivalent. For the purposes of determining equivalence, truncation of the mature sequence (*e.g.*, via a mutation which creates a spurious stop codon) should be disregarded. Sequence identity may be determined, *e.g.*, using the Jotun Hein method (Hein, J. (1990) *Methods Enzymol.* 183:626-645). Identity between sequences can also be determined by other methods known in the art, *e.g.* by varying hybridization conditions.

The term "totipotent" refers to the capability of a cell to differentiate into all of the cell types of an adult organism.

The term "transformation" means introducing DNA into a suitable host cell so that the DNA is replicable, either as an extrachromosomal element, or by chromosomal integration.

The term "transfection" refers to the taking up of an expression vector by a suitable host cell, whether or not any coding sequences are in fact expressed. The term "infection" refers to the introduction of nucleic acids into a suitable host cell by use of a virus or viral vector.

As used herein, an "uptake modulating fragment," UMF, means a series of nucleotides which mediate the uptake of a linked DNA fragment into a cell. UMFs can be readily identified using known UMFs as a target sequence or target motif with the computer-based systems described below. The presence and activity of a UMF can be confirmed by attaching the suspected UMF to a marker sequence. The resulting nucleic acid molecule is then incubated with an appropriate host under appropriate conditions and the uptake of the marker sequence is determined. As described above, a UMF will increase the frequency of uptake of a linked marker sequence.

Each of the above terms is meant to encompass all that is described for each, unless the context dictates otherwise.

## 5.2 NUCLEIC ACIDS OF THE INVENTION

The invention is based on the discovery of a novel secreted GPCR-like polypeptide, the polynucleotides encoding the GPCR-like polypeptide and the use of these compositions for the diagnosis, treatment or prevention of cancers and other immunological disorders.

The isolated polynucleotides of the invention include, but are not limited to a

polynucleotide comprising any of the nucleotide sequences of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62; a fragment of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62; a polynucleotide comprising the full length protein coding sequence of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 (for example SEQ ID NO: 4, 13, 20, 29, 36, or 42); and a

polynucleotide comprising the nucleotide sequence encoding the mature protein coding sequence of the polynucleotides of any one of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62. The polynucleotides of the present invention also include, but are not limited to, a polynucleotide that hybridizes under stringent conditions to

(a) the complement of any of the nucleotides sequences of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62; (b) a polynucleotide encoding any one of the

polypeptides of SEQ ID NO: 4, 6-9, 13, 15-16, 20, 22-25, 29, 31-32, 36, 38-40, 42, 44-47, 61, or 63; (c) a polynucleotide which is an allelic variant of any polynucleotides recited above; (d) a polynucleotide which encodes a species homolog of any of the proteins recited above; or (e) a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of the polypeptides of SEQ ID NO: 4, 6-9, 13, 15-16, 20, 22-25, 29, 31-32, 36, 38-40, 42, 44-47, 61, or 63. Preferably the polynucleotides include a polynucleotide comprising the sequence set forth in nucleotides 1-1351 of SEQ ID NO: 19 or nucleotides 271-1351 of SEQ ID NO: 19. Preferably the polynucleotides include a polynucleotide comprising the sequence set forth in nucleotides 1-1668 of SEQ ID NO: 28, nucleotides 52-1668 of SEQ ID NO: 28, or nucleotides 2845-3993 of SEQ ID NO: 28. Domains of interest may depend on the nature of the encoded polypeptide; e.g., domains in receptor-like polypeptides include ligand-binding, extracellular, transmembrane, or cytoplasmic domains, or combinations thereof; domains in immunoglobulin-like proteins include the variable immunoglobulin-like domains; domains in enzyme-like polypeptides include catalytic and substrate binding domains; and domains in ligand polypeptides include receptor-binding domains.

The polynucleotides of the invention include naturally occurring or wholly or partially synthetic DNA, e.g., cDNA and genomic DNA, and RNA, e.g., mRNA. The polynucleotides may include all of the coding region of the cDNA or may represent a portion of the coding region of the cDNA.

The present invention also provides genes corresponding to the cDNA sequences disclosed herein. The corresponding genes can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include the preparation of probes or primers from the disclosed sequence information for identification and/or amplification of genes in appropriate genomic libraries or other sources of genomic materials. Further 5' and 3' sequence can be obtained using methods known in the art. For example, full length cDNA or genomic DNA that corresponds to any of the polynucleotides of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 can be obtained by screening appropriate cDNA or genomic DNA libraries under suitable hybridization conditions using any of the polynucleotides of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 or a portion thereof as a probe. Alternatively, the polynucleotides of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 may be used as the basis for suitable primer(s) that allow identification and/or amplification of genes in appropriate genomic DNA or cDNA libraries.

The nucleic acid sequences of the invention can be assembled from ESTs and sequences (including cDNA and genomic sequences) obtained from one or more public databases, such as dbEST, gbpi, and UniGene. The EST sequences can provide identifying sequence information, representative fragment or segment information, or novel segment information for the full-length gene.

The polynucleotides of the invention also provide polynucleotides including nucleotide sequences that are substantially equivalent to the polynucleotides recited above.

Polynucleotides according to the invention can have, e.g., at least about 65%, at least about 70%, at least about 75%, at least about 80%, more typically at least about 90%, and even more typically at least about 95%, sequence identity to a polynucleotide recited above.

Included within the scope of the nucleic acid sequences of the invention are nucleic acid sequence fragments that hybridize under stringent conditions to any of the nucleotide sequences of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62, or complements thereof, which fragment is greater than about 5 nucleotides, preferably 7 nucleotides, more preferably greater than 9 nucleotides and most preferably greater than 17 nucleotides. Fragments of, e.g. 15, 17, or 20 nucleotides or more that are selective for (i.e. specifically hybridize to any one of the polynucleotides of the invention) are contemplated. Probes capable of specifically hybridizing to a polynucleotide can differentiate polynucleotide sequences of the invention from other polynucleotide sequences in the same family of genes or can differentiate human genes from genes of other species, and are preferably based on unique nucleotide sequences.

The sequences falling within the scope of the present invention are not limited to these specific sequences, but also include allelic and species variations thereof. Allelic and species variations can be routinely determined by comparing the sequence provided in SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62, a representative fragment thereof, or a nucleotide sequence at least 90% identical, preferably 95% identical, to SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 with a sequence from another isolate of the same species. Furthermore, to accommodate codon variability, the invention includes nucleic acid molecules coding for the same amino acid sequences as do the specific ORFs disclosed herein. In other words, in the coding region of an ORF, substitution of one codon for another codon that encodes the same amino acid is expressly contemplated.

The nearest neighbor result for the nucleic acids of the present invention, including SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62, can be obtained by searching a database using an algorithm or a program. Preferably, a BLAST which stands for

Basic Local Alignment Search Tool is used to search for local sequence alignments (Altschul, S.F. J Mol. Evol. 36:290-300 (1993) and Altschul S.F. et al. J. Mol. Biol. 21:403-410 (1990))

Species homologs (or orthologs) of the disclosed polynucleotides and proteins are also provided by the present invention. Species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source from the desired species.

The invention also encompasses allelic variants of the disclosed polynucleotides or proteins; that is, naturally-occurring alternative forms of the isolated polynucleotide which also encode proteins which are identical, homologous or related to that encoded by the polynucleotides.

The nucleic acid sequences of the invention are further directed to sequences which encode variants of the described nucleic acids. These amino acid sequence variants may be prepared by methods known in the art by introducing appropriate nucleotide changes into a native or variant polynucleotide. There are two variables in the construction of amino acid sequence variants: the location of the mutation and the nature of the mutation. Nucleic acids encoding the amino acid sequence variants are preferably constructed by mutating the polynucleotide to encode an amino acid sequence that does not occur in nature. These nucleic acid alterations can be made at sites that differ in the nucleic acids from different species (variable positions) or in highly conserved regions (constant regions). Sites at such locations

will typically be modified in series, *e.g.*, by substituting first with conservative choices (*e.g.*, hydrophobic amino acid to a different hydrophobic amino acid) and then with more distant choices (*e.g.*, hydrophobic amino acid to a charged amino acid), and then deletions or insertions may be made at the target site. Amino acid sequence deletions generally range from about 1 to 30 residues, preferably about 1 to 10 residues, and are typically contiguous. Amino acid insertions include amino- and/or carboxyl-terminal fusions ranging in length from one to one hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Intrasequence insertions may range generally from about 1 to 10 amino residues, preferably from 1 to 5 residues. Examples of terminal insertions include the heterologous signal sequences necessary for secretion or for intracellular targeting in different host cells and sequences such as FLAG or poly-histidine sequences useful for purifying the expressed protein.

In a preferred method, polynucleotides encoding the novel amino acid sequences are changed via site-directed mutagenesis. This method uses oligonucleotide sequences to alter a polynucleotide to encode the desired amino acid variant, as well as sufficient adjacent

nucleotides on both sides of the changed amino acid to form a stable duplex on either side of the site being changed. In general, the techniques of site-directed mutagenesis are well known to those of skill in the art and this technique is exemplified by publications such as, Edelman et al., *DNA* 2:183 (1983). A versatile and efficient method for producing site-specific changes in a polynucleotide sequence was published by Zoller and Smith, *Nucleic Acids Res.* 10:6487-6500 (1982). PCR may also be used to create amino acid sequence variants of the novel nucleic acids. When small amounts of template DNA are used as starting material, primer(s) that differs slightly in sequence from the corresponding region in the template DNA can generate the desired amino acid variant. PCR amplification results in a population of product DNA fragments that differ from the polynucleotide template encoding the polypeptide at the position specified by the primer. The product DNA fragments replace the corresponding region in the plasmid and this gives a polynucleotide encoding the desired amino acid variant.

A further technique for generating amino acid variants is the cassette mutagenesis technique described in Wells et al., *Gene* 34:315 (1985); and other mutagenesis techniques well known in the art, such as, for example, the techniques in Sambrook et al., *supra*, and *Current Protocols in Molecular Biology*, Ausubel et al. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be used in the practice of the invention for the cloning and expression of these novel nucleic acids. Such DNA sequences include those which are capable of hybridizing to the appropriate novel nucleic acid sequence under stringent conditions.

Polynucleotides encoding preferred polypeptide truncations of the invention can be used to generate polynucleotides encoding chimeric or fusion proteins comprising one or more domains of the invention and heterologous protein sequences.

The polynucleotides of the invention additionally include the complement of any of the polynucleotides recited above. The polynucleotide can be DNA (genomic, cDNA, amplified, or synthetic) or RNA. Methods and algorithms for obtaining such polynucleotides are well known to those of skill in the art and can include, for example, methods for determining hybridization conditions that can routinely isolate polynucleotides of the desired sequence identities.

In accordance with the invention, polynucleotide sequences comprising the mature protein coding sequences corresponding to any one of SEQ ID NO: 4, 6-9, 13, 15-16, 20, 22-25, 29, 31-32, 36, 38-40, 42, 44-47, 61, or 63, or functional equivalents thereof, may be used to generate recombinant DNA molecules that direct the expression of that nucleic acid, or a

functional equivalent thereof, in appropriate host cells. Also included are the cDNA inserts of any of the clones identified herein.

A polynucleotide according to the invention can be joined to any of a variety of other nucleotide sequences by well-established recombinant DNA techniques (see Sambrook J et al. (1989) Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, NY).

Useful nucleotide sequences for joining to polynucleotides include an assortment of vectors, e.g., plasmids, cosmids, lambda phage derivatives, phagemids, and the like, that are well known in the art. Accordingly, the invention also provides a vector including a polynucleotide of the invention and a host cell containing the polynucleotide. In general, the vector contains an origin of replication functional in at least one organism, convenient restriction endonuclease sites, and a selectable marker for the host cell. Vectors according to the invention include expression vectors, replication vectors, probe generation vectors, and sequencing vectors. A host cell according to the invention can be a prokaryotic or eukaryotic cell and can be a unicellular organism or part of a multicellular organism.

The present invention further provides recombinant constructs comprising a nucleic acid having any of the nucleotide sequences of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 or a fragment thereof or any other polynucleotides of the invention. In one embodiment, the recombinant constructs of the present invention comprise a vector, such as a plasmid or viral vector, into which a nucleic acid having any of the nucleotide sequences of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 or a fragment thereof is inserted, in a forward or reverse orientation. In the case of a vector comprising one of the ORFs of the present invention, the vector may further comprise regulatory sequences, including for example, a promoter, operably linked to the ORF. Large numbers of suitable vectors and promoters are known to those of skill in the art and are commercially available for generating the recombinant constructs of the present invention. The following vectors are provided by way of example. Bacterial: pBs, phagescript, PsiX174, pBluescript SK, pBs KS, pNH8a, pNH16a, pNH18a, pNH46a (Stratagene); pTrc99A, pKK223-3, pKK233-3, pDR540, pRIT5 (Pharmacia). Eukaryotic: pWLeo, pSV2cat, pOG44, PXT1, pSG (Stratagene) pSVK3, pBPV, pMSG, pSVL (Pharmacia).

The isolated polynucleotide of the invention may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman et al., *Nucleic Acids Res.* 19, 4485-4490 (1991), in order to produce the protein recombinantly. Many suitable expression control sequences are known in the art. General methods of

expressing recombinant proteins are also known and are exemplified in R. Kaufman, *Methods in Enzymology* 185, 537-566 (1990). As defined herein "operably linked" means that the isolated polynucleotide of the invention and an expression control sequence are situated within a vector or cell in such a way that the protein is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

Promoter regions can be selected from any desired gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers. Two appropriate vectors are pKK232-8 and pCM7. Particular named bacterial promoters include lacI, lacZ, T3, T7, gpt, lambda PR, and trc. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art.

Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, e.g., the ampicillin resistance gene of *E. coli* and *S. cerevisiae* TRP1 gene, and a promoter derived from a highly expressed gene to direct transcription of a downstream structural sequence. Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), a-factor, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein into the periplasmic space or extracellular medium. Optionally, the heterologous sequence can encode a fusion protein including an amino terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product. Useful expression vectors for bacterial use are constructed by inserting a structural DNA sequence encoding a desired protein together with suitable translation initiation and termination signals in operable reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transformation include *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium* and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, although others may also be employed as a matter of choice.

As a representative but non-limiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3

(Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM 1 (Promega Biotech, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed. Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is induced or derepressed by appropriate means (*e.g.*, temperature shift or chemical induction) and cells are cultured for an additional period. Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Polynucleotides of the invention can also be used to induce immune responses. For example, as described in Fan et al., *Nat. Biotech.* 17:870-872 (1999), incorporated herein by reference, nucleic acid sequences encoding a polypeptide may be used to generate antibodies against the encoded polypeptide following topical administration of naked plasmid DNA or following injection, and preferably intramuscular injection of the DNA. The nucleic acid sequences are preferably inserted in a recombinant expression vector and may be in the form of naked DNA.

### 15 5.3 ANTISENSE

Another aspect of the invention pertains to isolated antisense nucleic acid molecules that are hybridizable to or complementary to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62, or fragments, analogs or derivatives thereof. An "antisense" nucleic acid comprises a nucleotide sequence that is complementary to a "sense" nucleic acid encoding a protein, *e.g.*, complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. In specific aspects, antisense nucleic acid molecules are provided that comprise a sequence complementary to at least about 10, 25, 50, 100, 250 or 500 nucleotides or an entire coding strand, or to only a portion thereof. Nucleic acid molecules encoding fragments, homologs, derivatives and analogs of a protein of any of SEQ ID NO: 4, 6-9, 13, 15-16, 20, 22-25, 29, 31-32, 36, 38-40, 42, 44-47, 61, or 63 or antisense nucleic acids complementary to a nucleic acid sequence of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 are additionally provided.

In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence of the invention. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues. In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence of the invention. The term

"noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions).

Given the coding strand sequences encoding a nucleic acid disclosed herein (*e.g.*, SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62, antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick or Hoogsteen base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of a mRNA, but more preferably is an oligonucleotide that is antisense to only a portion of the coding or noncoding region of a mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of a mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis or enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (*e.g.*, an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, *e.g.*, phosphorothioate derivatives and acridine substituted nucleotides can be used.

Examples of modified nucleotides that can be used to generate the antisense nucleic acid include: 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxynucleic acid (v), wybutosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxynucleic acid methyl ester, uracil-5-oxynucleic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)<sup>w</sup>, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (*i.e.*, RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a protein according to the invention to thereby inhibit expression of the protein, *e.g.*, by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule that binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention includes direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, *e.g.*, by linking the antisense nucleic acid molecules to peptides or antibodies that bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an  $\alpha$ -anomeric nucleic acid molecule. An  $\alpha$ -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual -units, the strands run parallel to each other (Gaultier *et al.* (1987) *Nucleic Acids Res* 15: 6625-6641). The antisense nucleic acid molecule can also comprise a 2'-O-methylribonucleotide (Inoue *et al.* (1987) *Nucleic Acids Res* 15: 6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.* (1987) *FEBS Lett* 215: 327-330).

#### 5.4 RIBOZYMES AND PNA MOIETIES

In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity that are capable of cleaving a single-stranded nucleic acid, such as a mRNA, to which they have a complementary region. Thus, ribozymes (*e.g.*, hammerhead ribozymes (described in Haselhoff and Gertach (1988) *Nature* 334:585-591)) can be used to catalytically cleave a mRNA transcripts to thereby inhibit translation of a mRNA. A ribozyme having specificity for a nucleic acid of the invention can be designed based upon the nucleotide sequence of a DNA disclosed herein (*i.e.*, SEQ ID NO:SEQ ID NO: 1-3; 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62). For

example, a derivative of a Tetrahymena L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a SECX-encoding mRNA. See, *e.g.*, Cech *et al.* U.S. Pat. No. 4,987,071, and Cech *et al.* U.S. Pat. No. 5,116,742. Alternatively, mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, *e.g.*, Bartel *et al.*, (1993) *Science* 261:1411-1418.

Alternatively, gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region (*e.g.*, promoter and/or enhancers) to form triple helical structures that prevent transcription of the gene in target cells. See generally, Helene. (1991) *Anticancer Drug Des.* 6: 569-84; Helene. *et al.* (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher (1992) *Bioassays* 14: 807-15.

In various embodiments, the nucleic acids of the invention can be modified at the base moiety, sugar moiety or phosphate backbone to improve, *e.g.*, the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acids can be modified to generate peptide nucleic acids (see Hyrup *et al.* (1996) *Bioorg Med Chem* 4: 5-23). As used herein, the terms "peptide nucleic acids" or "PNAs" refer to nucleic acid mimics, *e.g.*, DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described in Hyrup *et al.* (1996) above; Perry-O'Keefe *et al.* (1996) *PNAS* 93: 14670-675.

PNAs of the invention can be used in therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, *e.g.*, inducing transcription or translation arrest or inhibiting replication. PNAs of the invention can also be used, *e.g.*, in the analysis of single base pair mutations in a gene by, *e.g.*, PNA directed PCR clamping; as artificial restriction enzymes when used in combination with other enzymes, *e.g.*, S1 nucleases (Hyrup B. (1996) above); or as probes or primers for DNA sequence and hybridization (Hyrup *et al.* (1996), above; Perry-O'Keefe (1996), above).

In another embodiment, PNAs of the invention can be modified, *e.g.*, to enhance their stability or cellular uptake, by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras can be generated that may

combine the advantageous properties of PNA and DNA. Such chimeras allow DNA recognition enzymes, *e.g.*, RNase H and DNA polymerases, to interact with the DNA portion while the PNA portion would provide high binding affinity and specificity. PNA-DNA chimeras can be linked using linkers of appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and orientation (Hyrup (1996) above). The synthesis of PNA-DNA chimeras can be performed as described in Hyrup (1996) above and Finn *et al.* (1996) *Nucl Acids Res* 24: 3357-63. For example, a DNA chain can be synthesized on a solid support using standard phosphoramidite coupling chemistry, and modified nucleoside analogs, *e.g.*, 5'-(4-methoxytrityl)amino-5'-deoxy-thymidine phosphoramidite, can be used between the PNA and the 5' end of DNA (Mag *et al.* (1989) *Nucl Acid Res* 17: 5973-88). PNA monomers are then coupled in a stepwise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment (Finn *et al.* (1996) above). Alternatively, chimeric molecules can be synthesized with a 5' DNA segment and a 3' PNA segment. See, Petersen *et al.* (1975) *Bioorg Med Chem Lett* 5: 1119-1124.

In other embodiments, the oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, *e.g.*, Leisinger *et al.*, 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556; Lemaire *et al.*, 1987, *Proc. Natl. Acad. Sci.* 84:648-652; PCT Publication No. W088/09810) or the blood-brain barrier (see, *e.g.*, PCT Publication No. W089/10134). In addition, oligonucleotides can be modified with hybridization triggered cleavage agents (See, *e.g.*, Krol *et al.*, 1988, *BioTechniques* 6:958-976) or intercalating agents. (See, *e.g.*, Zon, 1988, *Pharm. Res.* 5: 539-549). To this end, the oligonucleotide may be conjugated to another molecule, *e.g.*, a peptide, a hybridization triggered cross-linking agent, a transport agent, a hybridization-triggered cleavage agent, etc.

## 5.5 HOSTS

The present invention further provides host cells genetically engineered to contain the polynucleotides of the invention. For example, such host cells may contain nucleic acids of the invention introduced into the host cell using known transformation, transfection or infection methods. The present invention still further provides host cells genetically engineered to express the polynucleotides of the invention, wherein such polynucleotides are in operative association with a regulatory sequence heterologous to the host cell which drives expression of the polynucleotides in the cell.

Knowledge of GPCR-like DNA sequences allows for modification of cells to permit, or increase, expression of GPCR-like polypeptide. Cells can be modified (*e.g.*, by homologous recombination) to provide increased GPCR-like polypeptide expression by replacing, in whole or in part, the naturally occurring GPCR-like promoter with all or part of a heterologous promoter so that the cells GPCR-like polypeptide is expressed at higher levels. The heterologous promoter is inserted in such a manner that it is operatively linked to GPCR-like encoding sequences. See, for example, PCT International Publication No. WO94/12650, PCT International Publication No. WO92/20808, and PCT International Publication No. WO91/09955. It is also contemplated that, in addition to heterologous promoter DNA, amplifiable marker DNA (*e.g.*, *ada*, *dhfr*, and the multifunctional CAD gene which encodes carbamyl phosphate synthase, aspartate transcarbamylase, and dihydroorotase) and/or intron DNA may be inserted along with the heterologous promoter DNA. If linked to the GPCR-like coding sequence, amplification of the marker DNA by standard selection methods results in co-amplification of the GPCR-like coding sequences in the cells.

The host cell can be a higher eukaryotic host cell, such as a mammalian cell, a lower eukaryotic host cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the recombinant construct into the host cell can be effected by calcium phosphate transfection, DEAE, dextran-mediated transfection, or electroporation (Davis, L. *et al.*, *Basic Methods in Molecular Biology* (1986)). The host cells containing one of the polynucleotides of the invention, can be used in conventional manners to produce the gene product encoded by the isolated fragment (in the case of an ORF) or can be used to produce a heterologous protein under the control of the EMF.

Any host/vector system can be used to express one or more of the ORFs of the present invention. These include, but are not limited to, eukaryotic hosts such as HeLa cells, CV-1 cell, COS cells, 293 cells, and Sf9 cells, as well as prokaryotic host such as *E. coli* and *B. subtilis*. The most preferred cells are those which do not normally express the particular polypeptide or protein or which expresses the polypeptide or protein at low natural level. Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook, *et al.*, in *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor, New York (1989), the disclosure of which is hereby incorporated by reference.

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman, Cell 23:175 (1981). Other cell lines capable of expressing a compatible vector are, for example, the C127, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK or Jurkat cells. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early promoter, enhancer, splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements. Recombinant polypeptides and proteins produced in bacterial culture are usually isolated by initial extraction from cell pellets, followed by one or more salting-out, aqueous ion exchange or size exclusion chromatography steps. Protein refolding steps can be used, as necessary, in completing configuration of the mature protein. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps. Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

Alternatively, it may be possible to produce the protein in lower eukaryotes such as yeast or insects or in prokaryotes such as bacteria. Potentially suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Potentially suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins. If the protein is made in yeast or bacteria, it may be necessary to modify the protein produced therein, for example by phosphorylation or glycosylation of the appropriate sites, in order to obtain the functional protein. Such covalent attachments may be accomplished using known chemical or enzymatic methods.

In another embodiment of the present invention, cells and tissues may be engineered to express an endogenous gene comprising the polynucleotides of the invention under the control of inducible regulatory elements, in which case the regulatory sequences of the endogenous gene may be replaced by homologous recombination. As described herein, gene targeting can be used to replace a gene's existing regulatory region with a regulatory sequence isolated from

a different gene or a novel regulatory sequence synthesized by genetic engineering methods. Such regulatory sequences may be comprised of promoters, enhancers, scaffold-attachment regions, negative regulatory elements, transcriptional initiation sites, regulatory protein binding sites or combinations of said sequences. Alternatively, sequences which affect the structure or stability of the RNA or protein produced may be replaced, removed, added, or otherwise modified by targeting. These sequences include polyadenylation signals, mRNA stability elements, splice sites, leader sequences for enhancing or modifying transport or secretion properties of the protein, or other sequences which alter or improve the function or stability of protein or RNA molecules.

The targeting event may be a simple insertion of the regulatory sequence, placing the gene under the control of the new regulatory sequence, *e.g.*, inserting a new promoter or enhancer or both upstream of a gene. Alternatively, the targeting event may be a simple deletion of a regulatory element, such as the deletion of a tissue-specific negative regulatory element. Alternatively, the targeting event may replace an existing element; for example, a tissue-specific enhancer can be replaced by an enhancer that has broader or different cell-type specificity than the naturally occurring elements. Here, the naturally occurring sequences are deleted and new sequences are added. In all cases, the identification of the targeting event may be facilitated by the use of one or more selectable marker genes that are contiguous with the targeting DNA, allowing for the selection of cells in which the exogenous DNA has integrated into the host cell genome. The identification of the targeting event may also be facilitated by the use of one or more marker genes exhibiting the property of negative selection, such that the negatively selectable marker is linked to the exogenous DNA, but configured such that the negatively selectable marker flanks the targeting sequence, and such that a correct homologous recombination event with sequences in the host cell genome does not result in the stable integration of the negatively selectable marker. Markers useful for this purpose include the Herpes Simplex Virus thymidine kinase (TK) gene or the bacterial xanthine-guanine phosphoribosyl-transferase (*gpt*) gene.

The gene targeting or gene activation techniques which can be used in accordance with this aspect of the invention are more particularly described in U.S. Patent No. 5,272,071 to Chappel; U.S. Patent No. 5,578,461 to Sherwin et al.; International Application No. PCT/US92/09627 (WO93/09222) by Selden et al.; and International Application No. PCT/US90/06436 (WO91/06667) by Skoultschi et al., each of which is incorporated by reference herein in its entirety.



## 5.6 POLYPEPTIDES OF THE INVENTION

The isolated polypeptides of the invention include, but are not limited to, a polypeptide comprising: the amino acid sequence set forth as any one of SEQ ID NO: 4, 6-9, 13, 15-16, 20, 22-25, 29, 31-32, 36, 38-40, 42, 44-47, 61, or 63 or an amino acid sequence encoded by any one of the nucleotide sequences SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 or the corresponding full length or mature protein. Polypeptides of the invention also include polypeptides preferably with biological or immunological activity that are encoded by: (a) a polynucleotide having any one of the nucleotide sequences set forth in SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 or (b) polynucleotides encoding any one of the amino acid sequences set forth as SEQ ID NO: 4, 6-9, 13, 15-16, 20, 22-25, 29, 31-32, 36, 38-40, 42, 44-47, 61, or 63 or (c) polynucleotides that hybridize to the complement of the polynucleotides of either (a) or (b) under stringent hybridization conditions. The invention also provides biologically active or immunologically active variants of any of the amino acid sequences set forth as SEQ ID NO: 4, 6-9, 13, 15-16, 20, 22-25, 29, 31-32, 36, 38-40, 42, 44-47, 61, or 63 or the corresponding full length or mature protein; and "substantial equivalents" thereof (e.g., with at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, typically at least about 95%, more typically at least about 98%, or most typically at least about 99% amino acid identity) that retain biological activity. Polypeptides encoded by allelic variants may have a similar, increased, or decreased activity compared to polypeptides comprising SEQ ID NO: 4, 6-9, 13, 15-16, 20, 22-25, 29, 31-32, 36, 38-40, 42, 44-47, 61, or 63. Preferably the polypeptides include a polypeptide comprising the sequence set forth in amino acid residues 1-360 of SEQ ID NO: 20. Preferably the polypeptides include a polypeptide comprising the sequence set forth in amino acid residues 1-539 of SEQ ID NO: 29 or the sequence set forth in amino acid residues 932-1314 of SEQ ID NO: 29.

Fragments of the proteins of the present invention which are capable of exhibiting biological activity are also encompassed by the present invention. Fragments of the protein may be in linear form or they may be cyclized using known methods, for example, as described in H. U. Saragovi, et al., *Bio/Technology* 10, 773-778 (1992) and in R. S. McDowell, et al., *J. Amer. Chem. Soc.* 114, 9245-9253 (1992), both of which are incorporated herein by reference. Such fragments may be fused to carrier molecules such as immunoglobulins for many purposes, including increasing the valency of protein binding sites.

The present invention also provides both full-length and mature forms (for example, without a signal sequence or precursor sequence) of the disclosed proteins. The protein coding

sequence is identified in the sequence listing by translation of the disclosed nucleotide sequences. The mature form of such protein may be obtained by expression of a full-length polynucleotide in a suitable mammalian cell or other host cell. The sequence of the mature form of the protein is also determinable from the amino acid sequence of the full-length form. Where proteins of the present invention are membrane bound, soluble forms of the proteins are also provided. In such forms, part or all of the regions causing the proteins to be membrane bound are deleted so that the proteins are fully secreted from the cell in which it is expressed.

Protein compositions of the present invention may further comprise an acceptable carrier, such as a hydrophilic, e.g., pharmaceutically acceptable, carrier.

The present invention further provides isolated polypeptides encoded by the nucleic acid fragments of the present invention or by degenerate variants of the nucleic acid fragments of the present invention. By "degenerate variant" is intended nucleotide fragments which differ from a nucleic acid fragment of the present invention (e.g., an ORF) by nucleotide sequence but, due to the degeneracy of the genetic code, encode an identical polypeptide sequence. Preferred nucleic acid fragments of the present invention are the ORFs that encode proteins.

A variety of methodologies known in the art can be utilized to obtain any one of the isolated polypeptides or proteins of the present invention. At the simplest level, the amino acid sequence can be synthesized using commercially available peptide synthesizers. The synthetically-constructed protein sequences, by virtue of sharing primary, secondary or tertiary structural and/or conformational characteristics with proteins may possess biological properties in common therewith, including protein activity. This technique is particularly useful in producing small peptides and fragments of larger polypeptides. Fragments are useful, for example, in generating antibodies against the native polypeptide. Thus, they may be employed as biologically active or immunological substitutes for natural, purified proteins in screening of therapeutic compounds and in immunological processes for the development of antibodies.

The polypeptides and proteins of the present invention can alternatively be purified from cells which have been altered to express the desired polypeptide or protein. As used herein, a cell is said to be altered to express a desired polypeptide or protein when the cell, through genetic manipulation, is made to produce a polypeptide or protein which it normally does not produce or which the cell normally produces at a lower level. One skilled in the art can readily adapt procedures for introducing and expressing either recombinant or synthetic sequences into eukaryotic or prokaryotic cells in order to generate a cell which produces one of the polypeptides or proteins of the present invention.

The invention also relates to methods for producing a polypeptide comprising growing a culture of host cells of the invention in a suitable culture medium, and purifying the protein from the cells or the culture in which the cells are grown. For example, the methods of the invention include a process for producing a polypeptide in which a host cell containing a suitable expression vector that includes a polynucleotide of the invention is cultured under conditions that allow expression of the encoded polypeptide. The polypeptide can be recovered from the culture, conveniently from the culture medium, or from a lysate prepared from the host cells and further purified. Preferred embodiments include those in which the protein produced by such process is a full length or mature form of the protein.

In an alternative method, the polypeptide or protein is purified from bacterial cells which naturally produce the polypeptide or protein. One skilled in the art can readily follow known methods for isolating polypeptides and proteins in order to obtain one of the isolated polypeptides or proteins of the present invention. These include, but are not limited to, immuno-chromatography, HPLC, size-exclusion chromatography, ion-exchange chromatography, and immuno-affinity chromatography. See, e.g., Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag (1994); Sambrook, et al., in *Molecular Cloning: A Laboratory Manual*; Ausubel et al., *Current Protocols in Molecular Biology*. Polypeptide fragments that retain biological/immunological activity include fragments comprising greater than about 100 amino acids, or greater than about 200 amino acids, and fragments that encode specific protein domains.

The purified polypeptides can be used in *in vitro* binding assays which are well known in the art to identify molecules which bind to the polypeptides. These molecules include but are not limited to, for e.g., small molecules, molecules from combinatorial libraries, antibodies or other proteins. The molecules identified in the binding assay are then tested for antagonist or agonist activity in *in vivo* tissue culture or animal models that are well known in the art. In brief, the molecules are titrated into a plurality of cell cultures or animals and then tested for either cell/animal death or prolonged survival of the animal/cells.

In addition, the peptides of the invention or molecules capable of binding to the peptides may be complexed with toxins, e.g., ricin or cholera, or with other compounds that are toxic to cells. The toxin-binding molecule complex is then targeted to a tumor or other cell by the specificity of the binding molecule for SEQ ID NO: 4, 6-9, 13, 15-16, 20, 22-25, 29, 31-32, 36, 38-40, 42, 44-47, 61, or 63.

The protein of the invention may also be expressed as a product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells containing a nucleotide sequence encoding the protein.

The proteins provided herein also include proteins characterized by amino acid sequences similar to those of purified proteins but into which modification are naturally provided or deliberately engineered. For example, modifications, in the peptide or DNA sequence, can be made by those skilled in the art using known techniques. Modifications of interest in the protein sequences may include the alteration, substitution, replacement, insertion or deletion of a selected amino acid residue in the coding sequence. For example, one or more of the cysteine residues may be deleted or replaced with another amino acid to alter the conformation of the molecule. Techniques for such alteration, substitution, replacement, insertion or deletion are well known to those skilled in the art (see, e.g., U.S. Pat. No. 4,518,584). Preferably, such alteration, substitution, replacement, insertion or deletion retains the desired activity of the protein. Regions of the protein that are important for the protein function can be determined by various methods known in the art including the alanine-scanning method which involved systematic substitution of single or strings of amino acids with alanine, followed by testing the resulting alanine-containing variant for biological activity. This type of analysis determines the importance of the substituted amino acid(s) in biological activity. Regions of the protein that are important for protein function may be determined by the eMATRIX program.

Other fragments and derivatives of the sequences of proteins which would be expected to retain protein activity in whole or in part and are useful for screening or other immunological methodologies may also be easily made by those skilled in the art given the disclosures herein. Such modifications are encompassed by the present invention.

The protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, e.g., Invitrogen, San Diego, Calif., U.S.A. (the MaxBat™ kit), and such methods are well known in the art, as described in Summers and Smith, *Texas Agricultural Experiment Station Bulletin No. 1555* (1987), incorporated herein by reference. As used herein, an insect cell capable of expressing a polynucleotide of the present invention is "transformed."

The protein of the invention may be prepared by culturing transformed host cells under culture conditions suitable to express the recombinant protein. The resulting expressed protein

may then be purified from such culture (*i.e.*, from culture medium or cell extracts) using known purification processes, such as gel filtration and ion exchange chromatography. The purification of the protein may also include an affinity column containing agents which will bind to the protein; one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl™ or Cibacrom blue 3GA Sepharose™, one or more steps involving hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or immunoaffinity chromatography.

Alternatively, the protein of the invention may also be expressed in a form which will facilitate purification. For example, it may be expressed as a fusion protein, such as those of maltose binding protein (MBP), glutathione-S-transferase (GST) or thioredoxin (TRX), or as a His tag. Kits for expression and purification of such fusion proteins are commercially available from New England BioLab (Beverly, Mass.), Pharmacia (Piscataway, N.J.) and Invitrogen, respectively. The protein can also be tagged with an epitope and subsequently purified by using a specific antibody directed to such epitope. One such epitope ("FLAG®") is commercially available from Kodak (New Haven, Conn.).

Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, *e.g.*, silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the protein. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a substantially homogeneous isolated recombinant protein. The protein thus purified is substantially free of other mammalian proteins and is defined in accordance with the present invention as an "isolated protein."

The polypeptides of the invention include analogs (variants). The polypeptides of the invention include GPCR-like analogs. This embraces fragments of GPCR-like polypeptide of the invention, as well GPCR-like polypeptides which comprise one or more amino acids deleted, inserted, or substituted. Also, analogs of the GPCR-like polypeptide of the invention embrace fusions of the GPCR-like polypeptides or modifications of the GPCR-like polypeptides, wherein the GPCR-like polypeptide or analog is fused to another moiety or moieties, *e.g.*, targeting moiety or another therapeutic agent. Such analogs may exhibit improved properties such as activity and/or stability. Examples of moieties which may be fused to the GPCR-like polypeptide or an analog include, for example, targeting moieties which provide for the delivery of polypeptide to neurons, *e.g.*, antibodies to central nervous system; or antibodies to receptor and ligands expressed on neuronal cells. Other moieties

which may be fused to GPCR-like polypeptide include therapeutic agents which are used for treatment, for example anti-depressant drugs or other medications for neurological disorders. Also, GPCR-like polypeptides may be fused to neuron growth modulators, and other chemokines for targeted delivery.

## 5.6.1 DETERMINING POLYPEPTIDE AND POLYNUCLEOTIDE IDENTITY AND SIMILARITY

Preferred identity and/or similarity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in computer programs including, but are not limited to, the GCG program package, including GAP (Devereux, J., et al., *Nucleic Acids Research* 12(1):387 (1984); Genetics Computer Group, University of Wisconsin, Madison, WI), BLASTP, BLASTN, BLASTX, FASTA (Altschul, S.F. et al., *J. Molec. Biol.* 215:403-410 (1990), PSI-BLAST (Altschul S.F. et al., *Nucleic Acids Res.* vol. 25, pp. 3389-3402, herein incorporated by reference), the eMatrix software (Wu et al., *J. Comp. Biol.*, vol. 6, pp. 219-235 (1999), herein incorporated by reference), eMotif software (Nevill-Manning et al, *ISMB-97*, vol 4, pp. 202-209, herein incorporated by reference), the GeneAtlas software (Molecular Simulations Inc. (MSI), San Diego, CA) (Sanchez and Sali (1998) *Proc. Natl. Acad. Sci.*, 95, 13597-13602; Kison DH et al. (2000) "Remote homology detection using structural modeling - an evaluation" Submitted; Fischer and Eisenberg (1996) *Protein Sci.* 5, 947-955), Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark), Pfam, which are multiple protein sequence alignment and hidden Markov models of common protein domains (Wang et al (2000) submitted and Bateman et al (2000) *Nucleic Acid Res.* 28, 263-266) and the Kyte-Doolittle hydrophobicity prediction algorithm (*J. Mol.Biol.* 157, pp. 105-31 (1982), incorporated herein by reference). The BLAST programs are publicly available from the National Center for Biotechnology Information (NCBI) and other sources (BLAST Manual, Altschul, S., et al. NCB NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215:403-410 (1990).

## 5.7 CHIMERIC AND FUSION PROTEINS

The invention also provides chimeric or fusion proteins. As used herein, a "chimeric protein" or "fusion protein" comprises a polypeptide of the invention operatively linked to another polypeptide. Within a fusion protein the polypeptide according to the invention can

correspond to all or a portion of a protein according to the invention. In one embodiment, a fusion protein comprises at least one biologically active portion of a protein according to the invention. In another embodiment, a fusion protein comprises at least two biologically active portions of a protein according to the invention. Within the fusion protein, the term "operatively linked" is intended to indicate that the polypeptide(s) according to the invention and the other polypeptide(s) are fused in-frame to each other. The polypeptide can be fused to the N-terminus or C-terminus or in the middle.

For example, in one embodiment a fusion protein comprises a polypeptide according to the invention operably linked to the extracellular domain of a second protein.

In another embodiment, the fusion protein is a GST-fusion protein in which the polypeptide sequences of the invention are fused to the C-terminus of the GST (i.e., glutathione S-transferase) sequences.

In another embodiment, the fusion protein is an immunoglobulin fusion protein in which the polypeptide sequences according to the invention comprise one or more domains fused to sequences derived from a member of the immunoglobulin protein family. The immunoglobulin fusion proteins of the invention can be incorporated into pharmaceutical compositions and administered to a subject to inhibit an interaction between a ligand and a protein of the invention on the surface of a cell, to thereby suppress signal transduction *in vivo*. The immunoglobulin fusion proteins can be used to affect the bioavailability of a cognate ligand. Inhibition of the ligand/protein interaction may be useful therapeutically for both the treatment of proliferative and differentiative disorders, *e.g.*, cancer as well as modulating the invention can be used to bind and to dimerize 2 receptors and thereby transduce an intracellular signal. The immunoglobulin fusion proteins may also be used as immunogens to produce antibodies in a subject, to purify ligands, and in screening assays to identify molecules that inhibit the interaction of a polypeptide of the invention with a ligand.

A chimeric or fusion protein of the invention can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, *e.g.*, by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using

anchor primers that give rise to complementary overhangs between two consecutive gene fragments that can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, Ausubel et al. (eds.) CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST polypeptide): A nucleic acid encoding a polypeptide of the invention can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the protein of the invention.

## 5.8 GENE THERAPY

Mutations in the polynucleotides of the invention gene may result in loss of normal function of the encoded protein. The invention thus provides gene therapy to restore normal activity of the polypeptides of the invention; or to treat disease states involving polypeptides of the invention. Delivery of a functional gene encoding polypeptides of the invention to appropriate cells is effected *ex vivo*, *in situ*, or *in vivo* by use of vectors, and more particularly viral vectors (*e.g.*, adenovirus, adeno-associated virus, or a retrovirus), or *ex vivo* by use of physical DNA transfer methods (*e.g.*, liposomes or chemical treatments). See, for example, Anderson, Nature, supplement to vol. 392, no. 6679, pp.25-20 (1998). For additional reviews of gene therapy technology see Friedmann, Science, 244: 1275-1281 (1989); Verma, Scientific American: 68-84 (1990); and Miller, Nature, 357: 455-460 (1992). Introduction of any one of the nucleotides of the present invention or a gene encoding the polypeptides of the present invention can also be accomplished with extrachromosomal substrates (transient expression) or artificial chromosomes (stable expression). Cells may also be cultured *ex vivo* in the presence of proteins of the present invention in order to proliferate or to produce a desired effect on or activity in such cells. Treated cells can then be introduced *in vivo* for therapeutic purposes. Alternatively, it is contemplated that in other human disease states, preventing the expression of or inhibiting the activity of polypeptides of the invention will be useful in treating the disease states. It is contemplated that antisense therapy or gene therapy could be applied to negatively regulate the expression of polypeptides of the invention.

Other methods inhibiting expression of a protein include the introduction of antisense molecules to the nucleic acids of the present invention, their complements, or their translated RNA sequences, by methods known in the art. Further, the polypeptides of the present invention can be inhibited by using targeted deletion methods, or the insertion of a negative regulatory element such as a silencer, which is tissue specific.

The present invention still further provides cells genetically engineered *in vivo* to express the polynucleotides of the invention, wherein such polynucleotides are in operative association with a regulatory sequence heterologous to the host cell which drives expression of the polynucleotides in the cell. These methods can be used to increase or decrease the expression of the polynucleotides of the present invention.

Knowledge of DNA sequences provided by the invention allows for modification of cells to permit, increase, or decrease, expression of endogenous polypeptide. Cells can be modified (e.g., by homologous recombination) to provide increased polypeptide expression by replacing, in whole or in part, the naturally occurring promoter with all or part of a heterologous promoter so that the cells express the protein at higher levels. The heterologous promoter is inserted in such a manner that it is operatively linked to the desired protein encoding sequences. See, for example, PCT International Publication No. WO 94/12650, PCT International Publication No. WO 92/20808, and PCT International Publication No. WO 91/09955. It is also contemplated that, in addition to heterologous promoter DNA, amplifiable marker DNA (e.g., *ada*, *dhfr*, and the multifunctional CAD gene which encodes carbamyl phosphate synthase, aspartate transcarbamylase, and dihydroorotase) and/or intron DNA may be inserted along with the heterologous promoter DNA. If linked to the desired protein coding sequence, amplification of the marker DNA by standard selection methods results in co-amplification of the desired protein coding sequences in the cells.

In another embodiment of the present invention, cells and tissues may be engineered to express an endogenous gene comprising the polynucleotides of the invention under the control of inducible regulatory elements, in which case the regulatory sequences of the endogenous gene may be replaced by homologous recombination. As described herein, gene targeting can be used to replace a gene's existing regulatory region with a regulatory sequence isolated from a different gene or a novel regulatory sequence synthesized by genetic engineering methods. Such regulatory sequences may be comprised of promoters, enhancers, scaffold-attachment regions, negative regulatory elements, transcriptional initiation sites, regulatory protein binding sites or combinations of said sequences. Alternatively, sequences which affect the structure or stability of the RNA or protein produced may be replaced, removed, added, or otherwise modified by targeting. These sequences include polyadenylation signals, mRNA stability elements, splice sites, leader sequences for enhancing or modifying transport or secretion properties of the protein, or other sequences which alter or improve the function or stability of protein or RNA molecules.

The targeting event may be a simple insertion of the regulatory sequence, placing the gene under the control of the new regulatory sequence, *e.g.*, inserting a new promoter or enhancer or

both upstream of a gene. Alternatively, the targeting event may be a simple deletion of a regulatory element, such as the deletion of a tissue-specific negative regulatory element. Alternatively, the targeting event may replace an existing element; for example, a tissue-specific enhancer can be replaced by an enhancer that has broader or different cell-type specificity than the naturally occurring elements. Here, the naturally occurring sequences are deleted and new sequences are added. In all cases, the identification of the targeting event may be facilitated by the use of one or more selectable marker genes that are contiguous with the targeting DNA, allowing for the selection of cells in which the exogenous DNA has integrated into the cell genome. The identification of the targeting event may also be facilitated by the use of one or more marker genes exhibiting the property of negative selection, such that the negatively selectable marker is linked to the exogenous DNA, but configured such that the negatively selectable marker flanks the targeting sequence, and such that a correct homologous recombination event with sequences in the host cell genome does not result in the stable integration of the negatively selectable marker. Markers useful for this purpose include the Herpes Simplex Virus thymidine kinase (TK) gene or the bacterial xanthine-guanine phosphoribosyl-transferase (*gpt*) gene.

The gene targeting or gene activation techniques which can be used in accordance with this aspect of the invention are more particularly described in U.S. Patent No. 5,272,071 to Chappel; U.S. Patent No. 5,578,461 to Sherwin et al.; International Application No. PCT/US92/09627 (WO93/09222) by Seiden et al.; and International Application No. PCT/US90/06436 (WO91/06667) by Skoultschi et al., each of which is incorporated by reference herein in its entirety.

## 5.9 TRANSGENIC ANIMALS

In preferred methods to determine biological functions of the polypeptides of the invention *in vivo*, one or more genes provided by the invention are either over expressed or inactivated in the germ line of animals using homologous recombination [Capecchi, Science 244:1288-1292 (1989)]. Animals in which the gene is over expressed, under the regulatory control of exogenous or endogenous promoter elements, are known as transgenic animals. Animals in which an endogenous gene has been inactivated by homologous recombination are referred to as "knockout" animals. Knockout animals, preferably non-human mammals, can be prepared as described in U.S. Patent No. 5,557,032, incorporated herein by reference.

Transgenic animals are useful to determine the roles polypeptides of the invention play in biological processes, and preferably in disease states. Transgenic animals are useful as model

systems to identify compounds that modulate lipid metabolism. Transgenic animals, preferably non-human mammals, are produced using methods as described in U.S. Patent No. 5,489,743 and PCT Publication No. WO94/28122, incorporated herein by reference.

Transgenic animals can be prepared wherein all or part of a promoter of the polynucleotides of the invention is either activated or inactivated to alter the level of expression of the polypeptides of the invention. Inactivation can be carried out using homologous recombination methods described above. Activation can be achieved by supplementing or even replacing the homologous promoter to provide for increased protein expression. The homologous promoter can be supplemented by insertion of one or more heterologous enhancer elements known to confer promoter activation in a particular tissue.

The polynucleotides of the present invention also make possible the development, through, e.g., homologous recombination or knock out strategies; of animals that fail to express functional GPCR-like polypeptide or that express a variant of GPCR-like polypeptide. Such animals are useful as models for studying the *in vivo* activities of GPCR-like polypeptide as well as for studying modulators of the GPCR-like polypeptide.

In preferred methods to determine biological functions of the polypeptides of the invention *in vivo*, one or more genes provided by the invention are either over expressed or inactivated in the germ line of animals using homologous recombination [Capecchi, Science 244:1288-1292 (1989)]. Animals in which the gene is over expressed, under the regulatory control of exogenous or endogenous promoter elements, are known as transgenic animals. Animals in which an endogenous gene has been inactivated by homologous recombination are referred to as "knockout" animals. Knockout animals, preferably non-human mammals, can be prepared as described in U.S. Patent No. 5,557,032, incorporated herein by reference.

Transgenic animals are useful to determine the roles polypeptides of the invention play in biological processes, and preferably in disease states. Transgenic animals are useful as model systems to identify compounds that modulate lipid metabolism. Transgenic animals, preferably non-human mammals, are produced using methods as described in U.S. Patent No. 5,489,743 and PCT Publication No. WO94/28122, incorporated herein by reference.

Transgenic animals can be prepared wherein all or part of the polynucleotides of the invention promoter is either activated or inactivated to alter the level of expression of the polypeptides of the invention. Inactivation can be carried out using homologous recombination methods described above. Activation can be achieved by supplementing or even replacing the homologous promoter to provide for increased protein expression. The homologous promoter

can be supplemented by insertion of one or more heterologous enhancer elements known to confer promoter activation in a particular tissue.

## 5.10 USES AND BIOLOGICAL ACTIVITY OF HUMAN GPCR-LIKE POLYPEPTIDE

The polynucleotides and proteins of the present invention are expected to exhibit one or more of the uses or biological activities (including those associated with assays cited herein) identified herein. Uses or activities described for proteins of the present invention may be provided by administration or use of such proteins or of polynucleotides encoding such proteins (such as, for example, in gene therapies or vectors suitable for introduction of DNA).

The mechanism underlying the particular condition or pathology will dictate whether the polypeptides of the invention, the polynucleotides of the invention or modulators (activators or inhibitors) thereof would be beneficial to the subject in need of treatment. Thus, "therapeutic compositions of the invention" include compositions comprising isolated polynucleotides (including recombinant DNA molecules, cloned genes and degenerate variants thereof) or polypeptides of the invention (including full length protein, mature protein and truncations or domains thereof), or compounds and other substances that modulate the overall activity of the target gene products, either at the level of target gene/protein expression or target protein activity. Such modulators include polypeptides, analogs, (variants), including fragments and fusion proteins, antibodies and other binding proteins; chemical compounds that directly or indirectly activate or inhibit the polypeptides of the invention (identified, e.g., via drug screening assays as described herein); antisense polynucleotides and polynucleotides suitable for triple helix formation; and in particular antibodies or other binding partners that specifically recognize one or more epitopes of the polypeptides of the invention.

The polypeptides of the present invention may likewise be involved in cellular activation or in one of the other physiological pathways described herein.

### 5.10.1 RESEARCH USES AND UTILITIES

The polynucleotides provided by the present invention can be used by the research community for various purposes. The polynucleotides can be used to express recombinant protein for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in disease states); as molecular weight markers on gels; as chromosome markers or tags (when labeled) to identify chromosomes or to map related gene positions; to compare with endogenous DNA sequences in patients to identify

potential genetic disorders; as probes to hybridize and thus discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; as a probe to "subtract-out" known sequences in the process of discovering other novel

polynucleotides; for selecting and making oligomers for attachment to a "gene chip" or other support, including for examination of expression patterns; to raise anti-protein antibodies using DNA immunization techniques; and as an antigen to raise anti-DNA antibodies or elicit another immune response. Where the polynucleotide encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, that described in Gyuris et al., Cell 75:791-803 (1993)) to identify polynucleotides encoding the other protein with which binding occurs or to identify inhibitors of the binding interaction.

The polypeptides provided by the present invention can similarly be used in assays to determine biological activity, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the corresponding polypeptide is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Proteins involved in these binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

The polypeptides of the invention are also useful for making antibody substances that are specifically immunoreactive with GPCR-like proteins. Antibodies and portions thereof (e.g., Fab fragments) which bind to the polypeptides of the invention can be used to identify the presence of such polypeptides in a sample. Such determinations are carried out using any suitable immunoassay format, and any polypeptide of the invention that is specifically bound by the antibody can be employed as a positive control.

Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as research products.

Methods for performing the uses listed above are well known to those skilled in the art.

References disclosing such methods include without limitation "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E. F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S. L. and A. R. Kimmel eds., 1987.

### 5.10.2 NUTRITIONAL USES

Polynucleotides and polypeptides of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein or amino acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In such cases the polypeptide or polynucleotide of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the polypeptide or polynucleotide of the invention can be added to the medium in or on which the microorganism is cultured.

Additionally, the polypeptides of the invention can be used as molecular weight markers, and as a food supplement. A polypeptide consisting of SEQ ID NO: 4, for example, has a molecular mass of approximately 93 kDa in its unprocessed and unglycosylated state. Protein food supplements are well known and the formulation of suitable food supplements including polypeptides of the invention is within the level of skill in the food preparation art.

### 5.10.3 CYTOKINE AND CELL PROLIFERATION/DIFFERENTIATION

#### ACTIVITY

A polypeptide of the present invention may exhibit activity relating to cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may induce production of other cytokines in certain cell populations. A

polynucleotide of the invention can encode a polypeptide exhibiting such attributes. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor-dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of therapeutic compositions of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+ (preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e, CMK, HUVETC, and Caco. Therapeutic compositions of the invention can be used in the following:

Assays for T-cell or thymocyte proliferation include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H.

Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-

Interscience (Chapter 3. *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7,

Immunologic studies in Humans); Takai et al., *J. Immunol.* 137:3494-3500, 1986; Bertagnolli et al., *J. Immunol.* 145:1706-1712, 1990; Bertagnolli et al., *Cellular Immunology* 133:327-341, 1991; Bertagnolli, et al., *J. Immunol.* 149:3778-3783, 1992; Bowman et al., *J. Immunol.* 152:1756-1761, 1994.

5 Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: Polyclonal T cell stimulation, Kruisbeek, A. M. and Shevach, E. M. In *Current Protocols in Immunology*. J. E. e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto, 1994; and Measurement of mouse and human interleukin- $\gamma$ , Schreiber, R. D. In *Current Protocols in Immunology*. J. E. e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto, 1994.

10 Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without limitation, those described in: Measurement of Human and Murine Interleukin 2 and Interleukin 4, Bottomly, K., Davis, L. S. and Lipsky, P. E. In *Current Protocols in Immunology*. J. E. e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto, 1991; deVries et al., *J. Exp. Med.* 173:1205-1211, 1991; Moreau et al., *Nature* 336:690-692, 1988; Greenberger et al., *Proc. Natl. Acad. Sci. U.S.A.* 80:2931-2938, 1983; Measurement of mouse and human interleukin 6--Nordan, R. In *Current Protocols in Immunology*. J. E. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto, 1991; Smith et al., *Proc. Natl. Acad. Sci. U.S.A.* 83:1857-1861, 1986; Measurement of human Interleukin 11--Bennett, F., Giannotti, J., Clark, S. C. and Turner, K. J. In *Current Protocols in Immunology*. J. E. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto, 1991; Measurement of mouse and human Interleukin 9-Ciarletta, A., Giannotti, J., Clark, S. C. and Turner, K. J. In *Current Protocols in Immunology*. J. E. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto, 1991.

25 Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors; Chapter 7, Immunologic studies in Humans); Weinberger et al., *Proc. Natl. Acad. Sci. USA* 77:6091-6095, 1980; Weinberger et al., *Eur. J. Immunol.* 11:405-411, 1981; Takai et al., *J. Immunol.* 137:3494-3500, 1986; Takai et al., *J. Immunol.* 140:508-512, 1988.

#### 5.10.4 STEM CELL GROWTH FACTOR ACTIVITY

5 A polypeptide of the present invention may exhibit stem cell growth factor activity and be involved in the proliferation, differentiation and survival of pluripotent and totipotent stem cells including primordial germ cells, embryonic stem cells, hematopoietic stem cells and/or germ line stem cells. Administration of the polypeptide of the invention to stem cells *in vivo* or *ex vivo* may maintain and expand cell populations in a totipotent or pluripotential state which would be useful for re-engineering damaged or diseased tissues, transplantation, manufacture of bio-pharmaceuticals and the development of bio-sensors. The ability to produce large quantities of human cells has important working applications for the production of human proteins which currently must be obtained from non-human sources or donors, implantation of cells to treat diseases such as Parkinson's, Alzheimer's and other neurodegenerative diseases; tissues for grafting such as bone marrow, skin, cartilage, tendons, bone, muscle (including cardiac muscle), blood vessels, cornea, neural cells, gastrointestinal cells and others; and organs for transplantation such as kidney, liver, pancreas (including islet cells), heart and lung.

15 It is contemplated that multiple different exogenous growth factors and/or cytokines may be administered in combination with the polypeptide of the invention to achieve the desired effect, including any of the growth factors listed herein, other stem cell maintenance factors, and specifically including stem cell factor (SCF), leukemia inhibitory factor (LIF), Flt-3 ligand (Flt-3L), any of the interleukins, recombinant soluble IL-6 receptor fused to IL-6, macrophage inflammatory protein 1-alpha (MIP-1-alpha), G-CSF, GM-CSF, thrombopoietin (TPO), platelet factor 4 (PF-4), platelet-derived growth factor (PDGF), neural growth factors and basic fibroblast growth factor (bFGF).

25 Since totipotent stem cells can give rise to virtually any mature cell type, expansion of these cells in culture will facilitate the production of large quantities of mature cells. Techniques for culturing stem cells are known in the art and administration of polypeptides of the invention, optionally with other growth factors and/or cytokines, is expected to enhance the survival and proliferation of the stem cell populations. This can be accomplished by direct administration of the polypeptide of the invention to the culture medium. Alternatively, stroma cells transfected with a polynucleotide that encodes for the polypeptide of the invention can be used as a feeder layer for the stem cell populations in culture or *in vivo*. Stromal support cells



for feeder layers may include embryonic bone marrow fibroblasts, bone marrow stromal cells, fetal liver cells, or cultured embryonic fibroblasts (see U.S. Patent No. 5,690,926).

Stem cells themselves can be transfected with a polynucleotide of the invention to induce autocrine expression of the polypeptide of the invention. This will allow for generation of undifferentiated totipotent/pluripotent stem cell lines that are useful as is or that can then be differentiated into the desired mature cell types. These stable cell lines can also serve as a source of undifferentiated totipotent/pluripotent mRNA to create cDNA libraries and templates for polymerase chain reaction experiments. These studies would allow for the isolation and identification of differentially expressed genes in stem cell populations that regulate stem cell proliferation and/or maintenance.

Expansion and maintenance of totipotent stem cell populations will be useful in the treatment of many pathological conditions. For example, polypeptides of the present invention may be used to manipulate stem cells in culture to give rise to neuroepithelial cells that can be used to augment or replace cells damaged by illness, autoimmune disease, accidental damage or genetic disorders. The polypeptide of the invention may be useful for inducing the proliferation of neural cells and for the regeneration of nerve and brain tissue, i.e. for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders which involve degeneration, death or trauma to neural cells or nerve tissue. In addition, the expanded stem cell populations can also be genetically altered for gene therapy purposes and to decrease host rejection of replacement tissues after grafting or implantation.

Expression of the polypeptide of the invention and its effect on stem cells can also be manipulated to achieve controlled differentiation of the stem cells into more differentiated cell types. A broadly applicable method of obtaining pure populations of a specific differentiated cell type from undifferentiated stem cell populations involves the use of a cell-type specific promoter driving a selectable marker. The selectable marker allows only cells of the desired type to survive. For example, stem cells can be induced to differentiate into cardiomyocytes (Wobus et al., *Differentiation*, 48: 173-182, (1991); Klug et al., *J. Clin. Invest.*, 98(1): 216-224, (1998)) or skeletal muscle cells (Browder, L. W. In: *Principles of Tissue Engineering eds.* Lanza et al., Academic Press (1997)). Alternatively, directed differentiation of stem cells can be accomplished by culturing the stem cells in the presence of a differentiation factor such as retinoic acid and an antagonist of the polypeptide of the invention which would inhibit the effects of endogenous stem cell factor activity and allow differentiation to proceed.

*In vitro* cultures of stem cells can be used to determine if the polypeptide of the invention exhibits stem cell growth factor activity. Stem cells are isolated from any one of various cell sources (including hematopoietic stem cells and embryonic stem cells) and cultured on a feeder layer, as described by Thompson et al. *Proc. Natl. Acad. Sci., U.S.A.*, 92: 7844-7848 (1995), in the presence of the polypeptide of the invention alone or in combination with other growth factors or cytokines. The ability of the polypeptide of the invention to induce stem cells proliferation is determined by colony formation on semi-solid support e.g. as described by Bernstein et al., *Blood*, 77: 2316-2321 (1991).

#### 5.10.5 HEMATOPOIESIS REGULATING ACTIVITY

A polypeptide of the present invention may be involved in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell disorders. Even marginal biological activity in support of colony forming cells or of factor-dependent cell lines indicates involvement in regulating hematopoiesis, e.g. in supporting the growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (i.e., traditional CSF activity) useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and generally for use in place of or complementary to platelet transfusions; and/or in supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either *in-vivo* or *ex-vivo* (i.e., in conjunction with bone marrow transplantation or with peripheral progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated for gene therapy.

Therapeutic compositions of the invention can be used in the following:  
Suitable assays for proliferation and differentiation of various hematopoietic lines, including those assays cited above.

Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. *Cellular Biology* 15:141-151, 1995; Keller et al., *Molecular and Cellular Biology* 13:473-486, 1993; McClanahan et al., *Blood* 81:2903-2915, 1993.

Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M. G. In *Culture of Hematopoietic Cells*. R. I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, N.Y. 1994;

10 Hirayama et al., *Proc. Natl. Acad. Sci. USA* 89:5907-5911, 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I. K. and Briddell, R. A. In *Culture of Hematopoietic Cells*. R. I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, N.Y. 1994; Neben et al., *Experimental Hematology* 22:353-359, 1994;

15 Cobblestone area forming cell assay, Ploemacher, R. E. In *Culture of Hematopoietic Cells*. R. I. Freshney, et al. eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, N.Y. 1994; Long term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and Allen, T. In *Culture of Hematopoietic Cells*. R. I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, N.Y. 1994; Long term culture initiating cell assay, Sutherland, H. J. In *Culture of Hematopoietic Cells*. R. I. Freshney, et al. eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, N.Y. 1994.

#### 5.10.6 TISSUE GROWTH ACTIVITY

A polypeptide of the present invention also may be involved in bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as in wound healing and tissue repair and replacement, and in healing of burns, incisions and ulcers.

A polypeptide of the present invention which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Compositions of a polypeptide, antibody, binding partner, or other modulator of the invention may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. De novo bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery.

A polypeptide of this invention may also be involved in attracting bone-forming cells, stimulating growth of bone-forming cells, or inducing differentiation of progenitors of bone-forming cells. Treatment of osteoporosis, osteoarthritis, bone degenerative disorders, or periodontal disease, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes may also be possible using the composition of the invention.

Another category of tissue regeneration activity that may involve the polypeptide of the present invention is tendon/ligament formation. Induction of tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. De novo tendon/ligament-like tissue formation induced by a composition of the present invention contributes to the repair of congenital, trauma induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions of the present invention may provide an environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The compositions of the present invention may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, i.e. for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders, which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a composition may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated in accordance with the present invention include mechanical and traumatic

disorders, such as spinal cord disorders, head trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a composition of the invention.

5 Compositions of the invention may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

10 Compositions of the present invention may also be involved in the generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac) and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues.

Part of the desired effects may be by inhibition or modulation of fibrotic scarring may allow normal tissue to regenerate. A polypeptide of the present invention may also exhibit angiogenic activity.

15 A composition of the present invention may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage.

A composition of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

20 Therapeutic compositions of the invention can be used in the following:

Assays for tissue generation activity include, without limitation, those described in: International Patent Publication No. WO95/16035 (bone, cartilage, tendon); International Patent Publication No. WO95/05846 (nerve, neuronal); International Patent Publication No. WO91/07491 (skin, endothelium).

25 Assays for wound healing activity include, without limitation, those described in: Winter, Epidermal Wound Healing, pp. 71-112 (Maibach, H. I. and Rovee, D. T., eds.), Year Book Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, J. Invest. Dermatol 71:382-84 (1978).

### 30 5.10.7 IMMUNE FUNCTION STIMULATING OR SUPPRESSING ACTIVITY

A polypeptide of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A polynucleotide of the invention can encode a polypeptide exhibiting such activities. A protein may be useful in the treatment of various immune deficiencies and disorders

(including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of the present invention, including infections by HIV, hepatitis viruses, herpes viruses, mycobacteria, Leishmania spp., malaria spp. and various fungal infections such as candidiasis. Of course, in this regard, proteins of the present invention may also be useful where a boost to the immune system generally may be desirable, i.e., in the treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example, connective tissue disease, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease. Such a protein (or antagonists thereof, including antibodies) of the present invention may also be useful in the treatment of allergic reactions and conditions (e.g., anaphylaxis, serum sickness, drug reactions, food allergies, insect venom allergies, mastocytosis, allergic rhinitis, hypersensitivity pneumonitis, urticaria, angioedema, eczema, atopic dermatitis, allergic contact dermatitis, erythema multiforme, Stevens-Johnson syndrome, allergic conjunctivitis, atopic keratoconjunctivitis, vernal keratoconjunctivitis, giant papillary conjunctivitis and contact allergies), such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions, in which immune suppression is desired (including, for example, organ transplantation), may also be treatable using a protein (or antagonists thereof) of the present invention. The therapeutic effects of the polypeptides or antagonists thereof on allergic reactions can be evaluated by *in vivo* animals models such as the cumulative contact enhancement test (Lastbom et al., Toxicology 125: 59-66, 1998), skin prick test (Hoffmann et al., Allergy 54: 446-54, 1999), guinea pig skin sensitization test (Vohr et al., Arch. Toxicol. 73: 501-9), and murine local lymph node assay (Kimber et al., J. Toxicol. Environ. Health 53: 563-79).

Using the proteins of the invention it may also be possible to modulate immune responses, in a number of ways. Down regulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The functions of activated T cells may be inhibited by suppressing T cell responses

or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or energy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions (such as, for example, B7)), e.g., preventing high level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys the transplant. The administration of a therapeutic composition of the invention may prevent cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, a lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term tolerance by B lymphocyte antigen-blocking reagents may avoid the necessity of repeated administration of these blocking reagents. To achieve sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens.

The efficacy of particular therapeutic compositions in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans. Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins *in vivo* as described in Lenschow et al., *Science* 257:789-792 (1992) and Turka et al., *Proc. Natl. Acad. Sci. USA*, 89:11102-11105 (1992). In addition, murine models of GVHD (see Paul et al., *Fundamental Immunology*, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of therapeutic compositions of the invention on the development of that disease.

Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and which promote the production of cytokines and autoantibodies involved in the pathology of the diseases. Preventing the activation of

autoreactive T cells may reduce or eliminate disease symptoms. Administration of reagents which block stimulation of T cells can be used to inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which may be involved in the disease process. Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from the disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythematosus in MRL/lpr/lpr mice or NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and murine experimental myasthenia gravis (see Paul et al., *Fundamental Immunology*, Raven Press, New York, 1989, pp. 840-856).

Upregulation of an antigen function (e.g., a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy. Upregulation of immune responses may be in the form of enhancing an existing immune response or eliciting an initial immune response. For example, enhancing an immune response may be useful in cases of viral infection, including systemic viral diseases such as influenza, the common cold, and encephalitis.

Alternatively, anti-viral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells *in vitro* with viral antigen-pulsed APCs either expressing a peptide of the present invention or together with a stimulatory form of a soluble peptide of the present invention and reintroducing the *in vitro* activated T cells into the patient. Another method of enhancing anti-viral immune responses would be to isolate infected cells from a patient, transfect them with a nucleic acid encoding a protein of the present invention as described herein such that the cells express all or a portion of the protein on their surface, and reintroduce the transfected cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to, and thereby activate, T cells *in vivo*.

A polypeptide of the present invention may provide the necessary stimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which lack MHC class I or MHC class II molecules, or which fail to reexpress sufficient amounts of MHC class I or MHC class II molecules, can be transfected with nucleic acid encoding all or a portion of (e.g., a cytoplasmic-domain truncated portion) of an MHC class I alpha chain protein and  $\beta$  microglobulin protein or an MHC class II alpha chain protein and an MHC class II beta chain protein to thereby express MHC class I or MHC

class II proteins on the cell surface. Expression of the appropriate class I or class II MHC in conjunction with a peptide having the activity of a B lymphocyte antigen (e.g., B7-1, B7-2, B7-3) induces a T cell mediated immune response against the transfected tumor cell.

Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA encoding a peptide having the activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bowman et al., J. Virology 61:1992-1998; Bertagnoli et al., Cellular Immunology 133:327-341, 1991; Brown et al., J. Immunol. 153:3079-3092, 1994.

Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, J. Immunol. 144:3028-3033, 1990; and Assays for B cell function: In vitro antibody production, Mond, J. J. and Brunswick, M. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.

Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnoli et al., J. Immunol. 149:3778-3783, 1992.

Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al., Journal of Experimental Medicine 173:549-559, 1991; Macatonia et al., Journal of Immunology 154:5071-5079, 1995; Porgador et al., Journal of Experimental Medicine 182:255-260, 1995; Nair et al., Journal of Virology 67:4062-4069, 1993; Huang et al., Science 264:961-965, 1994; Macatonia et al., Journal of Experimental Medicine 169:1255-1264, 1989; Bhardwaj et al., Journal of Clinical Investigation 94:797-807, 1994; and Inaba et al., Journal of Experimental Medicine 172:631-640, 1990.

Assays for lymphocyte survival/apoptosis (which will identify, among others, proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in: Darzynkiewicz et al., Cytometry 13:795-808, 1992; Gorczyca et al., Leukemia 7:659-670, 1993; Gorczyca et al., Cancer Research 53:1945-1951, 1993; Itoh et al., Cell 66:233-243, 1991; Zacharchuk, Journal of Immunology 145:4037-4045, 1990; Zamai et al., Cytometry 14:891-897, 1993; Gorczyca et al., International Journal of Oncology 1:639-648, 1992.

Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., Blood 84:111-117, 1994; Fine et al., Cellular Immunology 155:111-122, 1994; Galy et al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad. Sci. USA 88:7548-7551, 1991.

#### 5.10.8 ACTIVIN/INHIBIN ACTIVITY

A polypeptide of the present invention may also exhibit activin- or inhibin-related activities. A polynucleotide of the invention may encode a polypeptide exhibiting such characteristics. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins are characterized by their ability to stimulate the release of follicle stimulating hormone (FSH). Thus, a polypeptide of the present invention, alone or in heterodimers with a member of the inhibin family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the polypeptide of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example,

U.S. Pat. No. 4,798,885. A polypeptide of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as, but not limited to, cows, sheep and pigs.

5 The activity of a polypeptide of the invention may, among other means, be measured by the following methods.

Assays for activin/inhibin activity include, without limitation, those described in: Vale et al., *Endocrinology* 91:562-572, 1972; Ling et al., *Nature* 321:779-782, 1986; Vale et al., *Nature* 321:776-779, 1986; Mason et al., *Nature* 318:659-663, 1985; Forage et al., *Proc. Natl. Acad. Sci. USA* 83:3091-3095, 1986.

#### 5.10.9 CHEMOTACTIC/CHEMOKINETIC ACTIVITY

A polypeptide of the present invention may be involved in chemotactic or chemokinetic activity for mammalian cells, including, for example, monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells. A polynucleotide of the invention can encode a polypeptide exhibiting such attributes. Chemotactic and chemokinetic receptor activation can be used to mobilize or attract a desired cell population to a desired site of action. Chemotactic or chemokinetic compositions (e.g. proteins, antibodies, binding partners, or modulators of the invention) provide particular advantages in treatment of wounds and other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

Therapeutic compositions of the invention can be used in the following:

30 Assays for chemotactic activity (which will identify proteins that induce or prevent chemotaxis) consist of assays that measure the ability of a protein to induce the migration of cells across a membrane as well as the ability of a protein to induce the adhesion of one cell population to another cell population. Suitable assays for movement and adhesion include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan,

A. M. Kruisbeek, D. H. Margules, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 6.12, Measurement of alpha and beta Chemokines 6.12.1-6.12.28; Taub et al. *J. Clin. Invest.* 95:1370-1376, 1995; Lind et al. *APMIS* 103:140-146, 1995; Muller et al. *Eur. J. Immunol.* 25:1744-1748; Gruber et al. *J. of Immunol.* 152:5860-5867, 1994; Johnston et al. *J. of Immunol.* 153:1762-1768, 1994.

#### 5.10.10 HEMOSTATIC AND THROMBOLYTIC ACTIVITY

A polypeptide of the invention may also be involved in hemostasis or thrombolysis or thrombosis. A polynucleotide of the invention can encode a polypeptide exhibiting such attributes. Compositions may be useful in treatment of various coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A composition of the invention may also be useful for dissolving or inhibiting formation of thromboses and for treatment and prevention of conditions resulting therefrom (such as, for example, infarction of cardiac and central nervous system vessels (e.g., stroke).

Therapeutic compositions of the invention can be used in the following:

Assay for hemostatic and thrombolytic activity include, without limitation, those described in: Linet et al., *J. Clin. Pharmacol.* 26:131-140, 1986; Burdick et al., *Thrombosis Res.* 45:413-419, 1987; Humphrey et al., *Fibrinolysis* 5:71-79 (1991); Schaub, *Prostaglandins* 35:467-474, 1988.

#### 5.10.11 CANCER DIAGNOSIS AND THERAPY

Polypeptides of the invention may be involved in cancer cell generation, proliferation or metastasis. Detection of the presence or amount of polynucleotides or polypeptides of the invention may be useful for the diagnosis and/or prognosis of one or more types of cancer. For example, the presence or increased expression of a polynucleotide/polypeptide of the invention may indicate a hereditary risk of cancer, a precancerous condition, or an ongoing malignancy. Conversely, a defect in the gene or absence of the polypeptide may be associated with a cancer condition. Identification of single nucleotide polymorphisms associated with cancer or a predisposition to cancer may also be useful for diagnosis or prognosis.

Cancer treatments promote tumor regression by inhibiting tumor cell proliferation, inhibiting angiogenesis (growth of new blood vessels that is necessary to support tumor growth) and/or prohibiting metastasis by reducing tumor cell motility or invasiveness.

Therapeutic compositions of the invention may be effective in adult and pediatric oncology

including in solid phase tumors/malignancies, locally advanced tumors, human soft tissue sarcomas, metastatic cancer, including lymphatic metastases, blood cell malignancies including multiple myeloma, acute and chronic leukemias, and lymphomas, head and neck cancers including mouth cancer, larynx cancer and thyroid cancer, lung cancers including small cell carcinoma and non-small cell cancers, breast cancers including small cell carcinoma and ductal carcinoma, gastrointestinal cancers including esophageal cancer, stomach cancer, colon cancer, colorectal cancer and polyps associated with colorectal neoplasia, pancreatic cancers, liver cancer, urologic cancers including bladder cancer and prostate cancer, malignancies of the female genital tract including ovarian carcinoma, uterine (including endometrial) cancers, and solid tumor in the ovarian follicle, kidney cancers including renal cell carcinoma, brain cancers including intrinsic brain tumors, neuroblastoma, astrocytic brain tumors, gliomas, metastatic tumor cell invasion in the central nervous system, bone cancers including osteomas, skin cancers including malignant melanoma, tumor progression of human skin keratinocytes, squamous cell carcinoma, basal cell carcinoma, hemangiopericytoma and Kaposi's sarcoma.

Polypeptides, polynucleotides, or modulators of polypeptides of the invention (including inhibitors and stimulators of the biological activity of the polypeptide of the invention) may be administered to treat cancer. Therapeutic compositions can be administered in therapeutically effective dosages alone or in combination with adjuvant cancer therapy such as surgery, chemotherapy, radiotherapy, thermotherapy, and laser therapy, and may provide a beneficial effect, e.g. reducing tumor size, slowing rate of tumor growth, inhibiting metastasis, or otherwise improving overall clinical condition, without necessarily eradicating the cancer.

The composition can also be administered in therapeutically effective amounts as a portion of an anti-cancer cocktail. An anti-cancer cocktail is a mixture of the polypeptide or modulator of the invention with one or more anti-cancer drugs in addition to a pharmaceutically acceptable carrier for delivery. The use of anti-cancer cocktails as a cancer treatment is routine. Anti-cancer drugs that are well known in the art and can be used as a treatment in combination with the polypeptide or modulator of the invention include:

Actinomycin D, Aminoglutethimide, Asparaginase, Bleomycin, Busulfan, Carboplatin, Carmustine, Chlorambucil, Cisplatin (cis-DDP), Cyclophosphamide, Cytarabine HCl (Cytosine arabinoside), Dacarbazine, Dactinomycin, Daunorubicin HCl, Doxorubicin HCl, Estramustine phosphate sodium, Etoposide (V16-213), Floxuridine, 5-Fluorouracil (5-Fu), Flutamide, Hydroxyurea (hydroxycarbamide), Ifosfamide, Interferon Alpha-2a, Interferon Alpha-2b, Leuprolide acetate (LHRH-releasing factor analog), Lomustine, Mechlorethamine HCl (nitrogen mustard), Melphalan, Mercaptopurine, Mesna, Methotrexate (MTX),

Mitomycin, Mitoxantrone HCl, Octreotide, Plicamycin, Procarbazine HCl, Streptozocin, Tamoxifen citrate, Thioguanine, Thiotepa, Vinblastine sulfate, Vincristine sulfate, Amsacrine, Azacitidine, Hexamethylmelamine, Interleukin-2, Mitoguanzone, Pentostatin, Semustine, Teniposide, and Vindesine sulfate.

In addition, therapeutic compositions of the invention may be used for prophylactic treatment of cancer. There are hereditary conditions and/or environmental situations (e.g. exposure to carcinogens) known in the art that predispose an individual to developing cancers. Under these circumstances, it may be beneficial to treat these individuals with therapeutically effective doses of the polypeptide of the invention to reduce the risk of developing cancers.

*In vitro* models can be used to determine the effective doses of the polypeptide of the invention as a potential cancer treatment. These *in vitro* models include proliferation assays of cultured tumor cells, growth of cultured tumor cells in soft agar (see Freshney, (1987) Culture of Animal Cells: A Manual of Basic Technique, Wiley-Liss, New York, NY Ch 18 and Ch 21), tumor systems in nude mice as described in Giovanella et al., J. Natl. Can. Inst., 52: 921-30 (1974), mobility and invasive potential of tumor cells in Boyden Chamber assays as described in Pilkington et al., Anticancer Res., 17: 4107-9 (1997), and angiogenesis assays such as induction of vascularization of the chick chorioallantoic membrane or induction of vascular endothelial cell migration as described in Ribatta et al., Intl. J. Dev. Biol., 40: 1189-97 (1999) and Li et al., Clin. Exp. Metastasis, 17:423-9 (1999), respectively. Suitable tumor cell lines are available, e.g. from American Type Tissue Culture Collection catalogs.

#### 5.10.12 RECEPTOR/LIGAND ACTIVITY

A polypeptide of the present invention may also demonstrate activity as receptor, receptor ligand or inhibitor or agonist of receptor/ligand interactions. A polynucleotide of the invention can encode a polypeptide exhibiting such characteristics. Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses. Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction. A protein of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions.

The activity of a polypeptide of the invention may, among other means, be measured by the following methods:

Suitable assays for receptor-ligand activity include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley- Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static conditions 7.28.1- 7.28.22), Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 1987; Bierer et al., J. Exp. Med. 168:1145-1156, 1988; Rosenstein et al., J. Exp. Med. 169:149-160 1989; Stoltzenberg et al., J. Immunol. Methods 175:59-68, 1994; Stitt et al., Cell 80:661-670, 1995.

By way of example, the polypeptides of the invention may be used as a receptor for a ligand(s) thereby transmitting the biological activity of that ligand(s). Ligands may be identified through binding assays, affinity chromatography, dihybrid screening assays, BIAcore assays, gel overlay assays, or other methods known in the art.

Studies characterizing drugs or proteins as agonist or antagonist or partial agonists or a partial antagonist require the use of other proteins as competing ligands. The polypeptides of the present invention or ligand(s) thereof may be labeled by being coupled to radioisotopes, colorimetric molecules or a toxin molecules by conventional methods. ("Guide to Protein Purification" Murray P. Deutscher (ed) Methods in Enzymology Vol. 182 (1990) Academic Press, Inc. San Diego). Examples of radioisotopes include, but are not limited to, tritium and carbon-14. Examples of colorimetric molecules include, but are not limited to, fluorescent molecules such as fluorescamine, or rhodamine or other colorimetric molecules. Examples of toxins include, but are not limited, to ricin.

### 5.10.13 DRUG SCREENING

This invention is particularly useful for screening chemical compounds by using the novel polypeptides or binding fragments thereof in any of a variety of drug screening techniques. The polypeptides or fragments employed in such a test may either be free in solution, affixed to a solid support, borne on a cell surface or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the polypeptide or a fragment thereof. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may measure, for example, the formation of complexes between polypeptides of the invention or fragments and

the agent being tested or examine the diminution in complex formation between the novel polypeptides and an appropriate cell line, which are well known in the art.

Sources for test compounds that may be screened for ability to bind to or modulate (i.e., increase or decrease) the activity of polypeptides of the invention include (1) inorganic and organic chemical libraries, (2) natural product libraries, and (3) combinatorial libraries comprised of either random or mimetic peptides, oligonucleotides or organic molecules.

Chemical libraries may be readily synthesized or purchased from a number of commercial sources, and may include structural analogs of known compounds or compounds that are identified as "hits" or "leads" via natural product screening.

The sources of natural product libraries are microorganisms (including bacteria and fungi), animals, plants or other vegetation, or marine organisms, and libraries of mixtures for screening may be created by: (1) fermentation and extraction of broths from soil, plant or marine microorganisms or (2) extraction of the organisms themselves. Natural product libraries include polyketides, non-ribosomal peptides, and (non-naturally occurring) variants thereof. For a review, see *Science* 282:63-68 (1998).

Combinatorial libraries are composed of large numbers of peptides, oligonucleotides or organic compounds and can be readily prepared by traditional automated synthesis methods, PCR, cloning or proprietary synthetic methods. Of particular interest are peptide and oligonucleotide combinatorial libraries. Still other libraries of interest include peptide, protein, peptidomimetic, multiparallel synthetic collection, recombinatorial, and polypeptide libraries. For a review of combinatorial chemistry and libraries created therefrom, see Myers, *Curr. Opin. Biotechnol.* 8:701-707 (1997). For reviews and examples of peptidomimetic libraries, see Al-Obeidi et al., *Mol. Biotechnol.* 9(3):205-23 (1998); Hruby et al., *Curr Opin Chem Biol.* 1(1):114-19 (1997); Dörner et al., *Bioorg Med Chem.* 4(5):709-15 (1996) (alkylated dipeptides).

Identification of modulators through use of the various libraries described herein permits modification of the candidate "hit" (or "lead") to optimize the capacity of the "hit" to bind a polypeptide of the invention. The molecules identified in the binding assay are then tested for antagonist or agonist activity in *in vivo* tissue culture or animal models that are well known in the art. In brief, the molecules are titrated into a plurality of cell cultures or animals and then tested for either cell/animal death or prolonged survival of the animal/cells.

The binding molecules thus identified may be complexed with toxins, e.g., ricin or cholera, or with other compounds that are toxic to cells such as radioisotopes. The toxin-binding molecule complex is then targeted to a tumor or other cell by the specificity of the



binding molecule for a polypeptide of the invention. Alternatively, the binding molecules may be complexed with imaging agents for targeting and imaging purposes.

#### 5.10.14 ASSAY FOR RECEPTOR ACTIVITY

5 The invention also provides methods to detect specific binding of a polypeptide e.g. a ligand or a receptor. The art provides numerous assays particularly useful for identifying previously unknown binding partners for receptor polypeptides of the invention. For example, expression cloning using mammalian or bacterial cells, or dihybrid screening assays can be used to identify polynucleotides encoding binding partners. As another example, affinity chromatography with the appropriate immobilized polypeptide of the invention can be used to isolate polypeptides that recognize and bind polypeptides of the invention. There are a number of different libraries used for the identification of compounds, and in particular small molecule, that modulate (*i.e.*, increase or decrease) biological activity of a polypeptide of the invention. Ligands for receptor polypeptides of the invention can also be identified by adding exogenous ligands, or cocktails of ligands to two cells populations that are genetically identical except for the expression of the receptor of the invention: one cell population expresses the receptor of the invention whereas the other does not. The response of the two cell populations to the addition of ligands(s) are then compared. Alternatively, an expression library can be co-expressed with the polypeptide of the invention in cells and assayed for an autocrine response to identify potential ligand(s). As still another example, BIAcore assays, gel overlay assays, or other methods known in the art can be used to identify binding partner polypeptides, including, (1) organic and inorganic chemical libraries, (2) natural product libraries, and (3) combinatorial libraries comprised of random peptides, oligonucleotides or organic molecules.

25 The role of downstream intracellular signaling molecules in the signaling cascade of the polypeptide of the invention can be determined. For example, a chimeric protein in which the cytoplasmic domain of the polypeptide of the invention is fused to the extracellular portion of a protein, whose ligand has been identified, is produced in a host cell. The cell is then incubated with the ligand specific for the extracellular portion of the chimeric protein, thereby activating the chimeric receptor. Known downstream proteins involved in intracellular signaling can then be assayed for expected modifications *i.e.* phosphorylation. Other methods known to those in the art can also be used to identify signaling molecules involved in receptor activity.

#### 5.10.15 ANTI-INFLAMMATORY ACTIVITY

Compositions of the present invention may also exhibit anti-inflammatory activity. The anti-inflammatory activity may be achieved by providing a stimulus to cells involved in the inflammatory response, by inhibiting or promoting cell-cell interactions (such as, for example, cell adhesion), by inhibiting or promoting chemolaxis of cells involved in the inflammatory process, inhibiting or promoting cell extravasation, or by stimulating or suppressing production of other factors which more directly inhibit or promote an inflammatory response.

5 Compositions with such activities can be used to treat inflammatory conditions including chronic or acute conditions, including without limitation intimation associated with infection (such as septic shock, sepsis or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine-induced lung injury, inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF or IL-1. Compositions of the invention may also be useful to treat anaphylaxis and hypersensitivity to an antigenic substance or material. Compositions of this invention may be utilized to prevent or treat conditions such as, but not limited to, sepsis, acute pancreatitis, endotoxin shock, cytokine induced shock, rheumatoid arthritis, chronic inflammatory arthritis, pancreatic cell damage from diabetes mellitus type 1, graft versus host disease, inflammatory bowel disease, inflammation associated with pulmonary disease, other autoimmune disease or inflammatory disease, an antiproliferative agent such as for acute or chronic myelogenous leukemia or in the prevention of premature labor secondary to intrauterine infections.

#### 5.10.16 LEUKEMIAS

15 Leukemias and related disorders may be treated or prevented by administration of a therapeutic that promotes or inhibits function of the polynucleotides and/or polypeptides of the invention. Such leukemias and related disorders include but are not limited to acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia, myeloblastic, promyelocytic, myelomonocytic, monocytic, erythroleukemia, chronic leukemia, chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia (for a review of such disorders, see Fishman et al., 1985, Medicine, 2d Ed., J.B. Lippincott Co., Philadelphia).

#### 5.10.17 NERVOUS SYSTEM DISORDERS

30 Nervous system disorders, involving cell types which can be tested for efficacy of intervention with compounds that modulate the activity of the polynucleotides and/or polypeptides of the invention, and which can be treated upon thus observing an indication of

therapeutic utility, include but are not limited to nervous system injuries, and diseases or disorders which result in either a disconnection of axons, a diminution or degeneration, of neurons, or demyelination. Nervous system lesions which may be treated in a patient (including human and non-human mammalian patients) according to the invention include but are not limited to the following lesions of either the central (including spinal cord, brain) or peripheral nervous systems:

(i) traumatic lesions, including lesions caused by physical injury or associated with surgery, for example, lesions which sever a portion of the nervous system, or compression injuries;

(ii) ischemic lesions, in which a lack of oxygen in a portion of the nervous system results in neuronal injury or death, including cerebral infarction or ischemia, or spinal cord infarction or ischemia;

(iii) infectious lesions, in which a portion of the nervous system is destroyed or injured as a result of infection, for example, by an abscess or associated with infection by

human immunodeficiency virus, herpes zoster, or herpes simplex virus or with Lyme disease, tuberculosis, syphilis;

(iv) degenerative lesions, in which a portion of the nervous system is destroyed or injured as a result of a degenerative process including but not limited to degeneration

associated with Parkinson's disease, Alzheimer's disease, Huntington's chorea, or anyotrophic lateral sclerosis;

(v) lesions associated with nutritional diseases or disorders, in which a portion of the nervous system is destroyed or injured by a nutritional disorder or disorder of metabolism including but not limited to, vitamin B12 deficiency, folic acid deficiency, Wernicke disease, tobacco-alcohol amblyopia, Marchiafava-Bignami disease (primary degeneration of the corpus callosum), and alcoholic cerebellar degeneration;

(vi) neurological lesions associated with systemic diseases including but not limited to diabetes (diabetic neuropathy, Bell's palsy), systemic lupus erythematosus, carcinoma, or sarcoidosis;

(vii) lesions caused by toxic substances including alcohol, lead, or particular

neurotoxins; and

(viii) demyelinated lesions in which a portion of the nervous system is destroyed or injured by a demyelinating disease including but not limited to multiple sclerosis, human immunodeficiency virus-associated myelopathy, transverse myelopathy or various etiologies, progressive multifocal leukoencephalopathy, and central pontine myelinolysis.

Therapeutics which are useful according to the invention for treatment of a nervous system disorder may be selected by testing for biological activity in promoting the survival or differentiation of neurons. For example, and not by way of limitation, therapeutics which elicit any of the following effects may be useful according to the invention:

- 5 (i) increased survival time of neurons in culture;
- (ii) increased sprouting of neurons in culture or *in vivo*;
- (iii) increased production of a neuron-associated molecule in culture or *in vivo*, *e.g.*, choline acetyltransferase or acetylcholinesterase with respect to motor neurons; or
- (iv) decreased symptoms of neuron dysfunction *in vivo*.

10 Such effects may be measured by any method known in the art. In preferred, non-limiting embodiments, increased survival of neurons may be measured by the method set forth in Arakawa et al. (1990, J. Neurosci. 10:3507-3515); increased sprouting of neurons may be detected by methods set forth in Pestronk et al. (1980, Exp. Neurol. 70:65-82) or Brown et al. (1981, Ann. Rev. Neurosci. 4:17-42); increased production of neuron-associated molecules may be measured by bioassay, enzymatic assay, antibody binding, Northern blot assay, *etc.*, depending on the molecule to be measured; and motor neuron dysfunction may be measured by assessing the physical manifestation of motor neuron disorder, *e.g.*, weakness, motor neuron conduction velocity, or functional disability.

In specific embodiments, motor neuron disorders that may be treated according to the invention include but are not limited to disorders such as infarction, infection, exposure to toxin, trauma, surgical damage, degenerative disease or malignancy that may affect motor neurons as well as other components of the nervous system, as well as disorders that selectively affect neurons such as amyotrophic lateral sclerosis, and including but not limited to progressive spinal muscular atrophy, progressive bulbar palsy, primary lateral sclerosis, infantile and juvenile muscular atrophy, progressive bulbar paralysis of childhood (Fazio-Londe syndrome), poliomyelitis and the post polio syndrome, and Hereditary Motor sensory Neuropathy (Charcot-Marie-Tooth Disease).

#### 5.10.18 OTHER ACTIVITIES

30 A polypeptide of the invention may also exhibit one or more of the following additional activities or effects: inhibiting the growth, infection or function of, or killing, infectious agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or body part

size or shape (such as, for example, breast augmentation or diminution, change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism, processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, co-factors or other nutritional factors or component(s); effecting behavioral characteristics, including, without limitation, appetite, libido, stress, cognition (including cognitive disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic lineages; hormonal or endocrine activity; in the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and the ability to act as an antigen in a vaccine composition to raise an immune response against such protein or another material or entity which is cross-reactive with such protein.

15

#### 5.10.19 IDENTIFICATION OF POLYMORPHISMS

The demonstration of polymorphisms makes possible the identification of such polymorphisms in human subjects and the pharmacogenetic use of this information for diagnosis and treatment. Such polymorphisms may be associated with, e.g., differential predisposition or susceptibility to various disease states (such as disorders involving inflammation or immune response) or a differential response to drug administration, and this genetic information can be used to tailor preventive or therapeutic treatment appropriately. For example, the existence of a polymorphism associated with a predisposition to inflammation or autoimmune disease makes possible the diagnosis of this condition in humans by identifying the presence of the polymorphism.

25

Polymorphisms can be identified in a variety of ways known in the art which all generally involve obtaining a sample from a patient, analyzing DNA from the sample, optionally involving isolation or amplification of the DNA, and identifying the presence of the polymorphism in the DNA. For example, PCR may be used to amplify an appropriate fragment of genomic DNA which may then be sequenced. Alternatively, the DNA may be subjected to allele-specific oligonucleotide hybridization (in which appropriate oligonucleotides are hybridized to the DNA under conditions permitting detection of a single base mismatch) or to a single nucleotide extension assay (in which an oligonucleotide that hybridizes immediately adjacent to the position of the polymorphism is extended with one or more labeled

75

nucleotides). In addition, traditional restriction fragment length polymorphism analysis (using restriction enzymes that provide differential digestion of the genomic DNA depending on the presence or absence of the polymorphism) may be performed. Arrays with nucleotide sequences of the present invention can be used to detect polymorphisms. The array can comprise modified nucleotide sequences of the present invention in order to detect the nucleotide sequences of the present invention. In the alternative, any one of the nucleotide sequences of the present invention can be placed on the array to detect changes from those sequences.

Alternatively a polymorphism resulting in a change in the amino acid sequence could also be detected by detecting a corresponding change in amino acid sequence of the protein, e.g., by an antibody specific to the variant sequence.

10

#### 5.10.20 ARTHRITIS AND INFLAMMATION

The immunosuppressive effects of the compositions of the invention against rheumatoid arthritis is determined in an experimental animal model system. The experimental model system is adjuvant induced arthritis in rats, and the protocol is described by J. Holoshitz, et al., 1983, Science, 219:56, or by B. Waksman et al., 1963, Int. Arch. Allergy Appl. Immunol., 23:129. Induction of the disease can be caused by a single injection, generally intradermally, of a suspension of killed Mycobacterium tuberculosis in complete Freund's adjuvant (CFA). The route of injection can vary, but rats may be injected at the base of the tail with an adjuvant mixture. The polypeptide is administered in phosphate buffered solution (PBS) at a dose of about 1-5 mg/kg. The control consists of administering PBS only.

20

The procedure for testing the effects of the test compound would consist of intradermally injecting killed Mycobacterium tuberculosis in CFA followed by immediately administering the test compound and subsequent treatment every other day until day 24. At 14, 15, 18, 20, 22, and 24 days after injection of Mycobacterium CFA, an overall arthritis score may be obtained as described by J. Holoskitz above. An analysis of the data would reveal that the test compound would have a dramatic affect on the swelling of the joints as measured by a decrease of the arthritis score.

30

#### 5.11 THERAPEUTIC METHODS

The compositions (including polypeptide fragments, analogs, variants and antibodies or other binding partners or modulators including antisense polynucleotides) of the invention have

76

numerous applications in a variety of therapeutic methods. Examples of therapeutic applications include, but are not limited to, those exemplified herein.

#### 5.11.1 EXAMPLE

One embodiment of the invention is the administration of an effective amount of the GPCR-like polypeptides or other composition of the invention to individuals affected by a disease or disorder that can be modulated by regulating the peptides of the invention. While the mode of administration is not particularly important, parenteral administration is preferred. An exemplary mode of administration is to deliver an intravenous bolus. The dosage of GPCR-like polypeptides or other composition of the invention will normally be determined by the prescribing physician. It is to be expected that the dosage will vary according to the age, weight, condition and response of the individual patient. Typically, the amount of polypeptide administered per dose will be in the range of about 0.01 µg/kg to 100 mg/kg of body weight, with the preferred dose being about 0.1 µg/kg to 10 mg/kg of patient body weight. For parenteral administration, GPCR-like polypeptides of the invention will be formulated in an injectable form combined with a pharmaceutically acceptable parenteral vehicle. Such vehicles are well known in the art and examples include water, saline, Ringer's solution, dextrose solution, and solutions consisting of small amounts of the human serum albumin. The vehicle may contain minor amounts of additives that maintain the isotonicity and stability of the polypeptide or other active ingredient. The preparation of such solutions is within the skill of the art.

#### 5.12 PHARMACEUTICAL FORMULATIONS AND ROUTES OF

##### ADMINISTRATION

A protein or other composition of the present invention (from whatever source derived, including without limitation from recombinant and non-recombinant sources and including antibodies and other binding partners of the polypeptides of the invention) may be administered to a patient in need, by itself, or in pharmaceutical compositions where it is mixed with suitable carriers or excipient(s) at doses to treat or ameliorate a variety of disorders. Such a composition may optionally contain (in addition to protein or other active ingredient and a carrier) diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not

interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IFN, TNF $\alpha$ , TNF $\beta$ , G-CSF, Meg-CSF, thrombopoietin, stem cell factor, and erythropoietin. In further compositions, proteins of the invention may be combined with other agents beneficial to the treatment of the disease or disorder in question. These agents include various growth factors such as epidermal growth factor (EGF), platelet-derived growth factor (PDGF), transforming growth factors (TGF- $\alpha$  and TGF- $\beta$ ), insulin-like growth factor (IGF), as well as cytokines described herein.

The pharmaceutical composition may further contain other agents which either enhance the activity of the protein or other active ingredient or complement its activity or use in treatment. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with protein or other active ingredient of the invention, or to minimize side effects. Conversely, protein or other active ingredient of the present invention may be included in formulations of the particular clotting factor, cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the clotting factor, cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent (such as IL-1Ra, IL-1 Hy1, IL-1 Hy2, anti-TNF, corticosteroids, immunosuppressive agents). A protein of the present invention may be active in multimers (e.g., heterodimers or homodimers) or complexes with itself or other proteins. As a result, pharmaceutical compositions of the invention may comprise a protein of the invention in such multimeric or complexed form.

As an alternative to being included in a pharmaceutical composition of the invention including a first protein, a second protein or a therapeutic agent may be concurrently administered with the first protein (e.g., at the same time, or at differing times provided that therapeutic concentrations of the combination of agents is achieved at the treatment site). Techniques for formulation and administration of the compounds of the instant application may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition. A therapeutically effective dose further refers to that amount of the compound sufficient to result in amelioration of symptoms, e.g., treatment, healing, prevention or amelioration of the relevant medical condition, or an increase in rate of treatment, healing,

prevention or amelioration of such conditions. When applied to an individual active ingredient, administered alone, a therapeutically effective dose refers to that ingredient alone. When applied to a combination, a therapeutically effective dose refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

In practicing the method of treatment or use of the present invention, a therapeutically effective amount of protein or other active ingredient of the present invention is administered to a mammal having a condition to be treated. Protein or other active ingredient of the present invention may be administered in accordance with the method of the invention either alone or in combination with other therapies such as treatments employing cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines or other hematopoietic factors, protein or other active ingredient of the present invention may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering protein or other active ingredient of the present invention in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

#### 5.12.1 ROUTES OF ADMINISTRATION

Suitable routes of administration may, for example, include oral, rectal, transmucosal, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections. Administration of protein or other active ingredient of the present invention used in the pharmaceutical composition or to practice the method of the present invention can be carried out in a variety of conventional ways, such as oral ingestion, inhalation, topical application or cutaneous, subcutaneous, intraperitoneal, parenteral or intravenous injection. Intravenous administration to the patient is preferred.

Alternately, one may administer the compound in a local rather than systemic manner, for example, via injection of the compound directly into a arthritic joints or in fibrotic tissue, often in a depot or sustained release formulation. In order to prevent the scarring process frequently occurring as complication of glaucoma surgery, the compounds may be administered topically, for example, as eye drops. Furthermore, one may administer the drug in a targeted drug delivery system, for example, in a liposome coated with a specific antibody,

targeting, for example, arthritic or fibrotic tissue. The liposomes will be targeted to and taken up selectively by the afflicted tissue.

The polypeptides of the invention are administered by any route that delivers an effective dosage to the desired site of action. The determination of a suitable route of administration and an effective dosage for a particular indication is within the level of skill in the art. Preferably for wound treatment, one administers the therapeutic compound directly to the site. Suitable dosage ranges for the polypeptides of the invention can be extrapolated from these dosages or from similar studies in appropriate animal models. Dosages can then be adjusted as necessary by the clinician to provide maximal therapeutic benefit.

#### 5.12.2 COMPOSITIONS/FORMULATIONS

Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. These pharmaceutical compositions may be manufactured in a manner that is itself known, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapment or lyophilizing processes. Proper formulation is dependent upon the route of administration chosen. When a therapeutically effective amount of protein or other active ingredient of the present invention is administered orally, protein or other active ingredient of the present invention will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95 % protein or other active ingredient of the present invention, and preferably from about 25 to 90 % protein or other active ingredient of the present invention. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90 % by weight of protein or other active ingredient of the present invention, and preferably from about 1 to 50 % protein or other active ingredient of the present invention.

When a therapeutically effective amount of protein or other active ingredient of the present invention is administered by intravenous, cutaneous or subcutaneous injection, protein or other active ingredient of the present invention will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein or other active ingredient solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in addition to protein or other active ingredient of the present invention, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art. For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate. Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration. For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, *e.g.*, gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch. The compounds may be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, *e.g.*, in ampules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other

glycerides. In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

A pharmaceutical carrier for the hydrophobic compounds of the invention is a co-solvent system comprising benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. The co-solvent system may be the VPD co-solvent system. VPD is a solution of 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant polysorbate 80, and 65% w/v polyethylene glycol 300, made up to volume in absolute ethanol. The VPD co-solvent system (VPD:5W) consists of VPD diluted 1:1 with a 5% dextrose in water solution. This co-solvent system dissolves hydrophobic compounds well, and itself produces low toxicity upon systemic administration. Naturally, the proportions of a co-solvent system may be varied considerably without destroying its solubility and toxicity characteristics.

Furthermore, the identity of the co-solvent components may be varied: for example, other low-toxicity nonpolar surfactants may be used instead of polysorbate 80; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may replace polyethylene glycol, e.g. polyvinyl pyrrolidone; and other sugars or polysaccharides may substitute for dextrose. Alternatively, other delivery systems for hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed, although usually at the cost of greater toxicity. Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various types of sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein or other active ingredient stabilization may be employed.

The pharmaceutical compositions also may comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols. Many of the active ingredients of the invention may be

provided as salts with pharmaceutically compatible counter ions. Such pharmaceutically acceptable base addition salts are those salts which retain the biological effectiveness and properties of the free acids and which are obtained by reaction with inorganic or organic bases such as sodium hydroxide, magnesium hydroxide, ammonia, trialkylamine, dialkylamine, monoalkylamine, dibasic amino acids, sodium acetate, potassium benzoate, triethanol amine and the like.

The pharmaceutical composition of the invention may be in the form of a complex of the protein(s) or other active ingredient of present invention along with protein or peptide antigens. The protein and/or peptide antigen will deliver a stimulatory signal to both B and T lymphocytes. B lymphocytes will respond to antigen through their surface immunoglobulin receptor. T lymphocytes will respond to antigen through the T cell receptor (TCR) following presentation of the antigen by MHC proteins. MHC and structurally related proteins including those encoded by class I and class II MHC genes on host cells will serve to present the peptide antigen(s) to T lymphocytes. The antigen components could also be supplied as purified MHC-peptide complexes alone or with co-stimulatory molecules that can directly signal T cells. Alternatively antibodies able to bind surface immunoglobulin and other molecules on B cells as well as antibodies able to bind the TCR and other molecules on T cells can be combined with the pharmaceutical composition of the invention.

The pharmaceutical composition of the invention may be in the form of a liposome in which protein of the present invention is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, surfactides, lysolecithins, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Patent Nos. 4,235,871; 4,501,728; 4,837,028; and 4,737,323, all of which are incorporated herein by reference.

The amount of protein or other active ingredient of the present invention in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of protein or other active ingredient of the present invention with which to treat each individual patient. Initially, the attending physician will administer low doses of protein or other active ingredient of the present invention and observe the patient's response. Larger doses of protein or other active

ingredient of the present invention may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.01  $\mu\text{g}$  to about 100 mg (preferably about 0.1  $\mu\text{g}$  to about 10 mg, more preferably about 0.1  $\mu\text{g}$  to about 1 mg) of protein or other active ingredient of the present invention per kg body weight. For compositions of the present invention which are useful for bone, cartilage, tendon or ligament regeneration, the therapeutic method includes administering the composition topically, systemically, or locally as an implant or device. When administered, the therapeutic composition for use in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage or tissue damage. Topical administration may be suitable for wound healing and tissue repair. Therapeutically useful agents other than a protein or other active ingredient of the invention which may also optionally be included in the composition as described above, may alternatively or additionally, be administered simultaneously or sequentially with the composition in the methods of the invention. Preferably for bone and/or cartilage formation, the composition would include a matrix capable of delivering the protein-containing or other active ingredient-containing composition to the site of bone and/or cartilage damage, providing a structure for the developing bone and cartilage and optimally capable of being resorbed into the body. Such matrices may be formed of materials presently in use for other implanted medical applications.

The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance and interface properties. The particular application of the compositions will define the appropriate formulation. Potential matrices for the compositions may be biodegradable and chemically defined calcium sulfate, tricalcium phosphate, hydroxyapatite, polylactic acid, polyglycolic acid and polyanhydrides. Other potential materials are biodegradable and biologically well-defined, such as bone or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are nonbiodegradable and chemically defined, such as sintered hydroxyapatite, bioglass, aluminates, or other ceramics. Matrices may be comprised of combinations of any of the above mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalcium phosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size, particle shape, and biodegradability. Presently preferred is a 50:50 (mole weight)

copolymer of lactic acid and glycolic acid in the form of porous particles having diameters ranging from 150 to 800 microns. In some applications, it will be useful to utilize a sequestering agent, such as carboxymethyl cellulose or autologous blood clot, to prevent the protein compositions from disassociating from the matrix.

A preferred family of sequestering agents is cellulosic materials such as alkylcelluloses (including hydroxyalkylcelluloses), including methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropyl-methylcellulose, and carboxymethylcellulose, the most preferred being cationic salts of carboxymethylcellulose (CMC). Other preferred sequestering agents include hyaluronic acid, sodium alginate, poly(ethylene glycol), polyoxyethylene oxide, carboxyvinyl polymer and poly(vinyl alcohol). The amount of sequestering agent useful herein is 0.5-20 wt %, preferably 1-10 wt % based on total formulation weight, which represents the amount necessary to prevent desorption of the protein from the polymer matrix and to provide appropriate handling of the composition, yet not so much that the progenitor cells are prevented from infiltrating the matrix, thereby providing the protein the opportunity to assist the osteogenic activity of the progenitor cells. In further compositions, proteins or other active ingredient of the invention may be combined with other agents beneficial to the treatment of the bone and/or cartilage defect, wound, or tissue in question. These agents include various growth factors such as epidermal growth factor (EGF), platelet derived growth factor (PDGF), transforming growth factors (TGF- $\alpha$  and TGF- $\beta$ ), and insulin-like growth factor (IGF).

The therapeutic compositions are also presently valuable for veterinary applications. Particularly domestic animals and thoroughbred horses, in addition to humans, are desired patients for such treatment with proteins or other active ingredient of the present invention. The dosage regimen of a protein-containing pharmaceutical composition to be used in tissue regeneration will be determined by the attending physician considering various factors which modify the action of the proteins, e.g., amount of tissue weight desired to be formed, the site of damage, the condition of the damaged tissue, the size of a wound, type of damaged tissue (e.g., bone), the patient's age, sex, and diet, the severity of any infection, time of administration and other clinical factors. The dosage may vary with the type of matrix used in the reconstitution and with inclusion of other proteins in the pharmaceutical composition. For example, the addition of other known growth factors, such as IGF I (insulin like growth factor I), to the final composition, may also effect the dosage. Progress can be monitored by periodic



assessment of tissue/bone growth and/or repair, for example, X-rays, histomorphometric determinations and tetracycline labeling.

Polynucleotides of the present invention can also be used for gene therapy. Such polynucleotides can be introduced either *in vivo* or *ex vivo* into cells for expression in a mammalian subject. Polynucleotides of the invention may also be administered by other known methods for introduction of nucleic acid into a cell or organism (including, without limitation, in the form of viral vectors or naked DNA). Cells may also be cultured *ex vivo* in the presence of proteins of the present invention in order to proliferate or to produce a desired effect on or activity in such cells. Treated cells can then be introduced *in vivo* for therapeutic purposes.

### 5.12.3 EFFECTIVE DOSAGE

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve its intended purpose. More specifically, a therapeutically effective amount means an amount effective to prevent development of or to alleviate the existing symptoms of the subject being treated. Determination of the effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from appropriate *in vitro* assays. For example, a dose can be formulated in animal models to achieve a circulating concentration range that can be used to more accurately determine useful doses in humans. For example, a dose can be formulated in animal models to achieve a circulating concentration range that includes the  $IC_{50}$  as determined in cell culture (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of the protein's biological activity). Such information can be used to more accurately determine useful doses in humans.

A therapeutically effective dose refers to that amount of the compound that results in amelioration of symptoms or a prolongation of survival in a patient. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the  $LD_{50}$  (the dose lethal to 50% of the population) and the  $ED_{50}$  (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between  $LD_{50}$  and  $ED_{50}$ . Compounds which exhibit high therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the  $ED_{50}$  with little or no

toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. See, *e.g.*,

Fingl et al., 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p.1. Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the desired effects, or minimal effective concentration (MEC). The MEC will vary for each compound but can be estimated from *in vitro* data. Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. However, HPLC assays or bioassays can be used to determine plasma concentrations.

Dosage intervals can also be determined using MEC value. Compounds should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90%. In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

An exemplary dosage regimen for polypeptides or other compositions of the invention will be in the range of about 0.01  $\mu\text{g/kg}$  to 100  $\text{mg/kg}$  of body weight daily, with the preferred dose being about 0.1  $\mu\text{g/kg}$  to 25  $\text{mg/kg}$  of patient body weight daily, varying in adults and children. Dosing may be once daily, or equivalent doses may be delivered at longer or shorter intervals.

The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's age and weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

### 5.12.4 PACKAGING

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may, for example, comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

### 5.13 ANTIBODIES

Also included in the invention are antibodies to proteins, or fragments of proteins of the invention. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin (Ig) molecules, i.e., molecules that contain an antigen binding site that specifically binds (immunoreacts with) an antigen. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, F<sub>ab</sub>, F<sub>w</sub> and F<sub>ab</sub>'<sub>2</sub> fragments, and an F<sub>as</sub> expression library. In general, an antibody molecule obtained from humans relates to any of the classes IgG, IgM, IgA, IgE and IgD, which differ from one another by the nature of the heavy chain present in the molecule. Certain classes have subclasses as well, such as IgG<sub>1</sub>, IgG<sub>2</sub>, and others. Furthermore, in humans, the light chain may be a kappa chain or a lambda chain. Reference herein to antibodies includes a reference to all such classes, subclasses and types of human antibody species.

An isolated related protein of the invention may be intended to serve as an antigen, or a portion or fragment thereof, and additionally can be used as an immunogen to generate antibodies that immunospecifically bind the antigen, using standard techniques for polyclonal and monoclonal antibody preparation. The full-length protein can be used or, alternatively, the invention provides antigenic peptide fragments of the antigen for use as immunogens. An antigenic peptide fragment comprises at least 6 amino acid residues of the amino acid sequence of the full length protein, such as an amino acid sequence shown in SEQ ID NO: 4, and encompasses an epitope thereof such that an antibody raised against the peptide forms a specific immune complex with the full length protein or with any fragment that contains the epitope. Preferably, the antigenic peptide comprises at least 10 amino acid residues, or at least 15 amino acid residues, or at least 20 amino acid residues, or at least 30 amino acid residues. Preferred epitopes encompassed by the antigenic peptide are regions of the protein that are located on its surface; commonly these are hydrophilic regions.

In certain embodiments of the invention, at least one epitope encompassed by the antigenic peptide is a region of -related protein that is located on the surface of the protein, e.g., a hydrophilic region. A hydrophobicity analysis of the human related protein sequence will indicate which regions of a related protein are particularly hydrophilic and, therefore, are likely to encode surface residues useful for targeting antibody production. As a means for targeting antibody production, hydrophathy plots showing regions of hydrophilicity and hydrophobicity may be generated by any method well known in the art, including, for

example, the Kyte Doolittle or the Hopp Woods methods, either with or without Fourier transformation. See, e.g., Hopp and Woods, 1981, *Proc. Nat. Acad. Sci. USA* 78: 3824-3828; Kyte and Doolittle 1982, *J. Mol. Biol.* 157: 105-142, each of which is incorporated herein by reference in its entirety. Antibodies that are specific for one or more domains within an antigenic protein, or derivatives, fragments, analogs or homologs thereof, are also provided herein.

A protein of the invention, or a derivative, fragment, analog, homolog or ortholog thereof, may be utilized as an immunogen in the generation of antibodies that immunospecifically bind these protein components.

Various procedures known within the art may be used for the production of polyclonal or monoclonal antibodies directed against a protein of the invention, or against derivatives, fragments, analogs homologs or orthologs thereof (see, for example, *Antibodies: A Laboratory Manual*, Harlow E, and Lane D, 1988, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, incorporated herein by reference). Some of these antibodies are discussed below.

#### 5.13.1 Polyclonal Antibodies

For the production of polyclonal antibodies, various suitable host animals (e.g., rabbit, goat, mouse or other mammal) may be immunized by one or more injections with the native protein, a synthetic variant thereof, or a derivative of the foregoing. An appropriate immunogenic preparation can contain, for example, the naturally occurring immunogenic protein, a chemically synthesized polypeptide representing the immunogenic protein, or a recombinantly expressed immunogenic protein. Furthermore, the protein may be conjugated to a second protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. The preparation can further include an adjuvant. Various adjuvants used to increase the immunological response include, but are not limited to, Freund's (complete and incomplete), mineral gels (e.g., aluminum hydroxide), surface active substances (e.g., lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, dinitrophenol, etc.), adjuvants usable in humans such as Bacille Calmette-Guerin and Corynebacterium parvum, or similar immunostimulatory agents. Additional examples of adjuvants which can be employed include MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate).

The polyclonal antibody molecules directed against the immunogenic protein can be isolated from the mammal (e.g., from the blood) and further purified by well known

techniques, such as affinity chromatography using protein A or protein G, which provide primarily the IgG fraction of immune serum. Subsequently, or alternatively, the specific antigen which is the target of the immunoglobulin sought, or an epitope thereof, may be immobilized on a column to purify the immune specific antibody by immunoaffinity chromatography. Purification of immunoglobulins is discussed, for example, by D. Wilkinson (The Scientist, published by The Scientist, Inc., Philadelphia PA, Vol. 14, No. 8 (April 17, 2000), pp. 25-28).

### 5.13.2 Monoclonal Antibodies

The term "monoclonal antibody" (MAb) or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one molecular species of antibody molecule consisting of a unique light chain gene product and a unique heavy chain gene product. In particular, the complementarity determining regions (CDRs) of the monoclonal antibody are identical in all the molecules of the population. MAbs thus contain an antigen binding site capable of immunoreacting with a particular epitope of the antigen characterized by a unique binding affinity for it.

Monoclonal antibodies can be prepared using hybridoma methods, such as those described by Kohler and Milstein, *Nature*, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes can be immunized *in vitro*.

The immunizing agent will typically include the protein antigen, a fragment thereof or a fusion protein thereof. Generally, either peripheral blood lymphocytes are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, (1986) pp. 59-103). Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells can be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas

typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, J. Immunol., 133:3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York, (1987) pp. 51-63).

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunosorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980). Preferably, antibodies having a high degree of specificity and a high binding affinity for the target antigen are isolated.

After the desired hybridoma cells are identified, the clones can be subcloned by limiting dilution procedures and grown by standard methods. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells can be grown *in vivo* as ascites in a mammal.

The monoclonal antibodies secreted by the subclones can be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies can also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA can be placed into expression vectors,

which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also can be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; Morrison, *Nature* 368, 812-13 (1994)) or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

### 5.13.2 Humanized Antibodies

The antibodies directed against the protein antigens of the invention can further comprise humanized antibodies or human antibodies. These antibodies are suitable for administration to humans without engendering an immune response by the human against the administered immunoglobulin. Humanized forms of antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')<sub>2</sub> or other antigen-binding subsequences of antibodies) that are principally comprised of the sequence of a human immunoglobulin, and contain minimal sequence derived from a non-human immunoglobulin. Humanization can be performed following the method of Winter and co-workers (Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-327 (1988); Verhoeven et al., *Science*, 229:1534-1536 (1988)), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. (See also U.S. Patent No. 5,225,539.) In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies can also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the framework regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin (Jones et al., 1986; Riechmann et al., 1988; and Presta, *Curr. Op. Struct. Biol.*, 2:593-596 (1992)).

### 5.13.3 Human Antibodies

Fully human antibodies relate to antibody molecules in which essentially the entire sequences of both the light chain and the heavy chain, including the CDRs, arise from human genes. Such antibodies are termed "human antibodies", or "fully human antibodies" herein.

Human monoclonal antibodies can be prepared by the trioma technique; the human B-cell hybridoma technique (see Kozbor, et al., 1983 *Immunol Today* 4: 72) and the EBV hybridoma technique to produce human monoclonal antibodies (see Cole, et al., 1985 In: *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc., pp. 77-96). Human monoclonal antibodies may be utilized in the practice of the present invention and may be produced by using human hybridomas (see Cole, et al., 1983. *Proc Natl Acad Sci USA* 80: 2026-2030) or by transforming human B-cells with Epstein Barr Virus in vitro (see Cole, et al., 1985 In: *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc., pp. 77-96).

In addition, human antibodies can also be produced using additional techniques, including phage display libraries (Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991)). Similarly, human antibodies can be made by introducing human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in Marks et al. (*BioTechnology* 10, 779-783 (1992)); Lonberg et al. (*Nature* 368 856-859 (1994)); Morrison (*Nature* 368, 812-13 (1994)); Fishwild et al. (*Nature Biotechnology* 14, 845-51 (1996)); Neuberger (*Nature Biotechnology* 14, 826 (1996)); and Lonberg and Huszar (*Intern. Rev. Immunol.* 13 65-93 (1995)).

Human antibodies may additionally be produced using transgenic nonhuman animals which are modified so as to produce fully human antibodies rather than the animal's endogenous antibodies in response to challenge by an antigen. (See PCT publication WO94/02602). The endogenous genes encoding the heavy and light immunoglobulin chains in the nonhuman host have been inactivated, and active loci encoding human heavy and light chain immunoglobulins are inserted into the host's genome. The human genes are incorporated, for example, using yeast artificial chromosomes containing the requisite human DNA segments. An animal which provides all the desired modifications is then obtained as progeny by crossbreeding intermediate transgenic animals containing fewer than the full complement of the modifications. The preferred embodiment of such a nonhuman animal is a

mouse, and is termed the Xenomouse™ as disclosed in PCT publications WO 96/33735 and WO 96/34096. This animal produces B cells which secrete fully human immunoglobulins. The antibodies can be obtained directly from the animal after immunization with an immunogen of interest, as, for example, a preparation of a polyclonal antibody, or alternatively from immortalized B cells derived from the animal, such as hybridomas producing monoclonal antibodies. Additionally, the genes encoding the immunoglobulins with human variable regions can be recovered and expressed to obtain the antibodies directly, or can be further modified to obtain analogs of antibodies such as, for example, single chain Fv molecules.

10 An example of a method of producing a nonhuman host, exemplified as a mouse, lacking expression of an endogenous immunoglobulin heavy chain is disclosed in U.S. Patent No. 5,939,598. It can be obtained by a method including deleting the J segment genes from at least one endogenous heavy chain locus in an embryonic stem cell to prevent rearrangement of the locus and to prevent formation of a transcript of a rearranged immunoglobulin heavy chain locus, the deletion being effected by a targeting vector containing a gene encoding a selectable marker; and producing from the embryonic stem cell a transgenic mouse whose somatic and germ cells contain the gene encoding the selectable marker.

15 A method for producing an antibody of interest, such as a human antibody, is disclosed in U.S. Patent No. 5,916,771. It includes introducing an expression vector that contains a nucleotide sequence encoding a heavy chain into one mammalian host cell in culture, introducing an expression vector containing a nucleotide sequence encoding a light chain into another mammalian host cell, and fusing the two cells to form a hybrid cell. The hybrid cell expresses an antibody containing the heavy chain and the light chain.

20 In a further improvement on this procedure, a method for identifying a clinically relevant epitope on an immunogen, and a correlative method for selecting an antibody that binds immunospecifically to the relevant epitope with high affinity, are disclosed in PCT publication WO 99/53049.

#### 5.13.4 F<sub>ab</sub> Fragments and Single Chain Antibodies

30 According to the invention, techniques can be adapted for the production of single-chain antibodies specific to an antigenic protein of the invention (see e.g., U.S. Patent No. 4,946,778). In addition, methods can be adapted for the construction of F<sub>ab</sub> expression libraries (see e.g., Huse, et al., 1989 Science 246: 1275-1281) to allow rapid and effective identification of monoclonal F<sub>ab</sub> fragments with the desired specificity for a protein or

derivatives, fragments, analogs or homologs thereof. Antibody fragments that contain the idiotype to a protein antigen may be produced by techniques known in the art including, but not limited to: (i) an F<sub>ab</sub> fragment produced by pepsin digestion of an antibody molecule; (ii) an F<sub>ab</sub> fragment generated by reducing the disulfide bridges of an F<sub>ab</sub> fragment; (iii) an F<sub>ab</sub> fragment generated by the treatment of the antibody molecule with papain and a reducing agent and (iv) F<sub>v</sub> fragments.

#### 5.13.5 Bispecific Antibodies

30 Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for an antigenic protein of the invention. The second binding target is any other antigen, and advantageously is a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, *Nature*, 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker *et al.*, 1991 *EMBO J.*, 10:3655-3659.

40 Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh *et al.*, *Methods in Enzymology*, 121:210 (1986).

50 According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid

side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')<sub>2</sub> bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan et al., *Science* 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')<sub>2</sub> fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Additionally, Fab' fragments can be directly recovered from *E. coli* and chemically coupled to form bispecific antibodies. Shalaby et al., *J. Exp. Med.* 175:217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')<sub>2</sub> molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny et al., *J. Immunol.* 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger et al., *Proc. Natl. Acad. Sci. USA* 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The

fragments comprise a heavy-chain variable domain (V<sub>H</sub>) connected to a light-chain variable domain (V<sub>L</sub>) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V<sub>H</sub> and V<sub>L</sub> domains of one fragment are forced to pair with the complementary V<sub>L</sub> and V<sub>H</sub> domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFc) dimers has also been reported. See, Gruber et al., *J. Immunol.* 152:5368 (1994).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al., *J. Immunol.* 147:60 (1991).

Exemplary bispecific antibodies can bind to two different epitopes, at least one of which originates in the protein antigen of the invention. Alternatively, an anti-antigenic arm of an immunoglobulin molecule can be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g. CD2, CD3, CD28, or B7), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16) so as to focus cellular defense mechanisms to the cell expressing the particular antigen. Bispecific antibodies can also be used to direct cytotoxic agents to cells which express a particular antigen. These antibodies possess an antigen-binding arm and an arm which binds a cytotoxic agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA. Another bispecific antibody of interest binds the protein antigen described herein and further binds tissue factor (TF).

#### 5.13.6 Heteroconjugate Antibodies

Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360; WO 92/200373; EP 03089). It is contemplated that the antibodies can be prepared *in vitro* using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins can be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

#### 5.13.7 Effector Function Engineering

It can be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, e.g., the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) can be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated can have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron et al., *J. Exp. Med.*, 176: 1191-1195 (1992) and Shopes, *J. Immunol.*, 148: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity can also be prepared using heterobifunctional cross-linkers as described in Wolff et al. *Cancer Research*, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and can thereby have enhanced complement lysis and ADCC capabilities. See Stevenson et al., *Anti-Cancer Drug Design*, 3: 219-230 (1989).

#### 5.13.8 Immunoconjugates

The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolacca americana proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, saponaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include  $^{212}\text{Bi}$ ,  $^{131}\text{I}$ ,  $^{125}\text{I}$ ,  $^{90}\text{Y}$ , and  $^{186}\text{Re}$ .

Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimide HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene).

For example, a ricin immunotoxin can be prepared as described in Viteita et al., *Science*, 238:

1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminopentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionuclide to the antibody. See WO94/11026.

In another embodiment, the antibody can be conjugated to a "receptor" (such streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) that is in turn conjugated to a cytotoxic agent.

#### 5.14 COMPUTER READABLE SEQUENCES

In one application of this embodiment, a nucleotide sequence of the present invention can be recorded on computer readable media. As used herein, "computer readable media" refers to any medium which can be read and accessed directly by a computer. Such media include, but are not limited to: magnetic storage media, such as floppy discs, hard disc storage medium, and magnetic tape; optical storage media such as CD-ROM; electrical storage media such as RAM and ROM; and hybrids of these categories such as magnetic/optical storage media. A skilled artisan can readily appreciate how any of the presently known computer readable mediums can be used to create a manufacture comprising computer readable medium having recorded thereon a nucleotide sequence of the present invention. As used herein, "recorded" refers to a process for storing information on computer readable medium. A skilled artisan can readily adopt any of the presently known methods for recording information on computer readable medium to generate manufactures comprising the nucleotide sequence information of the present invention.

A variety of data storage structures are available to a skilled artisan for creating a computer readable medium having recorded thereon a nucleotide sequence of the present invention. The choice of the data storage structure will generally be based on the means chosen to access the stored information. In addition, a variety of data processor programs and formats can be used to store the nucleotide sequence information of the present invention on computer readable medium. The sequence information can be represented in a word processing text file, formatted in commercially-available software such as WordPerfect and Microsoft Word, or represented in the form of an ASCII file, stored in a database application, such as DB2, Sybase, Oracle, or the like. A skilled artisan can readily adapt any number of data processor structuring formats (e.g. text file or database) in order to obtain computer

readable medium having recorded thereon the nucleotide sequence information of the present invention.

By providing any of the nucleotide sequences SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 or a representative fragment thereof; or a nucleotide sequence at least 95% identical to any of the nucleotide sequences of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 in computer readable form, a skilled artisan can routinely access the sequence information for a variety of purposes. Computer software is publicly available which allows a skilled artisan to access sequence information provided in a computer readable medium. The examples which follow

demonstrate how software which implements the BLAST (Altschul et al., *J. Mol. Biol.* 215:403-410 (1990)) and BLAZE (Brudlag et al., *Comp. Chem.* 17:203-207 (1993)) search algorithms on a Sybase system is used to identify open reading frames (ORFs) within a nucleic acid sequence. Such ORFs may be protein encoding fragments and may be useful in producing commercially important proteins such as enzymes used in fermentation reactions and in the production of commercially useful metabolites.

As used herein, "a computer-based system" refers to the hardware means, software means, and data storage means used to analyze the nucleotide sequence information of the present invention. The minimum hardware means of the computer-based systems of the present invention comprises a central processing unit (CPU), input means, output means, and data storage means. A skilled artisan can readily appreciate that any one of the currently available computer-based systems are suitable for use in the present invention. As stated above, the computer-based systems of the present invention comprise a data storage means having stored therein a nucleotide sequence of the present invention and the necessary hardware means and software means for supporting and implementing a search means. As used herein, "data storage means" refers to memory which can store nucleotide sequence information of the present invention, or a memory access means which can access manufactures having recorded thereon the nucleotide sequence information of the present invention.

As used herein, "search means" refers to one or more programs which are implemented on the computer-based system to compare a target sequence or target structural motif with the sequence information stored within the data storage means. Search means are used to identify fragments or regions of a known sequence which match a particular target sequence or target motif. A variety of known algorithms are disclosed publicly and a variety of commercially available software for conducting search means are and can be used in the computer-based systems of the present invention. Examples of such software includes, but is not limited to,

Smith-Waterman, MacPattern (EMBL), BLASTN and BLASTA (NPOLYPEPTIDEIA). A skilled artisan can readily recognize that any one of the available algorithms or implementing software packages for conducting homology searches can be adapted for use in the present computer-based systems. As used herein, a "target sequence" can be any nucleic acid or amino acid sequence of six or more nucleotides or two or more amino acids. A skilled artisan can readily recognize that the longer a target sequence is, the less likely a target sequence will be present as a random occurrence in the database. The most preferred sequence length of a target sequence is from about 10 to 100 amino acids, or from about 30 to 300 nucleotide residues. However, it is well recognized that searches for commercially important fragments, such as sequence fragments involved in gene expression and protein processing, may be of shorter length.

As used herein, "a target structural motif," or "target motif," refers to any rationally selected sequence or combination of sequences in which the sequence(s) are chosen based on a three-dimensional configuration which is formed upon the folding of the target motif. There are a variety of target motifs known in the art. Protein target motifs include, but are not limited to, enzyme active sites and signal sequences. Nucleic acid target motifs include, but are not limited to, promoter sequences, hairpin structures and inducible expression elements (protein binding sequences).

## 20 5.15 TRIPLE HELIX FORMATION

In addition, the fragments of the present invention, as broadly described, can be used to control gene expression through triple helix formation or antisense DNA or RNA, both of which methods are based on the binding of a polynucleotide sequence to DNA or RNA. Polynucleotides suitable for use in these methods are usually 20 to 40 bases in length and are designed to be complementary to a region of the gene involved in transcription (triple helix - see Lee et al., *Nucl. Acids Res.* 6:3073 (1979); Cooney et al., *Science* 15241:456 (1988); and Dervan et al., *Science* 251:1360 (1991)) or to the mRNA itself (antisense - Olmno, *J. Neurochem.* 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). Triple helix-formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques have been demonstrated to be effective in model systems. Information contained in the sequences of the present invention is necessary for the design of an antisense or triple helix oligonucleotide.



## 5.16 DIAGNOSTIC ASSAYS AND KITS

The present invention further provides methods to identify the presence or expression of one of the ORFs of the present invention, or homolog thereof, in a test sample, using a nucleic acid probe or antibodies of the present invention, optionally conjugated or otherwise associated with a suitable label.

In general, methods for detecting a polynucleotide of the invention can comprise contacting a sample with a compound that binds to and forms a complex with the polynucleotide for a period sufficient to form the complex, and detecting the complex, so that if a complex is detected, a polynucleotide of the invention is detected in the sample. Such methods can also comprise contacting a sample under stringent hybridization conditions with nucleic acid primers that anneal to a polynucleotide of the invention under such conditions, and amplifying annealed polynucleotides, so that if a polynucleotide is amplified, a polynucleotide of the invention is detected in the sample.

In general, methods for detecting a polypeptide of the invention can comprise contacting a sample with a compound that binds to and forms a complex with the polypeptide for a period sufficient to form the complex, and detecting the complex, so that if a complex is detected, a polypeptide of the invention is detected in the sample.

In detail, such methods comprise incubating a test sample with one or more of the antibodies or one or more of the nucleic acid probes of the present invention and assaying for binding of the nucleic acid probes or antibodies to components within the test sample.

Conditions for incubating a nucleic acid probe or antibody with a test sample vary.

Incubation conditions depend on the format employed in the assay, the detection methods employed, and the type and nature of the nucleic acid probe or antibody used in the assay. One skilled in the art will recognize that any one of the commonly available hybridization, amplification or immunological assay formats can readily be adapted to employ the nucleic acid probes or antibodies of the present invention. Examples of such assays can be found in

Chard, T., *An Introduction to Radioimmunoassay and Related Techniques*, Elsevier Science Publishers, Amsterdam, The Netherlands (1986); Bullock, G.R. et al., *Techniques in Immunocytochemistry*, Academic Press, Orlando, FL Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985); Tijssen, P., *Practice and Theory of immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers, Amsterdam, The Netherlands (1985). The test samples of the present invention include cells, protein or membrane extracts of cells, or biological fluids such as sputum, blood, serum, plasma, or urine. The test sample used in the above-described method will vary based on the assay

format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed. Methods for preparing protein extracts or membrane extracts of cells are well known in the art and can be readily be adapted in order to obtain a sample which is compatible with the system utilized.

In another embodiment of the present invention, kits are provided which contain the necessary reagents to carry out the assays of the present invention. Specifically, the invention provides a compartment kit to receive, in close confinement, one or more containers which comprises: (a) a first container comprising one of the probes or antibodies of the present invention; and (b) one or more other containers comprising one or more of the following: wash reagents, reagents capable of detecting presence of a bound probe or antibody.

In detail, a compartment kit includes any kit in which reagents are contained in separate containers. Such containers include small glass containers, plastic containers or strips of plastic or paper. Such containers allow one to efficiently transfer reagents from one compartment to another compartment such that the samples and reagents are not cross-contaminated, and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers will include a container which will accept the test sample, a container which contains the antibodies used in the assay, containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, etc.), and containers which contain the reagents used to detect the bound antibody or probe. Types of detection reagents include labeled nucleic acid probes, labeled secondary antibodies, or in the alternative, if the primary antibody is labeled, the enzymatic, or antibody binding reagents which are capable of reacting with the labeled antibody. One skilled in the art will readily recognize that the disclosed probes and antibodies of the present invention can be readily incorporated into one of the established kit formats which are well known in the art.

## 5.17 MEDICAL IMAGING

The novel polypeptides and binding partners of the invention are useful in medical imaging of sites expressing the molecules of the invention (e.g., where the polypeptide of the invention is involved in the immune response, for imaging sites of inflammation or infection).

See, e.g., Kunkel et al., U.S. Pat. NO. 5,413,778. Such methods involve chemical attachment of a labeling or imaging agent, administration of the labeled polypeptide to a subject in a pharmaceutically acceptable carrier, and imaging the labeled polypeptide *in vivo* at the target site.

## 5.18 SCREENING ASSAYS

Using the isolated proteins and polynucleotides of the invention, the present invention further provides methods of obtaining and identifying agents which bind to a polypeptide encoded by an ORF corresponding to any of the nucleotide sequences set forth in SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62, or bind to a specific domain of the polypeptide encoded by the nucleic acid. In detail, said method comprises the steps of:

- (a) contacting an agent with an isolated protein encoded by an ORF of the present invention, or nucleic acid of the invention; and
  - (b) determining whether the agent binds to said protein or said nucleic acid.
- In general, therefore, such methods for identifying compounds that bind to a polynucleotide of the invention can comprise contacting a compound with a polynucleotide of the invention for a time sufficient to form a polynucleotide/compound complex, and detecting the complex, so that if a polynucleotide/compound complex is detected, a compound that binds to a polynucleotide of the invention is identified.

Likewise, in general, therefore, such methods for identifying compounds that bind to a polypeptide of the invention can comprise contacting a compound with a polypeptide of the invention for a time sufficient to form a polypeptide/compound complex, and detecting the complex, so that if a polypeptide/compound complex is detected, a compound that binds to a polynucleotide of the invention is identified.

Methods for identifying compounds that bind to a polypeptide of the invention can also comprise contacting a compound with a polypeptide of the invention in a cell for a time sufficient to form a polypeptide/compound complex, wherein the complex drives expression of a receptor gene sequence in the cell, and detecting the complex by detecting reporter gene sequence expression, so that if a polypeptide/compound complex is detected, a compound that binds a polypeptide of the invention is identified.

Compounds identified via such methods can include compounds which modulate the activity of a polypeptide of the invention (that is, increase or decrease its activity, relative to activity observed in the absence of the compound). Alternatively, compounds identified via such methods can include compounds which modulate the expression of a polynucleotide of the invention (that is, increase or decrease expression relative to expression levels observed in the absence of the compound). Compounds, such as compounds identified via the methods of the invention, can be tested using standard assays well known to those of skill in the art for their ability to modulate activity/expression.

The agents screened in the above assay can be, but are not limited to, peptides, carbohydrates, vitamin derivatives, or other pharmaceutical agents. The agents can be selected and screened at random or rationally selected or designed using protein modeling techniques.

For random screening, agents such as peptides, carbohydrates, pharmaceutical agents and the like are selected at random and are assayed for their ability to bind to the protein encoded by the ORF of the present invention. Alternatively, agents may be rationally selected or designed. As used herein, an agent is said to be "rationally selected or designed" when the agent is chosen based on the configuration of the particular protein. For example, one skilled in the art can readily adapt currently available procedures to generate peptides, pharmaceutical agents and the like, capable of binding to a specific peptide sequence, in order to generate rationally designed antipeptide peptides, for example see Hurby et al., Application of Synthetic Peptides: Antisense Peptides," In Synthetic Peptides, A User's Guide, W.H. Freeman, NY (1992), pp. 289-307, and Kasprzak et al., Biochemistry 28:9230-8 (1989), or pharmaceutical agents, or the like.

In addition to the foregoing, one class of agents of the present invention, as broadly described, can be used to control gene expression through binding to one of the ORFs or EMFs of the present invention. As described above, such agents can be randomly screened or rationally designed/selected. Targeting the ORF or EMF allows a skilled artisan to design sequence specific or element specific agents, modulating the expression of either a single ORF or multiple ORFs which rely on the same EMF for expression control. One class of DNA binding agents are agents which contain base residues which hybridize or form a triple helix formation by binding to DNA or RNA. Such agents can be based on the classic phosphodiester, ribonucleic acid backbone, or can be a variety of sulfhydryl or polymetric derivatives which have base attachment capacity.

Agents suitable for use in these methods usually contain 20 to 40 bases and are designed to be complementary to a region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1360 (1991)) or to the mRNA itself (antisense - Okano, J. Neurochem. 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). Triple helix-formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques have been demonstrated to be effective in model systems. Information contained in the sequences of the present

invention is necessary for the design of an antisense or triple helix oligonucleotide and other DNA binding agents.

Agents which bind to a protein encoded by one of the ORFs of the present invention can be used as a diagnostic agent. Agents which bind to a protein encoded by one of the ORFs of the present invention can be formulated using known techniques to generate a pharmaceutical composition.

## 5.19 USE OF NUCLEIC ACIDS AS PROBES

Another aspect of the subject invention is to provide for polypeptide-specific nucleic acid hybridization probes capable of hybridizing with naturally occurring nucleotide sequences. The hybridization probes of the subject invention may be derived from any of the nucleotide sequences SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62. Because the corresponding gene is only expressed in a limited number of tissues, a hybridization probe derived from any of the nucleotide sequences SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 can be used as an indicator of the presence of RNA of cell type of such a tissue in a sample.

Any suitable hybridization technique can be employed, such as, for example, *in situ* hybridization. PCR as described in US Patents Nos. 4,683,195 and 4,965,188 provides additional uses for oligonucleotides based upon the nucleotide sequences. Such probes used in PCR may be of recombinant origin, may be chemically synthesized, or a mixture of both. The probe will comprise a discrete nucleotide sequence for the detection of identical sequences or a degenerate pool of possible sequences for identification of closely related genomic sequences.

Other means for producing specific hybridization probes for nucleic acids include the cloning of nucleic acid sequences into vectors for the production of mRNA probes. Such vectors are known in the art and are commercially available and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerase as T7 or SP6 RNA polymerase and the appropriate radioactively labeled nucleotides. The nucleotide sequences may be used to construct hybridization probes for mapping their respective genomic sequences. The nucleotide sequence provided herein may be mapped to a chromosomal or specific regions of a chromosome using well known genetic and/or chromosomal mapping techniques. These techniques include *in situ* hybridization, linkage analysis against known chromosomal markers, hybridization screening with libraries or flow-sorted chromosomal preparations specific to known chromosomes, and the like. The technique of fluorescent *in situ* hybridization of chromosome spreads has been described, among other places, in Verma et

al (1988) Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York NY.

Fluorescent *in situ* hybridization of chromosomal preparations and other physical chromosome mapping techniques may be correlated with additional genetic map data.

5 Examples of genetic map data can be found in the 1994 Genome Issue of Science (265:1981f). Correlation between the location of a nucleic acid on a physical chromosomal map and a specific disease (or predisposition to a specific disease) may help delimit the region of DNA associated with that genetic disease. The nucleotide sequences of the subject invention may be used to detect differences in gene sequences between normal, carrier or affected individuals.

## 10 5.20 PREPARATION OF SUPPORT BOUND OLIGONUCLEOTIDES

Oligonucleotides, i.e., small nucleic acid segments, may be readily prepared by, for example, directly synthesizing the oligonucleotide by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer.

Support bound oligonucleotides may be prepared by any of the methods known to those of skill in the art using any suitable support such as glass, polystyrene or Teflon. One strategy is to precisely spot oligonucleotides synthesized by standard synthesizers. Immobilization can be achieved using passive adsorption (Inouye & Hondo, 1990 J. Clin Microbiol 28(6) 1462-72); using UV light (Nagata *et al.*, 1985; Dahlen *et al.*, 1987; Morrissey & Collins, Mol. Cell Probes 1989 3(2) 189-207) or by covalent binding of base modified DNA (Keller *et al.*, 1988; 1989); all references being specifically incorporated herein.

Another strategy that may be employed is the use of the strong biotin-streptavidin interaction as a linker. For example, Broude *et al.* (1994) Proc. Natl. Acad. Sci USA 91(8) 3072-6 describe the use of biotinylated probes, although these are duplex probes, that are immobilized on streptavidin-coated magnetic beads. Streptavidin-coated beads may be purchased from Dynal, Oslo. Of course, this same linking chemistry is applicable to coating any surface with streptavidin. Biotinylated probes may be purchased from various sources, such as, e.g., Operon Technologies (Alameda, CA).

Nunc Laboratories (Naperville, IL) is also selling suitable material that could be used.

Nunc Laboratories have developed a method by which DNA can be covalently bound to the microwell surface termed Covalink NH. Covalink NH is a polystyrene surface grafted with secondary amino groups (> NH) that serve as bridge-heads for further covalent coupling.

Covalink Modules may be purchased from Nunc Laboratories. DNA molecules may be bound

to CovaLink exclusively at the 5'-end by a phosphoramidate bond, allowing immobilization of more than 1 pmol of DNA (Rasmussen *et al.*, (1991) Anal Biochem 198(1) 138-42.

The use of CovaLink NH strips for covalent binding of DNA molecules at the 5'-end has been described (Rasmussen *et al.*, 1991). In this technology, a phosphoramidate bond is employed (Chui *et al.*, 1983 Nucleic Acids 11(18) 6513-29). This is beneficial as immobilization using only a single covalent bond is preferred. The phosphoramidate bond joins the DNA to the CovaLink NH secondary amino groups that are positioned at the end of spacer arms covalently grafted onto the polystyrene surface through a 2 nm long spacer arm. To link an oligonucleotide to CovaLink NH via an phosphoramidate bond, the oligonucleotide terminus must have a 5'-end phosphate group. It is, perhaps, even possible for biotin to be covalently bound to CovaLink and then streptavidin used to bind the probes.

More specifically, the linkage method includes dissolving DNA in water (7.5 ng/ul) and denaturing for 10 min. at 95°C and cooling on ice for 10 min. Ice-cold 0.1 M 1-methylimidazole, pH 7.0 (1-Melm), is then added to a final concentration of 10 mM 1-Melm. A ss DNA solution is then dispensed into CovaLink NH strips (75 ul/well) standing on ice.

Carbodiimide 0.2 M 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide(EDC), dissolved in 10 mM 1-Melm, is made fresh and 25 ul added per well. The strips are incubated for 5 hours at 50°C. After incubation the strips are washed using, e.g., Nunc-Immuno Wash; first the wells are washed 3 times, then they are soaked with washing solution for 5 min., and finally they are washed 3 times (where in the washing solution is 0.4 N NaOH, 0.25% SDS heated to 50°C).

It is contemplated that a further suitable method for use with the present invention is that described in PCT Patent Application WO 90/03382 (Southern & Maskos), incorporated herein by reference. This method of preparing an oligonucleotide bound to a support involves attaching a nucleoside 3'-reagent through the phosphate group by a covalent phosphodiester link to aliphatic hydroxyl groups carried by the support. The oligonucleotide is then synthesized on the supported nucleoside and protecting groups removed from the synthetic oligonucleotide chain under standard conditions that do not cleave the oligonucleotide from the support. Suitable reagents include nucleoside phosphoramidite and nucleoside hydrogen phosphorate.

An on-chip strategy for the preparation of DNA probe for the preparation of DNA probe arrays may be employed. For example, addressable laser-activated photodeprotection may be employed in the chemical synthesis of oligonucleotides directly on a glass surface, as described by Fodor *et al.* (1991) Science 251(4995) 767-73, incorporated herein by reference. Probes may also be immobilized on nylon supports as described by Van Ness *et al.* (1991) Nucleic Acids Res.

19(2) 3345-50; or linked to Teflon using the method of Durcan & Cavalier (1988) Anal Biochem 169(1) 104-8; all references being specifically incorporated herein.

To link an oligonucleotide to a nylon support, as described by Van Ness *et al.* (1991), requires activation of the nylon surface via alkylation and selective activation of the 5'-amine of oligonucleotides with cyanuric chloride.

One particular way to prepare support bound oligonucleotides is to utilize the light-generated synthesis described by Pease *et al.*, (1994) Proc. Natl. Acad. Sci USA 91(11) 5022-6. These authors used current photolithographic techniques to generate arrays of immobilized oligonucleotide probes (DNA chips). These methods, in which light is used to direct the synthesis of oligonucleotide probes in high-density, miniaturized arrays, utilize photolabile 5'-protected N-acyl-deoxynucleoside phosphoramidites, surface linker chemistry and versatile combinatorial synthesis strategies. A matrix of 256 spatially defined oligonucleotide probes may be generated in this manner.

## 5.21 PREPARATION OF NUCLEIC ACID FRAGMENTS

The nucleic acids may be obtained from any appropriate source, such as cDNAs, genomic DNA, chromosomal DNA, microdissected chromosome bands, cosmid or YAC inserts, and RNA, including mRNA without any amplification steps. For example, Sambrook *et al.* (1989) describes three protocols for the isolation of high molecular weight DNA from mammalian cells (p. 9.14-9.23).

DNA fragments may be prepared as clones in M13, plasmid or lambda vectors and/or prepared directly from genomic DNA or cDNA by PCR or other amplification methods. Samples may be prepared or dispensed in multiwell plates. About 100-1000 ng of DNA samples may be prepared in 2-500 ml of final volume.

The nucleic acids would then be fragmented by any of the methods known to those of skill in the art including, for example, using restriction enzymes as described at 9.24-9.28 of Sambrook *et al.* (1989), shearing by ultrasound and NaOH treatment.

Low pressure shearing is also appropriate, as described by Schriefer *et al.* (1990) Nucleic Acids Res. 18(24) 7455-6. In this method, DNA samples are passed through a small French pressure cell at a variety of low to intermediate pressures. A lever device allows controlled application of low to intermediate pressures to the cell. The results of these studies indicate that low-pressure shearing is a useful alternative to sonic and enzymatic DNA fragmentation methods.

One particularly suitable way for fragmenting DNA is contemplated to be that using the two base recognition endonuclease, CviII, described by Fitzgerald *et al.* (1992) Nucleic Acids

Res. 20(14) 3753-62. These authors described an approach for the rapid fragmentation and fractionation of DNA into particular sizes that they contemplated to be suitable for shotgun cloning and sequencing.

The restriction endonuclease *Cvi*II normally cleaves the recognition sequence PuGCPy between the G and C to leave blunt ends. Atypical reaction conditions, which alter the specificity of this enzyme (*Cvi*II\*\*), yield a quasi-random distribution of DNA fragments from the small molecule pUC19 (2688 base pairs). Fitzgerald *et al.* (1992) quantitatively evaluated the randomness of this fragmentation strategy, using a *Cvi*II\*\* digest of pUC19 that was size fractionated by a rapid gel filtration method and directly ligated, without end repair, to a lac Z minus M13 cloning vector. Sequence analysis of 76 clones showed that *Cvi*II\*\* restricts pyGCPy and PuGCPu, in addition to PuGCPy sites, and that new sequence data is accumulated at a rate consistent with random fragmentation.

As reported in the literature, advantages of this approach compared to sonication and agarose gel fractionation include: smaller amounts of DNA are required (0.2-0.5 ug instead of 2-5 ug); and fewer steps are involved (no preligation, end repair, chemical extraction, or agarose gel electrophoresis and elution are needed).

Irrespective of the manner in which the nucleic acid fragments are obtained or prepared, it is important to denature the DNA to give single stranded pieces available for hybridization. This is achieved by incubating the DNA solution for 2-5 minutes at 80-90°C. The solution is then cooled quickly to 2°C to prevent renaturation of the DNA fragments before they are contacted with the chip. Phosphate groups must also be removed from genomic DNA by methods known in the art.

## 5.22 PREPARATION OF DNA ARRAYS

Arrays may be prepared by spotting DNA samples on a support such as a nylon membrane. Spotting may be performed by using arrays of metal pins (the positions of which correspond to an array of wells in a microtiter plate) to repeated by transfer of about 20 nl of a DNA solution to a nylon membrane. By offset printing, a density of dots higher than the density of the wells is achieved. One to 25 dots may be accommodated in 1 mm<sup>2</sup>, depending on the type of label used. By avoiding spotting in some preselected number of rows and columns, separate subsets (subarrays) may be formed. Samples in one subarray may be the same genomic segment of DNA (or the same gene) from different individuals, or may be different, overlapped genomic clones. Each of the subarrays may represent replica spotting of the same samples. In one example, a selected gene segment may be amplified from 64 patients. For each patient, the

amplified gene segment may be in one 96-well plate (all 96 wells containing the same sample). A plate for each of the 64 patients is prepared. By using a 96-pin device, all samples may be spotted on one 8 x 12 cm membrane. Subarrays may contain 64 samples, one from each patient. Where the 96 subarrays are identical, the dot span may be 1 mm<sup>2</sup> and there may be a 1 mm space between subarrays.

Another approach is to use membranes or plates (available from NUNC, Naperville, Illinois) which may be partitioned by physical spacers e.g. a plastic grid molded over the membrane, the grid being similar to the sort of membrane applied to the bottom of multiwell plates, or hydrophobic strips. A fixed physical spacer is not preferred for imaging by exposure to flat phosphor-storage screens or x-ray films.

The present invention is illustrated in the following examples. Upon consideration of the present disclosure, one of skill in the art will appreciate that many other embodiments and variations may be made in the scope of the present invention. Accordingly, it is intended that the broader aspects of the present invention not be limited to the disclosure of the following examples. The present invention is not to be limited in scope by the exemplified embodiments which are intended as illustrations of single aspects of the invention, and compositions and methods which are functionally equivalent are within the scope of the invention. Indeed, numerous modifications and variations in the practice of the invention are expected to occur to those skilled in the art upon consideration of the present preferred embodiments. Consequently, the only limitations which should be placed upon the scope of the invention are those which appear in the appended claims.

All references cited within the body of the instant specification are hereby incorporated by reference in their entirety.

## 6.0 EXAMPLES

### EXAMPLE 1

#### Isolation of SEQ ID NO: 1, 10, 17, 26, and 33 from cDNA Libraries

A plurality of novel nucleic acids were obtained from cDNA libraries prepared from human leukocyte mRNA (GIBCO Laboratories) (SEQ ID NO: 1), from human adult liver mRNA (GIBCO) (SEQ ID NO: 10); from human adult kidney mRNA (GIBCO) (SEQ ID NO: 17); from human adult brain mRNA (GIBCO) (SEQ ID NO: 26) and from human adult kidney mRNA (Invitrogen) (SEQ ID NO: 33) using standard PCR, sequencing by hybridization sequence signature analysis, and Sanger sequencing techniques. The inserts of the libraries

were amplified with PCR using primers specific for vector sequences flanking the inserts.

These samples were spotted onto nylon membranes and interrogated with oligonucleotide probes to give sequence signatures. The clones were clustered into groups of similar or identical sequences, and single representative clones were selected from each group for gel sequencing. The 5' sequence of the amplified inserts was then deduced using the reverse M13 sequencing primer in a typical Sanger sequencing protocol. PCR products were purified and subjected to fluorescent dye terminator cycle sequencing. Single-pass gel sequencing was done using a 377 Applied Biosystems (ABI) sequencer. The insert was identified as a novel sequence not previously obtained from this library and not previously reported in public databases. These sequences were designated as SEQ ID NO: 1, 10, 17, 26, and 33.

#### EXAMPLE 2

##### ASSEMBLAGE OF SEQ ID NO: 2, 11, 18, 27, 34, 60, and 62

The nucleic acids of the present invention, designated as SEQ ID NO: 2, 11, 18, 27, and 34 were assembled using SEQ ID NO: 1, 10, 17, 26, and 33 as a seed, respectively. Then a recursive algorithm was used to extend the seed into an extended assemblage, by pulling additional sequences from different databases (i.e., Hyseq's database containing EST sequences, dbEST version 114, gb pri 114, and UniGene version 101) that belong to this assemblage. The algorithm terminated when there was no additional sequences from the above databases that would extend the assemblage. Inclusion of component sequences into the assemblage was based on a BLASTN hit to the extending assemblage with BLAST score greater than 300 and percent identity greater than 95%.

The nearest neighbor result for the assembled contigs were obtained by a FASTA version 3 search against Genpept release 114, 117, or 118 using FASTXY algorithm. FASTXY is an improved version of FASTA alignment which allows in-codon frame shifts. The nearest neighbor result showed the closest homologue for each assemblage from Genpept (and contains the translated amino acid sequences for which the assemblage encodes). The nearest neighbor results are set forth below:

SEQ ID NO:	Accession No.	Description	Smith-Waterman Score	% Identity
2	AF151799	Homo sapiens CGI-40 protein	424	71.605

11	AC005053	Homo sapiens match to ESTs AA316181 (NID: g3165221), AA032221 (NID: g1502183), and A1167942 (NID: g3701112)	881	49,580
18	AB018301	Homo sapiens KIAA0758 protein	6455	99,595
27	AF027826	Homo sapiens putative seven pass transmembrane protein	487	46,691
34	AC005587	Homo sapiens similar to mouse olfactory receptor 13, similar to P34984 (PID: g464305)	1450	71,382

Polypeptides were predicted to be encoded by SEQ ID NO: 2, 11, 18, 27, and 34 as set forth below. The polypeptides were predicted using a software program called FASTY (available from <http://fasta.bioch.virginia.edu>) which selects a polypeptide based on a comparison of translated novel polynucleotide to known polypeptides (W.R. Pearson, Methods in Enzymology, 183: 63-98 (1990), herein incorporated by reference).

Predicted beginning nucleotide location	Predicted end nucleotide location	Amino acid composition of the polypeptide encoded, wherein, (A=Alanine, C=Cysteine, D=Aspartic Acid, E= Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop Codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
66	453	VGEPYIDWDEFFELLRSRTAVRARKIPISDI* KTK AKQVVKLLSNRSQAVGILMSSLHLDKMDIQHA VYNLDNSVVDLETQLQALYENRAQSQDELE*IEKHG

		RSSKDKENAKSLDKPEQLYFLRLYE (SEQ ID NO: 9)
23	462	ARGGRLRWRLDDCLSAAESDTVA YEDLSEDTY QKKWKGLALSQRALHWNMLENDRSMASLAG RNMMESSELTPKQEIFKGSESSNSTSGGLFGVVP GGTETGDVCEDTFKELEGQPSNEEGSRLESDFLEI IDEDKKKSTKDRY (SEQ ID NO: 16)
166	4280	ASDQSGSQPGDHSAGQANQKLEDKMSPRRTTL CLMFITYSSKAALNWNYESITHPLSLHEHEPAGE EALRQKRAVATKSPTABEYTVNIEISFENASFLDP IKAYLSLSPFHGNNTDQITDILSINVTTCRPAG NEIWCSCETGYGWPRERCLHNLICQERDVFLPGH HCSCCLKELPPNGPCLLQEDVTLNMRVRLNVGF QEDLMNTSSALYRSYKTDLETAFRKGYGILPGFK GVTVTGFKSGSVVTVYEKTTTPPSLELIHKANEQ VVQSLNQTYKMDYNSFQAVTINESNFFVTPEIIFE GDTVSLVCEKEVLSNVSWRYEEQQLIQNSSRF SYTALFNMMTSVSKLTHNTTPGDAGEYVCKLIL DIFEYCKKKIDVMPQILANEEMKVMCDNNPVS LNCCSQGNVNWSKVEWKQEGKINPGTPTDIDS SCSRYTLKADGTQCPSGSGTTVIYTCFISAYGA RGSANIKVTFISVANLITTPDPISVSEQNFSIKCIS DVSNYDEVYWNWTSAGIKIYQRFYTRRYLDGAE SVLTVKSTSTREWNGTYHCIFRYKNSYSIATKDVI VHPLPLKLNIMVDPLEATVSCSGSHHIKCCIEEDG DYKVTFFHMGSSSLPAAKEVNVKKQVCYKHNFNA SSVSWCSKTVDVCCFTNAANNSVWSPSMKLN VPGENITCQDPVIGVGEFGKVIQKLCRFNSVPSSP EE/SPLGGTITYKCVGSQWGIEKRNDICISAPNSLL QMAKALIKSPSQDEMLPTYLKDLISIDKAEHEIS SSPGSLGAIJNILDLLSTVPTQVNSEMMTHVLSTV NVILGKPVNLTWKVLQQQWNTNQSSQLLHVSVERF SQALQSGDSPPLSFQNTNVQMSSTVIKSSHPETYQ QRVFPYFDLWGNVVIDKSYLENLQSDSSIVTMA FPRTLQAILAQDIQENNFASLVMITTVSHNTTMP

		FRISMFTKNNSPSGGETKCVFWNFRLANNTGOW DSSGCVVEEGDGDNVTCICDHLTFSILMSPDSDPD PSSLLGILLDIISYVGVGFSILSLAACLVEAVVW KSVTKNRTSYMRHTCIVNIAASLLVANTWFIGV AAIQDNRYLCKTACVAAITFFIHFFYLSVFFWML TLGLMLFYRLVFILHETSRSTQKAIACGLGYGCPL AISVITLGATQPREVYTRKNVCWLNWEDTKALL AFAIPALIVVNNITITVITKILRPSIGDKPKQEK SSLFQISKSIGVLTPLLGLTWGFGLTTPFPGTNLV FHIIPAILNVFQGLFILLFGCLWDLKVQEALLNKF SLSRWSSQHSKSTSLGSSTPVFSMSSPSRRFNLF GKTGTYNVSTPEATSSSLENSSSASSLLN (SEQ ID NO: 25)
1009	1208	VRGLGPRLPVFPKGLSVBEGGLSATTSFLLSA PSPSLHPAIPTRIRIYFPGADSPSLV/SRDSGLPPL TWRVTCLOLVACLPGLPALPPAVTLGLTAAYT TLYALLFFSVYAQLWLVLARMGHKRLSYQTVFL ALCLFVWAPLRTTTFST*FPKILPAPNN/SWGPLPF WLLYCCPVCLQFFTLTLMNLVYFAQVVFKA/KSE ASGPKMSRGLLAVRGAFVGASLLFLLVNVLCAV LVPCGAAAQPWALLVRVLVSDSLFVICALSLA ACLFLCRQAGALH*HLPGGQGRAAAALMPRCLLG LSAAVLRV*RTAERPKRHLGISAAAALPWPGRG (SEQ ID NO: 32)
1206	2266	RHLLTIFHKLKIYKTIKIDFKKRVTVQLLVFLF LCLFFSSEMVKNQTMVTEFLLGLLGPRIQMLL FGLFSLFYVFTLLGNGTILGLISLDSRLHTPMYFFL SHLAVVNIAYACNTVPQMLVNLHPAKPISFAGC MT*TEFLSFAHTECLLLVLMYSDRYVAICHPLR YFIIMTWKVCITLAITSWTCGSLLAMVHVSLILRL PFCGPREINHFFCEILSVLRACADTWNQVIFA ACMFILVQPLCLVLVSYSHILAAILRIQSGEGRRK AFSTCSSHLCVVGLFFGSAIVMYMAPKSRHPEEQ QKVLFLEYSSFPMLNPLVNLNRNVEVKGALRRA LCKESH (SEQ ID NO: 47)

## EXAMPLE 3

## ASSEMBLAGE OF SEQ ID NO: 4, 13, 20, 29, 36, or 42

Using PHRAP (Univ. of Washington), full-length gene cDNA sequences and its corresponding protein sequences were generated from the assemblage of SEQ ID NO: 2, 11, 18, 27, and 34. Any frame shifts and incorrect stop codons were corrected by hand editing. During editing, the sequence was checked using FASTY and/or BLAST against Genbank. (i.e. Genpept release 117 or 119). Other computer programs, which may have been used in the editing process, were phredPhrap and Consed (University of Washington) and ed-ready, ed-ext and cg-zip-2 (Hyseq, Inc.).

A polypeptide (SEQ ID NO: 4) was predicted to be encoded by SEQ ID NO: 3 as set forth below. The polypeptide was predicted using a software program called BLASTX which selects a polypeptide based on a comparison of translated novel polynucleotide to known polynucleotides. The initial methionine starts at position 1 of SEQ ID NO: 3 and the putative stop codon, TGA, begins at position 2482 of the nucleotide sequence.

The GPCR-like polypeptide of SEQ ID NO: 4 is an approximately 827-amino acid transmembrane protein with a predicted molecular mass of approximately 93-kDa unglycosylated. Hyseq's sequence database searches using the Pfam models that were categorized under G protein-coupled receptors and using the hmsearch program (hmsearch - search a sequence database with a profile HMM: HMMER 2.1.1 (Dec 1998) Washington University School of Medicine), SEQ ID NO 4 was found to be homologous to G protein-coupled receptor model sequences with an E-value of 0.011. The homologous sequence identified using Pfam hmsearch is shown in SEQ ID NO: 6. Further analyses with protein database searches with the BLASTP algorithm (Altschul S.F. et al., J. Mol. Evol. 36:290-300 (1993) and Altschul S.F. et al., J. Mol. Biol. 21:403-10 (1990), herein incorporated by reference) indicate that SEQ ID NO: 4 is homologous to human CGI-40 protein and with protein of clone CT748\_2.

Figure 1 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 3 (i.e. SEQ ID NO: 4) GPCR-like polypeptide and human CGI-40 protein (Lai et al. (2000) Genome Res. 10, 703-713) (SEQ ID NO: 48), indicating that the two

sequences share 76% similarity over 526 amino acid residues and 63% identity over the same 526 amino acid residues.

Figure 2 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 3 (i.e. SEQ ID NO: 4) GPCR-like polypeptide and protein of clone CT748\_2 (Patent Application No. WO9824905) (SEQ ID NO: 49), indicating that the two sequences share 97% similarity over 445 amino acid residues and 96% identity over the same 445 amino acid residues.

A predicted approximately nineteen-residue signal peptide is encoded from approximately residue 1 through residue 19 of SEQ ID NO: 4 (SEQ ID NO: 7). The extracellular portion is useful on its own. This can be confirmed by expression in mammalian cells and sequencing of the cleaved product. The signal peptide region was predicted using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). One of skill in the art will recognize that the actual cleavage site may be different than that predicted by the computer program.

A polypeptide (SEQ ID NO: 13) was predicted to be encoded by SEQ ID NO: 12 as set forth below. The polypeptide was predicted using a software program called BLASTX which selects a polypeptide based on a comparison of translated novel polynucleotide to known polynucleotides. The initial methionine starts at position 135 of SEQ ID NO: 12 and the putative stop codon, TGA, begins at position 1599 of the nucleotide sequence.

The GPCR-like polypeptide of SEQ ID NO: 13 is an approximately 488-amino acid transmembrane protein with a predicted molecular mass of approximately 55-kDa unglycosylated. Hyseq's sequence database searches using the Pfam models that were categorized under G protein-coupled receptors and using the hmsearch program (hmsearch - search a sequence database with a profile HMM: HMMER 2.1.1 (Dec 1998) Washington University School of Medicine), SEQ ID NO 13 was found to be homologous to G protein-coupled receptor model sequences with an E-value of 0.017. The homologous sequence identified using Pfam hmsearch is shown in SEQ ID NO: 15. Protein database searches with the BLASTP algorithm (Altschul S.F. et al., J. Mol. Evol. 36:290-300 (1993) and Altschul S.F. et al., J. Mol. Biol. 21:403-10 (1990), herein incorporated by reference) indicate that SEQ ID NO: 4 is homologous to six transmembrane epithelial antigen of prostate and with human STRAP-1 protein.

Figure 3 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 12 (i.e. SEQ ID NO: 13) GPCR-like polypeptide and human six transmembrane epithelial antigen of prostate (Hubert et al. (1999) Proc. Natl. Acad. Sci.



U.S.A. 96, 14523-14528) (SEQ ID NO: 50), indicating that the two sequences share 68 % similarity over 267 amino acid residues and 47 % identity over the same 267 amino acid residues.

Figure 4 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 12 (i.e. SEQ ID NO: 13) GPCR-like polypeptide and human STRAP-1 protein (Patent Application No. WO9962941) (SEQ ID NO: 51), indicating that the two sequences share 68 % similarity over 267 amino acid residues and 47 % identity over the same 267 amino acid residues.

A polypeptide (SEQ ID NO: 20) was predicted to be encoded by SEQ ID NO: 19 as set forth below. The polypeptide was predicted using a software program called BLASTX which selects a polypeptide based on a comparison of translated novel polynucleotide to known polynucleotides. The initial methionine starts at position 272 of SEQ ID NO: 19 and the putative stop codon, TAA, begins at position 4310 of the nucleotide sequence.

The GPCR-like polypeptide of SEQ ID NO: 20 is an approximately 1346-amino acid transmembrane protein with a predicted molecular mass of approximately 151-kDa unglycosylated. Hyseq's sequence database searches using the Pfam models that were categorized under G protein-coupled receptors and using the hmsearch program (hmsearch - search a sequence database with a profile HMM: HMMER 2.1.1 (Dec 1998) Washington University School of Medicine), SEQ ID NO 20 was found to be homologous to G protein-coupled receptor model sequences with an E-value of 2.7e-24. The homologous sequence identified using Pfam hmsearch is shown in SEQ ID NO: 22. Further analyses with protein database searches with the BLASTP algorithm (Altschul S.F. et al., J. Mol. Evol. 36:290-300 (1993) and Altschul S.F. et al., J. Mol. Biol. 21:403-10 (1990), herein incorporated by reference) indicate that SEQ ID NO: 20 is homologous to the rat seven transmembrane receptor and to the human brain-derived G protein-coupled receptor proteins.

Figure 5A, 5B, and 5C show the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 19 (i.e. SEQ ID NO: 20) GPCR-like polypeptide and rat seven transmembrane receptor protein (Abe et al, (1999) J. Biol. Chem. 274, 19957-19964) (SEQ ID NO: 52), indicating that the two sequences share 81 % similarity over 1354 amino acid residues and 72 % identity over the same 1354 amino acid residues.

Figure 6A, and 6B show the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 19 (i.e. SEQ ID NO: 20) GPCR-like polypeptide and human brain-derived G protein-coupled receptor protein (Patent Application No. WO200008053) (SEQ ID

NO: 53), indicating that the two sequences share 100 % similarity over 986 amino acid residues and 100 % identity over the same 986 amino acid residues.

A predicted approximately twenty one-residue signal peptide is encoded from approximately residue 1 through residue 21 of SEQ ID NO: 20 (SEQ ID NO: 23). The extracellular portion is useful on its own. This can be confirmed by expression in mammalian cells and sequencing of the cleaved product. The signal peptide region was predicted using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). One of skill in the art will recognize that the actual cleavage site may be different than that predicted by the computer program.

SEQ ID NO: 60 is very similar to SEQ ID NO: 19. A polypeptide (SEQ ID NO: 61) was predicted to be encoded by SEQ ID NO: 60. The initial methionine starts at 272 of SEQ ID NO: 60 and the putative stop codon begins at position 4310.

A polypeptide (SEQ ID NO: 29) was predicted to be encoded by SEQ ID NO: 28 as set forth below. The polypeptide was predicted using a software program called BLASTX which selects a polypeptide based on a comparison of translated novel polynucleotide to known polynucleotides. The initial methionine starts at position 52 of SEQ ID NO: 28 and the putative stop codon, TAA, begins at position 3994 of the nucleotide sequence.

The GPCR-like polypeptide of SEQ ID NO: 29 is an approximately 1314-amino acid transmembrane protein with a predicted molecular mass of approximately 147-kDa unglycosylated. Hyseq's sequence database searches with the Pfam models that were categorized under G protein-coupled receptors using the hmsearch program (hmsearch - search a sequence database with a profile HMM: HMMER 2.1.1 (Dec 1998) Washington University School of Medicine), SEQ ID NO 29 was found to be homologous to G protein-coupled receptor model sequences with an E-value of 0.0036. The homologous sequence identified using Pfam hmsearch is shown in SEQ ID NO: 31. Further analyses with protein database searches with the BLASTP algorithm (Altschul S.F. et al., J. Mol. Evol. 36:290-300 (1993) and Altschul S.F. et al., J. Mol. Biol. 21:403-10 (1990), herein incorporated by reference) indicate that SEQ ID NO: 29 is homologous to the putative seven pass transmembrane protein and to the human h-TRAAK polypeptide #1.

Figure 7 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 28 (i.e. SEQ ID NO: 29) GPCR-like polypeptide and putative seven pass transmembrane protein (Spangenberg et al (1998) Genomics 48, 178-185) (SEQ ID NO: 54), indicating that the two sequences share 72 % similarity over 323 amino acid residues and 57 % identity over the same 323 amino acid residues.

Figure 8 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 28 (i.e. SEQ ID NO: 29) GPCR-like polypeptide and human h-TRAAK polypeptide #1 (Patent Application No. WO200026253) (SEQ ID NO: 55), indicating that the two sequences share 100% similarity over 392 amino acid residues and 100% identity over the same 392 amino acid residues.

A polypeptide (SEQ ID NO: 36) was predicted to be encoded by SEQ ID NO: 35 as set forth below. The polypeptide was predicted using a software program called BLASTX which selects a polypeptide based on a comparison of translated novel polynucleotide to known polynucleotides. The initial methionine starts at position 833 of SEQ ID NO: 35 and the putative stop codon, TAA, begins at position 1415 of the nucleotide sequence.

The GPCR-like polypeptide of SEQ ID NO: 36 is an approximately 194-amino acid transmembrane protein with a predicted molecular mass of approximately 22-kDa unglycosylated. Hyseq's sequence database searches using the Pfam models that were categorized under G protein-coupled receptors and using the hmmsrch program (hmmsrch - search a sequence database with a profile HMM: HMMER 2.1.1 (Dec 1998) Washington University School of Medicine), SEQ ID NO 36 was found to be homologous to G protein-coupled receptor model sequences with an E-value of 1.8e-28. The homologous sequence identified using Pfam hmmsrch is shown in SEQ ID NO: 38. Further analyses with protein database searches with the BLASTP algorithm (Altschul S.F. et al., J. Mol. Evol. 36:290-300 (1993) and Altschul S.F. et al., J. Mol. Biol. 21:403-10 (1990), herein incorporated by reference) indicate that SEQ ID NO: 36 is homologous to the human olfactory receptor protein and to the human G protein-coupled receptor GPR1 polypeptide.

Figure 9 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 35 (i.e. SEQ ID NO: 36) GPCR-like polypeptide and human olfactory receptor (Rouquier et al. (1998) Nature Genet. 18 (3), 243-250) (SEQ ID NO: 56), indicating that the two sequences share 92% similarity over 166 amino acid residues and 87% identity over the same 166 amino acid residues.

Figure 10 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 35 (i.e. SEQ ID NO: 36) GPCR-like polypeptide and human G protein-coupled receptor GPR1 protein (Patent Application No. WO9630406) (SEQ ID NO: 57), indicating that the two sequences share 93% similarity over 171 amino acid residues of and 92% identity over the same 171 amino acid residues.

A predicted approximately thirty five-residue signal peptide is encoded from approximately residue 1 through residue 35 of SEQ ID NO: 36 (SEQ ID NO: 39). The extracellular portion

is useful on its own. This can be confirmed by expression in mammalian cells and sequencing of the cleaved product. The signal peptide region was predicted using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). One of skill in the art will recognize that the actual cleavage site may be different than that predicted by the computer program.

A polypeptide (SEQ ID NO: 42) was predicted to be encoded by SEQ ID NO: 41 as set forth below. The polypeptide was predicted using a software program called BLASTX which selects a polypeptide based on a comparison of translated novel polynucleotide to known polynucleotides. The initial methionine starts at position 485 of SEQ ID NO: 41 and the putative stop codon, TAA, begins at position 1409 of the nucleotide sequence.

The GPCR-like polypeptide of SEQ ID NO: 42 is an approximately 308-amino acid transmembrane protein with a predicted molecular mass of approximately 34-kDa unglycosylated. Hyseq's sequence database searches using the Pfam models that were categorized under G protein-coupled receptors and using the hmmsrch program (hmmsrch - search a sequence database with a profile HMM: HMMER 2.1.1 (Dec 1998) Washington University School of Medicine), SEQ ID NO 42 was found to be homologous to G protein-coupled receptor model sequences with an E-value of 1.1e-47. The homologous sequence identified using Pfam hmmsrch is shown in SEQ ID NO: 44. Further analyses with protein database searches with the BLASTP algorithm (Altschul S.F. et al., J. Mol. Evol. 36:290-300 (1993) and Altschul S.F. et al., J. Mol. Biol. 21:403-10 (1990), herein incorporated by reference) indicate that SEQ ID NO: 42 is homologous to the mouse olfactory receptor 13 polypeptide and to the human G protein-coupled receptor GPR1 polypeptide.

Figure 11 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 41 (i.e. SEQ ID NO: 42) GPCR-like polypeptide and human gene ACO05587, similar to mouse olfactory receptor 13 polypeptide (SEQ ID NO: 58), indicating that the two sequences share 81% similarity over 304 amino acid residues and 68% identity over the same 304 amino acid residues.

Figure 12 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO: 41 (i.e. SEQ ID NO: 42) GPCR-like polypeptide and human G protein-coupled receptor GPR1 polypeptide (Patent Application No. WO9630406) (SEQ ID NO: 59), indicating that the two sequences share 91% similarity over 287 amino acid residues and 90% identity over the same 287 amino acid residues.

A predicted approximately forty two-residue signal peptide is encoded from approximately residue 1 through residue 42 of SEQ ID NO: 42 (SEQ ID NO: 45). The extracellular portion

is useful on its own. This can be confirmed by expression in mammalian cells and sequencing of the cleaved product. The signal peptide region was predicted using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). One of skill in the art will recognize that the actual cleavage site may be different than that predicted by the computer program.

5

SEQ ID NO: 62 is similar to SEQ ID NO: 35 and 41. A polypeptide (SEQ ID NO: 63) was predicted to be encoded by SEQ ID NO: 62. The initial methionine starts at 1257 of SEQ ID NO: 62 and the putative stop codon begins at position 2187.

10

## EXAMPLE 4

A. Expression of SEQ ID NO: 4, 13, 20, 29, 36, 42, 61, or 63 in cells

Chinese Hamster Ovary (CHO) cells or other suitable cell types are grown in DMEM (ATCC) and 10% fetal bovine serum (FBS) (Gibco) to 70% confluence. Prior to transfection the media is changed to DMEM and 0.5% FCS. Cells are transfected with cDNAs for SEQ ID NO: 4, 13, 20, 29, 36, 42, 61, or 63 or with pBGal vector by the FuGENE-6 transfection reagent (Boehringer). In summary, 4  $\mu$ l of FuGENE-6 is diluted in 100  $\mu$ l of DMEM and incubated for 5 minutes. Then, this is added to 1  $\mu$ g of DNA and incubated for 15 minutes before adding it to a 35 mm dish of CHO cells. The CHO cells are incubated at 37°C with 5% CO<sub>2</sub>. After 24 hours, media and cell lysates are collected, centrifuged and dialyzed against assay buffer (15 mM Tris pH 7.6, 134 mM NaCl, 5 mM glucose, 3 mM CaCl<sub>2</sub> and MgCl<sub>2</sub>).

20

B. Expression Study Using SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62

The expression of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 in various tissues is analyzed using a semi-quantitative polymerase chain reaction-based technique. Human cDNA libraries are used as sources of expressed genes from tissues of interest (adult bladder, adult brain, adult heart, adult kidney, adult lymph node, adult liver, adult lung, adult ovary, adult placenta, adult rectum, adult spleen, adult testis, bone marrow, thymus, thyroid gland, fetal kidney, fetal liver, fetal liver-spleen, fetal skin, fetal brain, fetal leukocyte and macrophage). Gene-specific primers are used to amplify portions of SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 sequence from the samples. Amplified products are separated on an agarose gel, transferred and chemically linked to a nylon filter. The filter is then hybridized with a radioactively labeled (<sup>32</sup>P-dCTP) double-stranded probe generated from SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21,

123

26-28, 30, 33-35, 37, 41, 43, 60, or 62 using a Klenow polymerase, random-prime method. The filters are washed (high stringency) and used to expose a phosphorimaging screen for several hours. Bands indicate the presence of cDNA including SEQ ID NO: 1-3, 5, 10-12, 14, 17-19, 21, 26-28, 30, 33-35, 37, 41, 43, 60, or 62 sequences in a specific library, and thus mRNA expression in the corresponding cell type or tissue.

5

## CLAIMS

## WE CLAIM:

5 1. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of SEQ ID NO: 2-3, 5, 11-12, 14, 18-19, 21, 27-28, 30, 34-35, 37, 41, 43, 60, or 62, the translated protein coding portion thereof, the mature protein coding portion thereof, the extracellular portion thereof, or the active domain thereof.

10 2. An isolated polynucleotide encoding a polypeptide with biological activity, which polynucleotide hybridizes to the complement of a polynucleotide of claim 1 under stringent hybridization conditions.

15 3. An isolated polynucleotide encoding a polypeptide with biological activity, said polynucleotide having greater than about 90 % sequence identity with the polynucleotide of claim 1.

4. The polynucleotide of claim 1 which is a DNA sequence.

20 5. An isolated polynucleotide which comprises the complement of the polynucleotide of claim 1.

6. A vector comprising the polynucleotide of claim 1.

25 7. An expression vector comprising the polynucleotide of claim 1.

8. A host cell genetically engineered to express the polynucleotide of claim 1.

30 9. The host cell of claim 8 wherein the polynucleotide is in operative association with a regulatory sequence that controls expression of the polynucleotide in the host cell.

10. An isolated polypeptide comprising an amino acid sequence which is at least 80% identical to the amino acid sequence selected from the group consisting of SEQ ID NO: 4, 6-9, 13, 15-16, 20, 22-25, 29, 31-32, 36, 38-40, 42, 44-47, 61, or 63, the translated protein coding portion thereof, the mature protein coding portion thereof, the extracellular portion thereof, or the active domain thereof.

11. A composition comprising the polypeptide of claim 10 and a carrier.

12. A polypeptide, having GPCR-like activity, comprising at least ten consecutive amino acids from the polypeptide sequences selected from the group consisting of SEQ ID NO: 4, 6-9, 13, 15-16, 20, 22-25, 29, 31-32, 36, 38-40, 42, 44-47, 61, or 63.

13. The polypeptide of claim 12, comprising at least five consecutive amino acids from the polypeptide sequences selected from the group consisting of SEQ ID NO: 4, 6-9, 13, 15-16, 20, 22-25, 29, 31-32, 36, 38-40, 42, 44-47, 61, or 63.

14. A polynucleotide encoding a polypeptide according to claim 12.

15. A polynucleotide encoding a polypeptide according to claim 13.

16. A polynucleotide encoding a polypeptide according to claim 10.

17. An antibody specific for the polypeptide of claim 10.

18. A method for detecting the polynucleotide of claim 1 in a sample, comprising:

a) contacting the sample with a compound that binds to and forms a complex with the polynucleotide of claim 1 for a period sufficient to form the complex; and

b) detecting the complex, so that if a complex is detected, the polynucleotide of claim 1 is detected.

19. A method for detecting the polynucleotide of claim 1 in a sample, comprising:

- a) contacting the sample under stringent hybridization conditions with nucleic acid primers that anneal to the polynucleotide of claim 1 under such conditions;
- b) amplifying a product comprising at least a portion of the polynucleotide of claim 1; and
- c) detecting said product and thereby the polynucleotide of claim 1 in the sample.

5

20. The method of claim 19, wherein the polynucleotide comprises an RNA molecule and the method further comprises reverse transcribing an annealed RNA molecule into a cDNA polynucleotide.

10

21. A method for detecting the polypeptide of claim 10 in a sample, comprising:

- a) contacting the sample with a compound that binds to and forms a complex with the polypeptide under conditions and for a period sufficient to form the complex; and
- b) detecting formation of the complex, so that if a complex formation is detected, the polypeptide of claim 10 is detected.

20

22. A method for identifying a compound that binds to the polypeptide of claim 10, comprising:

- a) contacting the compound with the polypeptide of claim 10 under conditions and for a time sufficient to form a polypeptide/compound complex; and
- b) detecting the complex, so that if the polypeptide/compound complex is detected, a compound that binds to the polypeptide of claim 10 is identified.

25

23. A method for identifying a compound that binds to the polypeptide of claim 10, comprising:

- a) contacting the compound with the polypeptide of claim 10, in a cell, for a time sufficient to form a polypeptide/compound complex, wherein the complex drives expression of a reporter gene sequence in the cell; and
- b) detecting the complex by detecting reporter gene sequence expression, so that if the polypeptide/compound complex is detected, a compound that binds to the polypeptide of claim 10 is identified.

127

24. A method of producing a GPCR-like polypeptide, comprising, culturing the host cell of claim 8 under conditions sufficient to express the polypeptide in said cell; and

- b) isolating the polypeptide from the cell culture or cells of step (a).

5

25. A kit comprising the polypeptide of claim 10.

26. A nucleic acid array comprising the polynucleotide of claim 1 or a unique segment of the polynucleotide of claim 1 attached to a surface.

10

27. The array of claim 26, wherein the array detects full-matches to the polynucleotide or a unique segment of the polynucleotide of claim 1.

28. The array of claim 26, wherein the array detects mismatches to the polynucleotide or a unique segment of the polynucleotide of claim 1.

15

29. A method of treatment of a subject in need of enhanced activity or expression of GPCR-like polypeptide of claim 10 comprising administering to the subject a composition selected from the group consisting of:

- (a) a therapeutic amount of an agonist of said polypeptide;
- (b) a therapeutic amount of the polypeptide; and
- (c) a therapeutic amount of a polynucleotide encoding the polypeptide in a form and under conditions such that the polypeptide is produced, and a pharmaceutically acceptable carrier.

25

30. A method of treatment of a subject having need to inhibit activity or expression of GPCR-like polypeptide of claim 10 comprising administering to the subject a composition selected from the group consisting of:

- (a) a therapeutic amount of an antagonist to said polypeptide;
- (b) a therapeutic amount of a polynucleotide that inhibits the expression of the nucleotide sequence encoding said polypeptide; and
- (c) a therapeutic amount of a polypeptide that competes with the GPCR-like polypeptide for its ligand and a pharmaceutically acceptable carrier.

30

128

BLASTP ALIGNMENT OF SEQ ID NO: 4, G PROTEIN-COUPLED RECEPTOR-LIKE POLYPEPTIDE (IDENTIFIED AS GPCR-LIKE) WITH HUMAN CGI-40 PROTEIN SEQ ID NO: 48

Query: G protein-coupled receptor-like polypeptide (SEQ ID NO: 4)  
 Subject: g14929551 (AF151799) CGI-40 protein [Homo sapiens] (SEQ ID NO: 48)  
 Score = 1709 (606.7 bits), Expect = 9.4e-176, p = 9.4e-176  
 Identifiers = 136/526 (63%), Positives = 400/526 (76%)  
 Length = 845  
 Query: 283 QTNLQKRLKLVLPKIKKSVVPSLPVLPGLVGVHVLRFQKSID 342  
 Subject: 318 ENMR-QKKLTLVADLRACPSGCHPRV-LADSPGSSPREGYNGSGFKNVSGSTGVD- 374  
 Query: 343 SFGSDSQNMVASHPIAASPTSGSNVGTIDSSSSPFGKQSSSDGQPDTSVSE 402  
 Subject: 375 SAGTGDLSVYGQGHQDFKRLPSSQ---NRQICIAMGRSPFVGTNP--RVDSSVSE 428  
 Query: 403 SPFTMDISDKNIIRTKRFLYLSDLRSKDRRIYKSKKIYFNNITIAVAVLPVQL 462  
 Subject: 429 DDVPLTIDSDSKNVIIRTKQYLVADLRKDKRVLRKQYIYFNNITIAVAVLPVQL 488  
 Query: 463 VITQVAVVNSGNDICYYNPLCAHPLGLVLSAFTNLSLGLLGLLILVLPDILH 522  
 Subject: 489 VITQVAVVNSGNDICYYNPLCAHPLGLVLSAFTNLSLGLLGLLILVLPDILH 548  
 Query: 523 RALAKADIPAVRGVLPKHEGTLVYMGATLMDRGGLSACVHVCBNYSNGQFTSPMYIA 582  
 Subject: 549 RNLALRNDLCALGCGIPKHEGTLVYMGATLMDRGGLSACVHVCBNYSNGQFTSPMYIA 608  
 Query: 583 GLCLMLKYQTRHFDINASAVASFAVAVTINVTALGVFGKNDVMWVIFSAIHVLSLA 642  
 Subject: 609 GLCLMLKYQTRHFDINASAVASFAVAVTINVTALGVFGKNDVMWVIFSAIHVLSLA 668  
 Query: 643 LSTQVYMGARFTRIDLGTRHRAAVVFTDTCIQGSRPLVMDNVLTVVGNLVMSGPALPGL 702  
 Subject: 669 LSTQVYMGARFTRIDLGTRHRAAVVFTDTCIQGSRPLVMDNVLTVVGNLVMSGPALPGL 728  
 Query: 703 IYRPDPASVYMGATLGLVLCNLTLLVLAAYIIMKLSSEKVLFPVPLFCIVATVAMNAALYFP 762  
 Subject: 729 IYRPDPASVYLAIGIICNLTLLVLAAYIIMKLSSEKVLFPVPLFCIVTCTSVVWGPALPFP 788  
 Query: 763 QNLSWSWGPAPASRRKRNRCIILLDPDDHDIMHFLSATATLPSPFLV 808  
 Subject: 789 QCLSTWQKTPASRRKRNRCIILLDPDDHDIMHFLSATATLPSPFLV 834

FIG. 1

BLASTP ALIGNMENT OF SEQ ID NO: 4, G PROTEIN-COUPLED RECEPTOR-LIKE POLYPEPTIDE (IDENTIFIED AS GPCR-LIKE) WITH PROTEIN OF CLONE CT748\_2 SEQ ID NO: 49

Query: G protein-coupled receptor-like polypeptide (SEQ ID NO: 4)  
 Subject: M57901 Protein of clone CT748\_2. [Homo sapiens] (SEQ ID NO: 49)  
 Length = 479  
 Score = 2220 (786.5 bits), Expect = 9.7e-231, p = 9.7e-231  
 Identifiers = 431/445 (96%), Positives = 433/445 (97%)  
 Query: 388 GSPGQSDSDSSVSPDTPMDIESDKNIIRTKRFLYLSDLRSKDRRIYKSKKIYFNN 447  
 Subject: 36 GHRASQYQT-APVEESDPTMDIESDKNIIRTKRFLYLSDLRSKDRRIYKSKKIYFNN 94  
 Query: 448 IITIAVAVVNSGNDICYYNPLCAHPLGLVLSAFTNLSLGLLILVLPDILH 507  
 Subject: 95 IITIAVAVVNSGNDICYYNPLCAHPLGLVLSAFTNLSLGLLILVLPDILH 154  
 Query: 508 GFLPLIIVLRDILHRNALBAKDIPAVRGVLPKHEGTLVYMGATLMDRGGLSACVHVCBN 567  
 Subject: 155 GFLPLIIVLRDILHRNALBAKDIPAVRGVLPKHEGTLVYMGATLMDRGGLSACVHVCBN 214  
 Query: 568 YSNFQDPTSPMYMIAAGLCMLKLYQTRHFDINASAVASFAVAVTINVTALGVFGKNDVM 627  
 Subject: 215 YSNFQDPTSPMYMIAAGLCMLKLYQTRHFDINASAVASFAVAVTINVTALGVFGKNDVM 274  
 Query: 628 FVIFSAIHVLSLAFLSTQIYYMGARFTRIDLGTRHRAAVVFTDTCIQGSRPLVMD 682  
 Subject: 275 FVIFSAIHVLSLAFLSTQIYYMGARFTRIDLGTRHRAAVVFTDTCIQGSRPLVMD 334  
 Query: 683 RMVLLVGNLVMSGPALPGLIYRPDPASVYMGATLGLVLCNLTLLVLAAYIIMKLSSEKVLFP 742  
 Subject: 335 RMVLLVGNLVMSGPALPGLIYRPDPASVYMGATLGLVLCNLTLLVLAAYIIMKLSSEKVLFP 394  
 Query: 743 PLFCIVATVAMNAALYFPPLFCIVATVAMNAALYFPPLFCIVATVAMNAALYFP 802  
 Subject: 395 PLFCIVATVAMNAALYFPPLFCIVATVAMNAALYFPPLFCIVATVAMNAALYFP 454  
 Query: 803 PPSFLVLTLLDDDLVVRDQIPVF 827  
 Subject: 455 PPSFLVLTLLDDDLVVRDQIPVF 479

FIG. 2

BLASTP ALIGNMENT OF SEQ ID NO: 13, G PROTEIN-COUPLED RECEPTOR-LIKE POLYPEPTIDE (IDENTIFIED AS GPCR-LIKE) WITH SIX TRANSMEMBRANE EPIITHELIAL ANTIGEN OF PROSTATE SEQ ID NO: 50

Query: G protein-coupled receptor-like polypeptide (SEQ ID NO: 13)  
Sbjct: g16572948 (A186249) six transmembrane epithelial antigen of prostate [Homo sapiens] (SEQ ID NO: 50)  
Length = 339

Score = 724 (259.9 bits), Expect = 2.3e-71, P = 2.3e-71  
Identities = 126/267 (47%), Positives = 184/267 (68%)

Query: 208 LTPMKVPTLALGLFVAVNFRVDTLPYVQBSQNKFFKLPSVSVNTTLPCVAVYLL 267  
Sbjct: 67 LFPQWMLPIKIKIAIIVASLVTLYTLTRRVIHPLATSHQGYFYKIPILVINKVLPMVISTLL 126  
Query: 268 SLVYLPFGVLAALQLRRGRTKYQRPDMLDHRQIGILSPFCALHALVSPCLPLRR 327  
Sbjct: 127 ALVYLPFGVLAIVQLHNGRTKYKKPPHMLTRKQPGILSPFAVLAHAYSLSYPMRR 186  
Query: 328 AHRVDLVNLAVKQVLANKSKSHLMVBEVWRRRIYILSLGLALAGTSLSLAVTSLSPIANSIN 387  
Sbjct: 187 SYRYKLLNMAVYQVQVQNKEDAMWIRHDDVMRMRIYVSLGIVGLAIALAVTSIPSDSLT 246  
Query: 388 WRPSFVQSSISGLFVALVLTSLHTLTLYGWTAFAPESRKYKPYLLPPTFTLTLPVCCVILAKA 447  
Sbjct: 247 WRPSHYIQSKIGIVSLTLTGTHLILFAMNKMWDIKQFVMWYTPPTFMIAVFLPIVILIFRS 306  
Query: 448 LFLPCISRLAIRRMGRRESSTIKFT 474  
Sbjct: 307 ILFLPCILRKLKIRHGMEDVTKINKT 333

FIG. 3

BLASTP ALIGNMENT OF SEQ ID NO: 13, G PROTEIN-COUPLED RECEPTOR-LIKE POLYPEPTIDE (IDENTIFIED AS GPCR-LIKE) WITH HUMAN STRAP-1 PROTEIN SEQ ID NO: 51

Query: G protein-coupled receptor-like polypeptide (SEQ ID NO: 13)  
Sbjct: Y58194 Human STRAP-1 protein. [Homo sapiens] (SEQ ID NO: 51)  
Length = 339

Score = 724 (259.9 bits), Expect = 3.3e-72, P = 3.3e-72  
Identities = 126/267 (47%), Positives = 184/267 (68%)

Query: 208 LTPMKVPTLALGLFVAVNFRVDTLPYVQBSQNKFFKLPSVSVNTTLPCVAVYLL 267  
Sbjct: 67 LFPQWMLPIKIKIAIIVASLVTLYTLTRRVIHPLATSHQGYFYKIPILVINKVLPMVISTLL 126  
Query: 268 SLVYLPFGVLAALQLRRGRTKYQRPDMLDHRQIGILSPFCALHALVSPCLPLRR 327  
Sbjct: 127 ALVYLPFGVLAIVQLHNGRTKYKKPPHMLTRKQPGILSPFAVLAHAYSLSYPMRR 186  
Query: 328 AHRVDLVNLAVKQVLANKSKSHLMVBEVWRRRIYILSLGLALAGTSLSLAVTSLSPIANSIN 387  
Sbjct: 187 SYRYKLLNMAVYQVQVQNKEDAMWIRHDDVMRMRIYVSLGIVGLAIALAVTSIPSDSLT 246  
Query: 388 WRPSFVQSSISGLFVALVLTSLHTLTLYGWTAFAPESRKYKPYLLPPTFTLTLPVCCVILAKA 447  
Sbjct: 247 WRPSHYIQSKIGIVSLTLTGTHLILFAMNKMWDIKQFVMWYTPPTFMIAVFLPIVILIFRS 306  
Query: 448 LFLPCISRLAIRRMGRRESSTIKFT 474  
Sbjct: 307 ILFLPCILRKLKIRHGMEDVTKINKT 333

FIG. 4

BLASTS ALIGNMENT OF SEQ ID NO: 20, G PROTEIN-COUPLED RECEPTOR-LIKE  
POLYPEPTIDE (IDENTIFIED AS GPCR-LIKE) WITH SEVEN TRANSMEMBRANE RECEPTOR  
PROTEIN SEQ ID NO: 52

Query: G protein-coupled receptor-like polypeptide (Seq ID NO: 20)  
Subject: g15525078 (AB019120) seven transmembrane receptor [Rattus norvegicus] (Seq ID NO: 52)  
Length = 1349  
Score = 5072 (1790.5 bits), Expect = 0.0, p = 0.0  
Identifiers = 984/1354 (72%), Positives = 1108/1354 (81%)

FIG. 5A

Query:	1	MSRPTLTLCHAEIYSSAALNNWYSEI THPSLHSGAGALRKALRKAVAPKA
Query:	2	1 KMS R TL + AII SSAA + E + HPL L EER AGER LR KRYVA P
Query:	3	1 MESSRWTLVFLVLIYLISSSEALNSRPAEIVHPLI OEHLAGELRFRKAAVGPVA
Query:	4	61 ERYVNNIISFNASLPLDPIKALISLPPHANNPTDITLINTVCSPPAHMNC
Query:	5	61 ERYLVDSIFSNVSLPLRESIRAHNTSLRFPQGNCD ---LTSMATVTCPTGNDITC
Query:	6	121 SCDFYWRKNTLTCGDFRPLCPGHCSCCLPSPNGPCLDQ -VLMNRKALN
Query:	7	117 PCBKQYGPBEERCLSTLCGDEHNSALPGKYNCLKGLPQGPQCLPBTYILIKIKVRLN
Query:	8	180 VQDFDMLNNTSLRSXKTKDILYLFARQVGL LKPGKVLTCGSKSVAAVETKTP
Query:	9	177 IGFQDLENTSALYRSYKTKDILYBRAPAGVRLPGFRSVLTGKGSVAVIVASAA
Query:	10	240 SLE LHKANQGVDSITLTKKDNVSLTAKNSMFLVPIEISLQVTCACERLV
Query:	11	237 PLRGSIKHANKQVLIQNTLTKDNVSPQGPFSNRTKLVLEISQVNLTECSRNV
Query:	12	299 SSNSWYIEEODLQNSNSRSLT+LNN+ST+L+NN+ST+L+N DAG L + LDIR
Query:	13	297 SNTSMWYEGKRSIDQNSKDFSIHTSIINNTSLVTLTLTNFNTQHALYGCXNLDIR

BLASTS ALIGNMENT OF SEQ ID NO: 20, G PROTEIN-COUPLED RECEPTOR-LIKE  
POLYPEPTIDE (IDENTIFIED AS GPCR-LIKE) WITH SEVEN TRANSMEMBRANE RECEPTOR  
PROTEIN SEQ ID NO: 52

[illegible]

FIG 5B



BLASTP ALIGNMENT OF SEQ ID NO: 20, G PROTEIN-COUPLED RECEPTOR-LIKE  
POLYPEPTIDE (IDENTIFIED AS GPCR-LIKE) WITH SEVEN TRANSMEMBRANE RECEPTOR  
PROTEIN SEQ ID NO: 52

Query: 896 SIATVAPPTLALADODRNFASTLVTITVSHNTMTPTKNNSPSGCTKCV 955  
SIATVAPPTLALAD 0 + SIATVITVSHN PRISMTPKNN SCG + CV 955  
Query: 956 FMRNLTANTGWDSSCGVYZGDDN---VFCIDHLTSPSILMSPPSPSSLLGILT 1012  
FMRNLTANTGWDSSCGVYZGDDNDRVPCCKNHLTSPSILMSPPSPSSLLGILT 1015  
Query: 1013 DIISYGVFSTLSLAACLVAVAVWMSKSVTRNRTSYMRHCIVNIAALVATMTVVA 1072  
DIISYGVFSTLSLAACLVAVAVWMSKSVTRNRTSYMRHCIVNIAALVATMTVVA 1075  
Query: 1073 AIDNNKATLCTKACVATLTHIRFVLSVFFMMLTGILMLPYRRL+PILH+STQKALAF 1132  
AIDNNKATLCTKACVATLTHIRFVLSVFFMMLTGILMLPYRRL+PILH+STQKALAF 1135  
Query: 1133 CLGCGCLASIVITLCAATQPRVETRTKNCMLNWDRTALALVAVNITITV 1192  
CLGCGCLASIVITLCAATQPRVETRTKNCMLNWDRTALALVAVNITITV 1195  
Query: 1193 VITKILRPISGDKPKQKQKSSLPQISKSISGLVLPBLGLTWGFGTLVFNHIIIA 1252  
VITKILRPISGDKPKQKQKSSLPQISKSISGLVLPBLGLTWGFGTLVFNHIIIA 1255  
Query: 1253 ILTVFGCLTILPGLMDLKQVALLNFKSLRMSWSSQHSKSTSLSSSTPVSMSPIR 1312  
ILTVFGCLTILPGLMDLKQVALLNFKSLRMSWSSQHSKSTSLSSSTPVSMSPIR 1315  
Query: 1313 FNNLPKGTCTVNTVTEATLSSSLNNSASSSLIN 1346  
FNNLPKGTCTVNTVTEATLSSSLNNSASSSLIN 1349  
Query: 1316 FNNLPKGTCTVNTVTEATLSSSLNNSASSSLIN 1349  
FNNLPKGTCTVNTVTEATLSSSLNNSASSSLIN 1349

FIG. 5C

BLASTP ALIGNMENT OF SEQ ID NO: 20, G PROTEIN-COUPLED RECEPTOR-LIKE  
POLYPEPTIDE (IDENTIFIED AS GPCR-LIKE) WITH HUMAN BRAIN-DERIVED G PROTEIN-  
COUPLED RECEPTOR PROTEIN SEQ ID NO: 53

Query: G protein-coupled receptor-like polypeptide (Seq ID NO: 20)  
Subject: Y40440 Human brain-derived G-protein coupled receptor protein. [Homo sapiens] (Seq ID NO: 53)  
Length = 986  
Score = 5161 (1821.8 bits), Expect = 0.0, P = 0.0  
Identities = 986/986 (100%), Positives = 986/986 (100%)

Query: 420 CCKKIDVAPRIQILANBEMKVMCDNNPVSIMCCSQGNVMSKVMKQKGININIGTPTDI 60  
CCKKIDVAPRIQILANBEMKVMCDNNPVSIMCCSQGNVMSKVMKQKGININIGTPTDI 60  
Query: 421 DSSCSRYTLKADGTQCPSSGSGTIVITTCCEPISAVAGAFGASNAIKVITISVANLITTPDI 120  
DSSCSRYTLKADGTQCPSSGSGTIVITTCCEPISAVAGAFGASNAIKVITISVANLITTPDI 120  
Query: 481 SVSBQGNFSLKICISDVSNYDENVWNTSAGIKIYQRFYTRRYLDGABSVLTVITSTRMN 540  
SVSBQGNFSLKICISDVSNYDENVWNTSAGIKIYQRFYTRRYLDGABSVLTVITSTRMN 540  
Query: 541 GTCHCIFRYKNSYSIATKQVIVHPLPLKLNIMVDPLBATVSCSGSHIKKCCIEEDGYKV 600  
GTCHCIFRYKNSYSIATKQVIVHPLPLKLNIMVDPLBATVSCSGSHIKKCCIEEDGYKV 600  
Query: 601 TFHMSSSLPAAKEVKKQVCCYKHNFMNASVSMCSKTVDVCCCHPTNANNSVMSPTLN 660  
TFHMSSSLPAAKEVKKQVCCYKHNFMNASVSMCSKTVDVCCCHPTNANNSVMSPTLN 660  
Query: 720 LVFGENITCGDPVIGVGEFPGKVIQKLCRFNSVNPSSPSPPIGCTITTYKCVGSGEKKND 720  
LVFGENITCGDPVIGVGEFPGKVIQKLCRFNSVNPSSPSPPIGCTITTYKCVGSGEKKND 720  
Query: 780 ISAPINSLTGQNAKALIKSPSQDEMLPTTKDLKSLISIDKAEHRISSSPGLGALINILDL 780  
ISAPINSLTGQNAKALIKSPSQDEMLPTTKDLKSLISIDKAEHRISSSPGLGALINILDL 780  
Query: 840 STVPQVNSBEMTHVLSTVNLILGKPVLTNKKVILQGMNQSGLLHSEVRPSQALQSGD 840  
STVPQVNSBEMTHVLSTVNLILGKPVLTNKKVILQGMNQSGLLHSEVRPSQALQSGD 840  
Query: 841 STVPQVNSBEMTHVLSTVNLILGKPVLTNKKVILQGMNQSGLLHSEVRPSQALQSGD 840  
STVPQVNSBEMTHVLSTVNLILGKPVLTNKKVILQGMNQSGLLHSEVRPSQALQSGD 840

FIG. 6A



BLASTP ALIGNMENT OF SEQ ID NO: 29 G PROTEIN-COUPLED RECEPTOR-LIKE POLYPEPTIDE (IDENTIFIED AS GPCR-LIKE) WITH HUMAN h-TRAAK POLYPEPTIDE #1 SEQ ID NO: 55

Query: G protein-coupled receptor-like polypeptide (SEQ ID NO: 29)  
Sbjct: Y94425 Human h-TRAAK polypeptide #1 (Homo sapiens) (SEQ ID NO: 55)  
Score = 2062 (730.9 bits), Expect = 5.4e-214, P = 5.4e-214  
Length = 393  
Identities = 392/392 (100%), Positives = 392/392 (100%)

Query: 540 MRSITLALATLALVLTALVSGALVFRALDQPHBQQAQREKIGKVRERFLFAHPCVSDQELG  
Sbjct: 1 MRSITLALATLALVLTALVSGALVFRALDQPHBQQAQREKIGKVRERFLFAHPCVSDQELG  
Query: 600 LKIEVADLTCGADPPTNSTNSHSHAMDLGSAFFSGTIIITIGVGNVALRTDAGRLFC  
Sbjct: 61 LKIEVADLTCGADPPTNSTNSHSHAMDLGSAFFSGTIIITIGVGNVALRTDAGRLFC  
Query: 660 IYALVGIPIFGIILACGVDRLGSSILRHGIGHIEALFLKMHVPPFLVRLSAMTLILIG  
Sbjct: 121 IYALVGIPIFGIILACGVDRLGSSILRHGIGHIEALFLKMHVPPFLVRLSAMTLILIG  
Query: 720 LFLVLTPTFCYMDWMSKLEAIVFVIVTLITVGGPDYVAGADPRQDSPAYQPLVMWIL  
Sbjct: 181 LFLVLTPTFCYMDWMSKLEAIVFVIVTLITVGGPDYVAGADPRQDSPAYQPLVMWIL  
Query: 779 LFLVLTPTFCYMDWMSKLEAIVFVIVTLITVGGPDYVAGADPRQDSPAYQPLVMWIL  
Sbjct: 240 LFLVLTPTFCYMDWMSKLEAIVFVIVTLITVGGPDYVAGADPRQDSPAYQPLVMWIL  
Query: 839 LGLAVFASVLTITIGMILRVSRTRABKGGTLTAQASWMTGIVTAVTQRAAGAAPPEKE  
Sbjct: 780 LGLAVFASVLTITIGMILRVSRTRABKGGTLTAQASWMTGIVTAVTQRAAGAAPPEKE  
Query: 899 QELTPPPCAQDLGRSPSPPEKAQPPSPPTASALDYPSENLAFIDESSDTQSERGCP  
Sbjct: 301 QELTPPPCAQDLGRSPSPPEKAQPPSPPTASALDYPSENLAFIDESSDTQSERGCP  
Query: 900 LPRAPRGRRRPNPRKPVPRRGGRRNDKGV  
Sbjct: 361 LPRAPRGRRRPNPRKPVPRRGGRRNDKGV

FIG. 8

BLASTP ALIGNMENT OF SEQ ID NO: 36, G PROTEIN-COUPLED RECEPTOR-LIKE POLYPEPTIDE (IDENTIFIED AS GPCR-LIKE) WITH HUMAN OLFACTORY RECEPTOR POLYPEPTIDE SEQ ID NO: 56

Query: G protein-coupled receptor-like polypeptide (SEQ ID NO: 36)  
Sbjct: g12921716 (U86281) olfactory receptor [Homo sapiens] (SEQ ID NO: 56)  
Score = 779 (279.3 bits), Expect = 3.3e-77, P = 3.3e-77  
Identities = 146/166 (87%), Positives = 154/166 (92%)  
Length = 217

Query: 1 MSYDRYVAICHFLHYPIITMTWKVCITLALITSWTCGSLTAMVHVSILRLTPCCGPRHINHP  
Sbjct: 60 MSYDRYVAICHFLHYPIITMTWKVCITLALITSWTCGSLTAMVHVSILRLTPCCGPRHINHP  
Query: 52 MSYDRYVAICHFLHYPIITMTWKVCITLALITSWTCGSLTAMVHVSILRLTPCCGPRHINHP  
Sbjct: 111 MSYDRYVAICHFLHYPIITMTWKVCITLALITSWTCGSLTAMVHVSILRLTPCCGPRHINHP  
Query: 61 FCEILSVLRLACADVTALNOVITFAACMFILVGPICLTLVLSYSHTIILALRIQSGGRKKA  
Sbjct: 120 FCEILSVLRLACADVTALNOVITFAACMFILVGPICLTLVLSYSHTIILALRIQSGGRKKA  
Query: 112 FCEILSVLRLACADVTALNOVITFAACMFILVGPICLTLVLSYSHTIILALRIQSGGRKKA  
Sbjct: 171 FCEILSVLRLACADVTALNOVITFAACMFILVGPICLTLVLSYSHTIILALRIQSGGRKKA  
Query: 121 FSTCSSHLCLVAVGLPFGSAIVMYMAFKSRHPEEQKVLFLFYSSNP  
Sbjct: 166 FSTCSSHLCLVAVGLPFGSAIVMYMAFKSRHPEEQKVLFLFYSSNP  
Query: 172 FSTCSSHLCLVAVGLPFGSAIVMYMAFKSRHPEEQKVLFLFYSSNP  
Sbjct: 217 FSTCSSHLCLVAVGLPFGSAIVMYMAFKSRHPEEQKVLFLFYSSNP

FIG. 9

BLASTP ALIGNMENT OF SEQ ID NO: 36 G PROTEIN-COUPLED RECEPTOR-LIKE POLYPEPTIDE (IDENTIFIED AS GPCR-LIKE) WITH HUMAN G PROTEIN-COUPLED RECEPTOR GPR1

POLYPEPTIDE SEQ ID NO: 57

Query: G protein-coupled receptor-like polypeptide (SEQ ID NO: 36)  
 Subject: M04244 Human G-protein coupled receptor GPR1 (Homo sapiens) (SEQ ID NO: 57)  
 Score = 835 (299.0 bits), Expect = 5.7e-84, E = 5.7e-84  
 Identities = 159/171 (92%), Positives = 160/171 (93%)  
 Length = 296

Query: 1 MSYDRVAICHPLKRYEIIIMTKVCITLITMTWTCGSLTAAVHVSILRLPFCGPRRIINHP 60  
 MSYDRVAICHPLKRYEIIIMTKVCITLITMTWTCGSLTAAVHVSILRLPFCGPRRIINHP  
 Subject: 121 MSYDRVAICHPLKRYEIIIMTKVCITLITMTWTCGSLTAAVHVSILRLPFCGPRRIINHP 180  
 Query: 61 FCEILSVLRACADPTWLNQVIFPAACKEPILVGPICLVLVSYSHTLGLGIIQSGGRKA 120  
 FCEILSVLRACADPTWLNQVIFPAACKEPILVGPICLVLVSYSHTLGLGIIQSGGRKA  
 Subject: 181 FCEILSVLRACADPTWLNQVIFPAACKEPILVGPICLVLVSYSHTLGLGIIQSGGRKA 240  
 Query: 121 FSTCSSHLCVVGGLFPGSAIVMYAPKSRHREBQKVLFLFYSSRN-PMANP 170  
 FSTCSSHLCVVGGLFPGSAIVMYAPKSRHREBQKVLFL + PML P  
 Subject: 241 FSTCSSHLCVVGGLFPGSAIVMYAPKSRHREBQKVLFLFLIPLSTPMKP 291

FIG. 10

BLASTP ALIGNMENT OF SEQ ID NO: 42, G PROTEIN-COUPLED RECEPTOR-LIKE POLYPEPTIDE (IDENTIFIED AS GPCR-LIKE) WITH SIMILAR TO MOUSE OLFACTORY RECEPTOR 13 POLYPEPTIDE SEQ ID NO: 58

Query: G protein-coupled receptor-like polypeptide (SEQ ID NO: 42)  
 Subject: g14159884 (AC005587) similar to mouse olfactory receptor 13; similar to p34984 (PID:g464305) (Homo sapiens) (SEQ ID NO: 58)  
 Score = 1067 (380.7 bits), Expect = 1.0e-107, E = 1.0e-107  
 Identities = 209/304 (68%), Positives = 247/304 (81%)  
 Length = 310

Query: 1 MVRNQTWTEFLTLGLGPIRIQMLTFLGSLFYIFTLTGNGALICGLISDRLHTPMVF 60  
 M N T + EFLTLG + GPRIQMLTFLGSLFYIFTLTGNGALICGLISDRLHTPMVF  
 Subject: 1 MODNITSIREFLLGFPVCGPRIQMLTFLGSLFYIFTLTGNGALICGLISDRLHTPMVF 60  
 Query: 61 PLSHLAAVDIAYMTVP+ML NLTLPAPISFAG N QTFP + E ECTLLTV+MSYD 120  
 PLSHLAAVDIAYMTVP+ML NLTLPAPISFAG N QTFP + E ECTLLTV+MSYD  
 Subject: 61 PLSHLAAVDIAYMTVP+ML NLTLPAPISFAG N QTFP + E ECTLLTV+MSYD 120  
 Query: 121 RYVAICHPLKRYEIIIMTKVCITLITMTWTCGSLTAAVHVSILRLPFCGPRRIINHP 178  
 RYVAICHPLKRYEIIIMTKVCITLITMTWTCGSLTAAVHVSILRLPFCGPRRIINHP  
 Subject: 121 RYVAICHPLKRYEIIIMTKVCITLITMTWTCGSLTAAVHVSILRLPFCGPRRIINHP 180  
 Query: 179 LSVLRACADPTWLNQVIFPAACKEPILVGPICLVLVSYSHTLGLGIIQSGGRKA 238  
 LSVLRACADPTWLNQVIFPAACKEPILVGPICLVLVSYSHTLGLGIIQSGGRKA  
 Subject: 181 LSVLRACADPTWLNQVIFPAACKEPILVGPICLVLVSYSHTLGLGIIQSGGRKA 240  
 Query: 239 SSHLCVVGGLFPGSAIVMYAPKSRHREBQKVLFLFYSSRN-PMANP 298  
 SSHLCVVGGLFPGSAIVMYAPKSRHREBQKVLFLFYSSRN-PMANP  
 Subject: 241 SSHLCVVGGLFPGSAIVMYAPKSRHREBQKVLFLFYSSRN-PMANP 300  
 Query: 299 RYAL 302  
 Subject: 301 KXVL 304

FIG. 11

SEQUENCE LISTING

<110> HYSEQ, Inc.  
Yamazaki, Victoria  
Tang, Y. Tom  
Liu, Chenghua  
Zhou, Ping  
Wang, Dunrui  
Zhang, Jie  
Ren, Feiyan  
Asundi, Vinod  
Drmanac, Radoje T

<120> METHODS AND MATERIALS RELATING TO G PROTEIN-COUPLED RECEPTOR-LIKE (GPCR-LIKE) POLYPEPTIDES AND POLYNUCLEOTIDES

<130> HYS-37CIP

<140> not yet assigned

<141> 2000-12-22

<150> not yet assigned

<151> 2000-12-21

<150> US 09/729,739

<151> 2000-12-04

<150> US 09/653,450

<151> 2000-08-31

<150> US 09/620,312

<151> 2000-07-19

<150> US 09/598,042

<151> 2000-06-20

<150> US 09/552,317

<151> 2000-04-25

<150> US 09/488,725

<151> 2000-01-21

<160> 63

<170> Patent in version 3.0

<210> 1

<211> 412

<212> DNA

<213> Homo sapiens

<400> 1

ctttttacac atcatgaagc tccgcagctc tgaagaagtc ctccagtc cgcctctctg 60  
catcgtggcc accgctgtga tctgggctgc cgcctatat tttttcttc agaatctcag 120  
cagctgggag ggaactccg ccgaatccc ggagaagaac cgcgagtgca tctcgtgga 180  
ttctctgat gaccatgaca tctggcaact cctctctgct actcctctgt tttctctatt 240  
cttgatttg ttaacttgg atgatgacct tgatgtggtt cggagagacc agatccctgt 300  
ctctgaacc tccaacnta agagagggga gggagcgatc aatcttggtg ctgtttcaca 360

BLASTP ALIGNMENT OF SEQ ID NO: 42, G PROTEIN-COUPLED RECEPTOR-LIKE POLYPEPTIDE (IDENTIFIED AS GPCR-LIKE) WITH HUMAN G PROTEIN-COUPLED RECEPTOR GPR1 POLYPEPTIDE SEQ ID NO: 59

Query: G protein-coupled receptor-like polypeptide (SEQ ID NO: 42)  
Sbjct: M04244 Human G-protein coupled receptor GPR1. [Homo sapiens] (SEQ ID NO: 59)  
Length = 296  
Identical = 1128 (472.5 bits), Expect = 3.2e-136, P = 3.2e-136  
Positives = 259/287 (90%), Positives = 264/287 (91%)

Query: 1 MVRNQMTVEFLGLGFLGPRIQMLTFLGFLSPYIFTLTGNGAIIIGLISLDSRLHTPMYF 60  
Sbjct: 5 MVRNQMTVEFLGLGFLGPRIQMLTFLGFLSPYIFTLTGNGAIIIGLISLDSRLHTPMYF 64  
Query: 61 FLSHLAVVDAIVTNTVPQMTLNLHBAKISPAAGCMTQTFGLCLSPGHSCLLLVLMSYD 120  
Sbjct: 65 FLSHLAVVDAIVTNTVPQMTLNLHBAKISPAAGCMTFLLSR H+ECLLLVLMSYD 124  
Query: 121 RYVAICHPLEARYSVIPT--CCITLAIITSMTCGSLANVAVSLILALPFCGPRBINHPFCEI 178  
Sbjct: 125 RYVAICHPLEARYSVIPT--CCITLAIITSMTCGSLANVAVSLILALPFCGPRBINHPFCEI 184  
Query: 179 LSVLTLACADPTMLNQVILFPAACMPLVGLCLVAVSVSHLALAIRIQSGRGRRKAFSTC 238  
Sbjct: 185 LSVLTLACADPTMLNQVILFPAACMPLVGLCLVAVSVSHLALAIRIQSGRGRRKAFSTC 244  
Query: 239 SSHLCVVGTLFPFGSALIVMMAFKSRHHPPEEQKVLFLPLVSSFN-PM-LNP 284  
Sbjct: 245 SSHLCVVGTLFPFGSALIVMMAFKSRHHPPEEQKVLFLPLVSSFN-PM-LNP 291

FIG. 12

aaattaccag tgaccacagc aaagtaacca atgcagatg ctccactcac cc 412

<210> 2

<211> 749

<212> DNA

<213> Homo sapiens

<400> 2

ctttatcac atcatgaagc tccgagctc tgaaaagtc ctccagtc ccgtctctctg 60

catctggcc accgctgtga tctgggtgc cgccctatat tttttctcc agaactctcag 120

cagctggag ggaactccg ccgaatccg ggagaagaac cgcgagtga tctcgtgga 180

tttctcgat gaccatgaca tctggcactt cctctctgct actgctctgt tttttcatt. 240

cttggaattg ttaacttttg atgatgacct tcatgtggtt cggagagacc agatccctgt 300

ctctgaacc tccaacatta agagagggga gggagcgatc aatcttggtg ctgtttcaca 360

aaaattacag tgaccacagc aaagtaacca ctgcagatg ctccactcac cctctgtaga 420

gccactctg catctacaca ggaaggagag gggctgcggg agatttaaac ctgcaagaaa 480

ggaggcagaa ggggagccat gttttgagga cagacgcaaa cctgagagc tgagaacac 540

tgtctcttc catctgcagc tttgggagt gaaacagggat aggcactgca tccaagtcaa 600

ctcaccatct tggggctcct ccacacctca cggagacttg ccagcaatgg cagaatgctg 660

ctgcacactt tcttcaagt gttaccttc ccagaaaggc caagctcgtg gacttcttg 720

ccaaatactt gggtaggccc cgcgttccg 749

<210> 3

<211> 3188

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1) .. (2484)

<400> 3

atg cgc ggc tgc ctg cgg ctc gcg ctg ctc tgc gcg ctg ccc tgg ctc 48

Met Arg Gly Cys Leu Arg Leu Ala Leu Leu Cys Ala Leu Pro Trp Leu 15

1 5 10

ctg ctg cgc ggc tgc ccc ggg cac cgc ggc ggc aaa tcc ccc agg cag ccc 96

Leu Leu Ala Ala Ser Pro Gly His Pro Ala Lys Ser Pro Arg Gln Pro 30

20 25

ccg gca cgc cgc ggc ccc ttc gac gct gcc agg ggc gcc gat ttc 144

Pro Ala Pro Arg Arg Asp Pro Phe Asp Ala Ala Arg Gly Ala Asp Phe 45

35 40

gat cat gtc tac agc ggg gtc gtc aac ctc agc acc agc aac atc tac 192

Asp His Val Tyr Ser Gly Val Val Asn Leu Ser Thr Gln Asn Ile Tyr 60

50 55

tct ttc aac tac acc agc cag ccc gac cag gtc aca gcc gtc agg gtc 240

2

Ser Phe Asn Tyr Thr Ser Gln Pro Asp Gln Val Thr Ala Val Arg Val 80

65 70 75

tat gtc aac agt tcc tct gag aat ctc aac tac cgc gtc ctt gtt gtc 288

Tyr Val Asn Ser Ser Ser Glu Asn Leu Asn Tyr Pro Val Leu Val Val 95

85 90

gtt cgc cag cag aaa gag gtc ctg tcc tgg cag gtt cct ctg ctc ttc 336

Val Arg Gln Gln Lys Glu Val Leu Ser Trp Gln Val Pro Leu Leu Phe 110

100 105

caa gga cta tac cag agg agc tac aat tat caa gaa gtc agc cgc acc 384

Gln Gly Leu Tyr Gln Arg Ser Tyr Asn Tyr Gln Glu Val Ser Arg Thr 125

115 120

tta tgt ccc tca gaa gca acc aat gag acg gga ccc ttg cag caa ctg 432

Leu Cys Pro Ser Glu Ala Thr Asn Glu Thr Gly Pro Leu Gln Gln Leu 140

130 135

ata ttt gta gat gtc gca tcc atg gca ccc ctg ggt gct cag tac aaa 480

Ile Phe Val Asp Val Ala Ser Met Ala Pro Leu Gly Ala Gln Tyr Lys 160

145 150

ctg cta gtt acc agc ctg agc cac ttc cag ctc cgg aca aat gtt ggc 528

Leu Leu Val Thr Lys Leu Lys His Phe Gln Leu Arg Thr Asn Val Ala 175

165 170

ttt cac ttt act gcc agc ccc tct caa cct cag tat ttt cta tac aag 576

Phe His Phe Thr Ala Ser Pro Ser Gln Pro Gln Tyr Phe Leu Tyr Lys 190

180 185

ttt ccc aaa gac gtc gac tca gtt atc att aaa gtc gtc tct gaa atg 624

Phe Pro Lys Asp Val Asp Ser Val Ile Ile Lys Val Val Ser Glu Met 205

195 200

gct tat cca tgt tct gtt gtc tca gtc cag aat atc atg tgc cgc gtc 672

Ala Tyr Pro Cys Ser Val Val Ser Val Gln Asn Ile Met Cys Pro Val 220

210 215

tat gat ctc gac cac aat atg gtc gaa ttt aat ggt gtc tat cag tcc atg 720

Tyr Asp Leu Asp His Asn Val Glu Phe Asn Gly Val Tyr Gln Ser Met 240

225 230

acc aag aaa gct gcc atc acg cta cag aag aag gat ttt cca ggc gag 768

Thr Lys Lys Ala Ala Ile Thr Leu Gln Lys Lys Asp Phe Pro Gly Glu 255

245 250

cag ttc ttc gtc gta ttt gtc ata aag cct gaa gat tat gcc tgt gga 816

Gln Phe Phe Val Val Phe Val Ile Lys Pro Glu Asp Tyr Ala Cys Gly 270

260 265

gga tct ttc ttc atc cag gaa aag gaa aac cag acc tgg aat cta cag 864

Gly Ser Phe Phe Ile Gln Glu Lys Glu Asn Gln Thr Trp Asn Leu Gln 285

275 280

cga aaa aag aac ctt gaa gtc acc att gtc cct tcc att aaa gaa tct 912

Arg Lys Lys Asn Leu Glu Val Thr Ile Val Pro Ser Ile Lys Glu Ser 300

290 295

gtt tat gtc aaa tcc agt att ttc agt gtc ttc atc ttc ctg tcc ttc 960

Val Tyr Val Lys Ser Ser Leu Phe Ser Val Phe Ile Phe Leu Ser Phe 320

305 310 315

WO 01/3434	PCT/US00/34983	WO 01/3434	PCT/US00/34983
tac ttg gga tgc ctt ctt gtt ggg ttt gtt cat tat ctg agg ttt cag Tyr Leu Gly Cys Leu Leu Val Gly Phe Val His Tyr Leu Arg Phe Gln 335 330 335 1008	ttc atg tac atg atc gct ggc ctg tgc atg ctg aag ctc tat cag acc Phe Met Tyr Met Ile Ala Gly Leu Cys Met Leu Lys Leu Tyr Gln Thr 580 585 590 1776		
aga aaa tcc att gat gga agc ttt ggg tcc aat gat ggc tct gga aat Arg Lys Ser Ile Asp Gly Ser Phe Gly Ser Asn Asp Gly Ser Gly Asn 340 345 350 1056	cgc cac cca gac atc aat gcc agc gcc tac tct gcc tat gcc tcc ttt Arg His Pro Asp Ile Asn Ala Ser Ala Tyr Ser Ala Tyr Ala Ser Phe 595 600 605 1824		
atg gtg gca tct cat ccc att gct gcc agc aca ccc gaa ggg agc aat Met Val Ala Ser His Pro Ile Ala Ala Ser Thr Pro Gln Gly Ser Asn 355 360 365 1104	gct gtg gtc atc atg gtc acc gtc ctt gga gtg ttt gga aaa aat Ala Val Val Ile Met Val Thr Val Leu Gly Val Val Phe Gly Lys Asn 610 615 620 1872		
tat ggg aca ata gat gag tca agc tcc agt cct gga agg cag atg tcc Tyr Gly Thr Ile Asp Gln Ser Ser Ser Ser Pro Gly Arg Gln Met Ser 370 375 380 1152	gac gta tgg ttc tgg gtc atc ttc tct gca atc cac gtt ctg gcc tgg Asp Val Thr Phe Thr Val Ile Phe Ser Ala Ile His Val Leu Ala Ser 625 630 635 1920		
tcc tcc gat ggt ggg cca cgg ggc cag tca gac aca gac agc tcc gtg Ser Ser Asp Gly Gly Pro Pro Gly Gln Ser Asp Thr Asp Ser Ser Val 385 390 395 1200	cta gcc ctc agc acc cag ata tat tat atg ggt cgt ttc aag ata gat Leu Ala Leu Ser Thr Gln Ile Tyr Tyr Met Gly Arg Phe Lys Ile Asp 645 650 655 1968		
gag gag agc gac ttc gac acc atg coa gac att gag agt gat aaa aac Glu Gln Ser Asp Phe Asp Thr Met Pro Asp Ile Glu Ser Asp Lys Asn 405 410 415 1248	ttg gga att ttc cgg cgg gct gcc atg gtg ttc tac aca gac tgt atc Leu Gly Ile Phe Arg Arg Ala Ala Met Val Phe Tyr Thr Asp Cys Ile 660 665 670 2016		
atc atc cgg acc aag atg ttc ctt tac ctg tca gat ttg tcc agg aag Ile Ile Arg Thr Lys Met Phe Leu Tyr Leu Ser Asp Leu Ser Arg Lys 420 425 430 1296	cag cag tgt agc cga cct cta tat atg gat aga atg gtg ttg ctg gtt Gln Gln Cys Ser Arg Pro Leu Tyr Met Asp Arg Met Val Leu Leu Val 675 680 685 2064		
gac cgg aga att gtc agc aaa tat aaa att tat ttt tgg aac atc Asp Arg Arg Ile Val Ser Lys Lys Tyr Lys Ile Tyr Phe Thr Asn Ile 435 440 445 1344	gtg ggg aat ctg gtt aac tgg tcc ttc gcc ctc ttt gga ttg ata tac Val Gly Asn Leu Val Asn Thr Ser Phe Ala Leu Phe Gly Leu Ile Tyr 690 700 705 2112		
atc acc att gct gtg ttt tac cgg ctg ccc gtc atc cag ctg gtc att Ile Thr Ile Ala Val Phe Tyr Ala Leu Pro Val Ile Gln Leu Val Ile 450 455 460 1392	cgc ccc agg gac ttt gct tcc tac atg ctg ggc atc ttc atc tgt aac Arg Pro Arg Asp Phe Ala Ser Tyr Met Leu Gly Ile Phe Ile Cys Asn 705 710 715 2160		
acc tat cag aca gtg gta aat gtc act ggc aac cag gac atc tgt tac Thr Tyr Gln Thr Val Val Asn Val Thr Gly Asn Gln Asp Ile Cys Tyr 465 470 475 1440	ctt ttg ctg tac ctg gcc ttt tac atc atc atg aag ctc cgc agc tct Leu Leu Leu Tyr Leu Ala Phe Tyr Ile Ile Met Lys Leu Arg Ser Ser 725 730 735 2208		
tac aac ttc ctc tgt gct cac ccc ttg ggc gtc ctg agt gcc ttc aac Tyr Asn Phe Leu Cys Ala His Pro Leu Leu Gly Val Leu Ser Ala Phe Asn 485 490 495 1488	gaa aag gtc ctc cca gtc cgg ctc ttc tgc atc gtg gcc acc gct gtg Glu Lys Val Leu Pro Val Pro Leu Phe Cys Ile Val Ala Thr Ala Val 740 745 750 2256		
aac att ctc agc aat ctg ggc cac gtg ctt ctg ggc ttc ctc ttc ctg Asn Ile Leu Ser Asn Leu Gly His Val Val Leu Leu Phe Leu 500 505 510 1536	atg tgg gct gcc gcc cta tat ttt ttc ttc cag aat ctc agc agc tgg Met Thr Ala Ala Ala Leu Tyr Phe Phe Phe Gln Asn Leu Ser Ser Thr 755 760 765 2304		
ctg ata gtc ttg cgc cgc gac atc ctc cat cgg aga gcc ctg gaa gcc Leu Ile Val Leu Arg Arg Asp Ile Leu His Arg Arg Ala Leu Glu Ala 515 520 525 1584	gag gga act cgg gcc gaa tcc cgg gag aag aac cgc gag tgc att ctg Glu Gly Thr Pro Ala Glu Ser Arg Glu Lys Asn Arg Glu Cys Ile Leu 770 775 780 2352		
aag gac atc ttt gct gtg gag tac ggg att ccc aaa cac ttt ggt ctc Lys Asp Ile Phe Ala Val Glu Tyr Gly Ile Pro Lys His Phe Gly Leu 530 535 540 1632	ctg gat ttc ttc gat gac cat gac atc tgg cac ttc ctc tct gct act Leu Asp Phe Phe Asp Asp His Asp Ile Thr Phe His Phe Leu Ser Ala Thr 785 790 795 2400		
ttc tac gct atg ggc att gca ttg atg atg gaa ggg gtg ctc agt gct Phe Tyr Ala Met Gly Ile Ala Leu Met Met Glu Gly Val Leu Ser Ala 545 550 555 1680	gct ctg ttt ttc tca ttc ttg gtt ttg tta act ttg gat gat gac ctt Ala Leu Phe Phe Ser Phe Leu Val Leu Leu Thr Leu Asp Asp Leu 805 810 815 2448		
tgc tac cat gtc tgc cct aat tat tcc aac ttc caa ttc gac acc tcc Cys Tyr His Val Cys Pro Asn Tyr Ser Asn Phe Gln Phe Asp Thr Ser 565 570 575 1728	gat gtg gtt cgg aga gac cag atc cct gtc ttc tga acctcaaca Asp Val Val Arg Arg Asp Gln Ile Pro Val Phe 580 585 590 2494		

ttaagagagg ggaaggagagc atcaatcttg gtgctgtttc acaaaatta cagtaccac 2554  
 agcaaaagtaa ccaactgccag atgctccact caccctctgt agagccaggc tctgcattca 2614  
 cacaggaagg agaggggctg cggagattt aaacctgcaa gaaaggaggc agaagggag 2674  
 caatgttttg aggacagacg caaactgag gagctgagaa acacttgctc ctccatctg 2734  
 cagctttggg agtgaacag ggaaggcac tgcatacaag tcaactcacc atcttgggg 2794  
 cctccacc ctaacggaga ctgcoagca atggcagaat gctgctgcac acttccctc 2854  
 aagtgcacc ctgcccacaa aaggccagca gcttgacctt cctgcccaga aactgtgttg 2914  
 gcccttca cacccttgca acactgtg ctccagcaag aggatgtgat tctttagaat 2974  
 atggcgggga ggtgacccca ggcctgccc tactgggata gatgtttcaa tggcaccagc 3034  
 tagtcactc ccagaagaaa ctctgtatat tccccagg ttcttgatgc catcagaagg 3094  
 gctcaggagt ggggtttgtc acacattcct cttaacaagt aactgtcact gggaccgagt 3154  
 cctgggtgct tacatatccc ttctgtctt catc 3188

&lt;210&gt; 4

&lt;211&gt; 827

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 4

Met Arg Gly Cys Leu Arg Leu Ala Leu Leu Cys Ala Leu Pro Trp Leu  
1 5 10 15

Leu Leu Ala Ala Ser Pro Gly His Pro Ala Lys Ser Pro Arg Gln Pro  
20 25 30

Pro Ala Pro Arg Arg Asp Pro Phe Asp Ala Ala Arg Gly Ala Asp Phe  
35 40 45

Asp His Val Tyr Ser Gly Val Val Asn Leu Ser Thr Glu Asn Ile Tyr  
50 55 60

Ser Phe Asn Tyr Thr Ser Gln Pro Asp Gln Val Thr Ala Val Arg Val  
65 70 75 80

Tyr Val Asn Ser Ser Ser Glu Asn Leu Asn Tyr Pro Val Leu Val Val  
85 90 95

Val Arg Gln Gln Lys Glu Val Leu Ser Trp Gln Val Pro Leu Leu Phe  
100 105 110

Gln Gly Leu Tyr Gln Arg Ser Tyr Asn Tyr Gln Glu Val Ser Arg Thr  
115 120 125

Leu Cys Pro Ser Glu Ala Thr Asn Glu Thr Gly Pro Leu Gln Gln Leu  
130 135 140

Ile Phe Val Asp Val Ala Ser Met Ala Pro Leu Gly Ala Gln Tyr Lys  
145 150 155 160

Leu Leu Val Thr Lys Leu Lys His Phe Gln Leu Arg Thr Asn Val Ala  
165 170 175

Phe His Phe Thr Ala Ser Pro Ser Gln Pro Gln Tyr Phe Leu Tyr Lys  
180 185 190

Phe Pro Lys Asp Val Asp Ser Val Ile Ile Lys Val Val Ser Glu Met  
195 200 205

Ala Tyr Pro Cys Ser Val Val Ser Val Gln Asn Ile Met Cys Pro Val  
210 215 220

Tyr Asp Leu Asp His Asn Val Glu Phe Asn Gly Val Tyr Gln Ser Met  
225 230 235 240

Thr Lys Lys Ala Ala Ile Thr Leu Gln Lys Lys Asp Phe Pro Gly Glu  
245 250 255

Gln Phe Phe Val Val Phe Val Ile Lys Pro Glu Asp Tyr Ala Cys Gly  
260 265 270

Gly Ser Phe Phe Ile Gln Glu Lys Glu Asn Gln Thr Trp Asn Leu Gln  
275 280 285

Arg Lys Lys Asn Leu Glu Val Thr Ile Val Pro Ser Ile Lys Glu Ser  
290 295 300

Val Tyr Val Lys Ser Ser Leu Phe Ser Val Phe Ile Phe Leu Ser Phe  
305 310 315 320

Tyr Leu Gly Cys Leu Leu Val Gly Phe Val His Tyr Leu Arg Phe Gln  
325 330 335

Arg Lys Ser Ile Asp Gly Ser Phe Gly Ser Asn Asp Gly Ser Gly Asn  
340 345 350

Met Val Ala Ser His Pro Ile Ala Ala Ser Thr Pro Glu Gly Ser Asn  
355 360 365



WO 01/53454	PCT/US00/34983	WO 01/53454	PCT/US00/34983
Tyr Gly Thr Ile Asp Glu Ser Ser Ser Pro Gly Arg Gln Met Ser 370 375 380		Asp Val Trp Phe Trp Val Ile Phe Ser Ala Ile His Val Leu Ala Ser 635 640	
Ser Ser Asp Gly Gly Pro Pro Gly Gln Ser Asp Thr Asp Ser Ser Val 385 390 395		Leu Ala Leu Ser Thr Gln Ile Tyr Tyr Met Gly Arg Phe Lys Ile Asp 645 650 655	
Glu Glu Ser Asp Phe Asp Thr Met Pro Asp Ile Glu Ser Asp Lys Asn 400 405 410 415		Leu Gly Ile Phe Arg Arg Ala Ala Met Val Phe Tyr Thr Asp Cys Ile 660 665 670	
Ile Ile Arg Thr Lys Met Phe Leu Tyr Leu Ser Asp Leu Ser Arg Lys 420 425 430		Gln Gln Cys Ser Arg Pro Leu Tyr Met Asp Arg Met Val Leu Leu Val 675 680 685	
Asp Arg Arg Ile Val Ser Lys Lys Tyr Lys Ile Tyr Phe Trp Asn Ile 435 440 445		Val Gly Asn Leu Val Asn Trp Ser Phe Ala Leu Phe Gly Leu Ile Tyr 690 695 700	
Ile Thr Ile Ala Val Phe Tyr Ala Leu Pro Val Ile Gln Leu Val Ile 450 455 460		Arg Pro Arg Asp Phe Ala Ser Tyr Met Leu Gly Ile Phe Ile Cys Asn 705 710 715 720	
Thr Tyr Gln Thr Val Val Asn Val Thr Gly Asn Gln Asp Ile Cys Tyr 465 470 475 480		Leu Leu Leu Tyr Leu Ala Phe Tyr Ile Ile Met Lys Leu Arg Ser Ser 725 730 735	
Tyr Asn Phe Leu Cys Ala His Pro Leu Gly Val Leu Ser Ala Phe Asn 485 490 495		Glu Lys Val Leu Pro Val Pro Leu Phe Cys Ile Val Ala Thr Ala Val 740 745 750	
Asn Ile Leu Ser Asn Leu Gly His Val Leu Leu Gly Phe Leu Phe Leu 500 505 510		Met Trp Ala Ala Ala Leu Tyr Phe Phe Phe Gln Asn Leu Ser Ser Trp 755 760 765	
Leu Ile Val Leu Arg Arg Asp Ile Leu His Arg Arg Ala Leu Glu Ala 515 520 525		Glu Gly Thr Pro Ala Glu Ser Arg Glu Lys Asn Arg Glu Cys Ile Leu 770 775 780	
Lys Asp Ile Phe Ala Val Glu Tyr Gly Ile Pro Lys His Phe Gly Leu 530 535 540		Leu Asp Phe Phe Asp Asp His Asp Ile Trp His Phe Leu Ser Ala Thr 785 790 795 800	
Phe Tyr Ala Met Gly Ile Ala Leu Met Met Glu Gly Val Leu Ser Ala 545 550 555		Ala Leu Phe Phe Ser Phe Leu Val Leu Leu Thr Leu Asp Asp Asp Leu 805 810 815	
Cys Tyr His Val Cys Pro Asn Tyr Ser Asn Phe Gln Phe Asp Thr Ser 565 570 575		Asp Val Val Arg Arg Asp Gln Ile Pro Val Phe 820 825	
Phe Met Tyr Met Ile Ala Gly Leu Cys Met Leu Lys Leu Tyr Gln Thr 580 585 590		<210> 5	
Arg His Pro Asp Ile Asn Ala Ser Ala Tyr Ser Ala Tyr Ala Ser Phe 595 600 605		<211> 2484	
Ala Val Val Ile Met Val Thr Val Leu Gly Val Val Phe Gly Lys Asn 610 615 620		<212> DNA	
		<213> Homo sapiens	
		<400> 5	
		atgcgcggct gcctgcggct cgcgcgtctc tcgcgcctgc cctggtctct gctggcgag	60
		tcgcgcgggc acccgcgcaa atcccccagg cagcccccg caccgcgcgc gaccccttc	120

gagcgtgcc gagggcgaga ttctgatacat gttacagcgg ggggtggtgaa cctcagcaacc 180  
 gagaacatct actctttcaa ctacaccagc cagcccgacc aggtgacagc cgtgagggtg 240  
 tatgtgaaca gttctctcga gaattccaac taccgggtcc ttgtgtggtg tgcgcagcag 300  
 aaagagggtc tgcctcggca ggttctctcg ctcttccaag gactatacca gaggaagctac 360  
 aattataag aagtgagcgg cacttatgt cctcagaag caaccaatga gagggagccc 420  
 ttgcagcaac tgatatttgt agatgtcgca tccatggcac ccttgggtgc tcagtacaaa 480  
 ctgcctagta ccaagctgaa gcacttcag ctccggacaa atgttgctt tcaactttact 540  
 gccagccct ctcaacctca gtattttcta tacaagtttc ccaagacgtt ggaactcagtt 600  
 atcattaaag tgggtgtcga aatggcttat ccatgtttctg ttgtctcagt ccagaatatac 660  
 atgtgcccg tgtatgatct cagaccacaaat ggggaattta atgggtctta tcagtccatg 720  
 accaagaag ctgcctatcac gctacagaag aaggattttc caggcgagca gttctctgtg 780  
 gtatttga taaagctga agattatgcc tgtggaggat ctttctcat ccaggaaaaag 840  
 gaaaaccaga cctggaatct acagcgaaaaa agaacccttg aagtgaccat tgcctctcc 900  
 attaaagaat ctgtttatgt gaaatccagt cttttcagtg tottcatctt cctgtccttc 960  
 tacttggat gctctcttgt tgggttgggt cattatctga ggtttcagag aaaaatccatt 1020  
 gatggagct ttgggtccaa tgaatgctct ggaatatgg tggcatctca tcccatgtct 1080  
 gccagcacac ccgaaggagg caattatggg acaatagatg agtcaagctc cagtcctgga 1140  
 aggcagatgt cctctctcga tgggtggcca ccggggccagt cagacacaga cagctccgtg 1200  
 gagggagcgg acttcgacac catgcagac attgagagtg ataaaacat catcgggacc 1260  
 aagatgttcc tttaactgtc agatttgtcc aggaaggacc gagaattgt cagcaaaaaa 1320  
 tataaaattt atttttggaa catcatcacc attgtctgtt tttaagcgtt gccgtgatac 1380  
 cagcttgta ttacctatca gacagtggta aatgtcaacty gaaaccagga catctgttac 1440  
 tacaactcc ttgtgtctca ccccttgggc gtccctgagtg ccttcaacaa catctcagc 1500  
 aatctgggcc acgtgtctct gggcttctct ttccctgtga tagttctgcy ccgcgacatc 1560  
 ctccatcga gagccctgga agccaaggac acttttgtg tggagtagcyg gattcccaaa 1620  
 cactttggtc tcttctacgc tatggcatt gcaattgatga tggaaaggggt gctcagtgct 1680  
 tgcctaccatg ttgcctctaa ttattccaac ttccaattcy acactcctt catgtacatg 1740  
 atcgtctggcc tgtgcagctt gaagctctat cagaccggcc accagacat caatgccagc 1800  
 gctactctg cctatgcctc ctttgtctg gtcatcatgg tcacgtcct tggagtggtg 1860  
 ttggaaaaa atgacgtatg gttctgggtc atcttctcty caatccagct ttggcctcg 1920  
 ctagcctca gaccacagat atattatag gtcgtttca agatagattt gggaaatttc 1980  
 cggcggtgct ccatgtgtgt ctacacagac tgaatccagc agttagcgcc acctctatat 2040

atggatagaa tgggtgtgct ggttggggg aatctggta actggctctt cgcctctttt 2100  
 ggaattatat accgcccacg ggaatttgcct tccacatgc tgggcatctt catctgtaac 2160  
 cttttgtgt acctggcctt ttacatcatc atgaagctcc gcagctctga aaaggtcctc 2220  
 ccagtcgcgc tcttctgcat cgtggccacc gctgtgatgt gggctgcgc cctatattt 2280  
 ttcttcaga atctcagcag ctggaggga actccggcg aatccggga gaagaaccgc 2340  
 gagtgcattc tgcgtgattt cttcgatgac catgacatct ggcacttctt ctctgtact 2400  
 gctctgtttt tctcatctt ggttttcta actttggatg atgacctga tgtggttcgg 2460  
 agagaccaga tctctgtctt ctga 2484

<210> 6  
 <211> 108  
 <212> PRT  
 <213> Homo sapiens  
 <400> 6  
 Val Ile Phe Ser Ala Ile His Val Leu Ala Ser Leu Ala Leu Ser Thr  
 1 5 10 15  
 Gln Ile Tyr Tyr Met Gly Arg Phe Lys Ile Asp Leu Gly Ile Phe Arg  
 20 25 30  
 Arg Ala Ala Met Val Phe Tyr Thr Asp Cys Ile Gln Gln Cys Ser Arg  
 35 40 45  
 Pro Leu Tyr Met Asp Arg Met Val Leu Leu Val Val Gly Asn Leu Val  
 50 55 60  
 Asn Trp Ser Phe Ala Leu Phe Gly Leu Ile Tyr Arg Pro Arg Asp Phe  
 65 70 75 80  
 Ala Ser Tyr Met Leu Gly Ile Phe Ile Cys Asn Leu Leu Leu Tyr Leu  
 85 90 95  
 Ala Phe Tyr Ile Ile Met Lys Leu Arg Ser Ser Glu  
 100 105  
 <210> 7  
 <211> 19  
 <212> PRT  
 <213> Homo sapiens  
 <400> 7  
 Met Arg Gly Cys Leu Arg Leu Ala Leu Leu Cys Ala Leu Pro Trp Leu  
 1 5 10 15  
 Leu Leu Ala  
 <210> 8  
 <211> 808  
 <212> PRT  
 <213> Homo sapiens

&lt;400&gt; 8

Ala Ser Pro Gly His Pro Ala Lys Ser Pro Arg Gln Pro Pro Ala Pro  
1 5 10 15  
Arg Arg Asp Pro Phe Asp Ala Ala Arg Gly Ala Asp Phe Asp His Val  
20 25 30  
Tyr Ser Gly Val Val Asn Leu Ser Thr Glu Asn Ile Tyr Ser Phe Asn  
35 40 45  
Tyr Thr Ser Gln Pro Asp Gln Val Thr Ala Val Arg Val Tyr Val Asn  
50 55 60  
Ser Ser Ser Glu Asn Leu Asn Tyr Pro Val Leu Val Val Val Arg Gln  
65 70 75 80  
Gln Lys Glu Val Leu Ser Trp Gln Val Pro Leu Leu Phe Gln Gly Leu  
85 90 95  
Tyr Gln Arg Ser Tyr Asn Tyr Gln Glu Val Ser Arg Thr Leu Cys Pro  
100 105 110  
Ser Glu Ala Thr Asn Glu Thr Gly Pro Leu Gln Gln Leu Ile Phe Val  
115 120 125  
Asp Val Ala Ser Met Ala Pro Leu Gly Ala Gln Tyr Lys Leu Leu Val  
130 135 140  
Thr Lys Leu Lys His Phe Gln Leu Arg Thr Asn Val Ala Phe His Phe  
145 150 155 160  
Thr Ala Ser Pro Ser Gln Pro Gln Tyr Phe Leu Tyr Lys Phe Pro Lys  
165 170 175  
Asp Val Asp Ser Val Ile Ile Lys Val Val Ser Glu Met Ala Tyr Pro  
180 185 190  
Cys Ser Val Val Ser Val Gln Asn Ile Met Cys Pro Val Tyr Asp Leu  
195 200 205  
Asp His Asn Val Glu Phe Asn Gly Val Tyr Gln Ser Met Thr Lys Lys  
210 215 220  
Ala Ala Ile Thr Leu Gln Lys Lys Asp Phe Pro Gly Glu Gln Phe Phe  
225 230 235 240  
Val Val Phe Val Ile Lys Pro Glu Asp Tyr Ala Cys Gly Gly Ser Phe  
245 250 255  
Phe Ile Gln Glu Lys Glu Asn Gln Thr Trp Asn Leu Gln Arg Lys Lys  
260 265 270  
Asn Leu Glu Val Thr Ile Val Pro Ser Ile Lys Glu Ser Val Tyr Val  
275 280 285  
Lys Ser Ser Leu Phe Ser Val Phe Ile Phe Leu Ser Phe Tyr Leu Gly  
290 295 300  
Cys Leu Leu Val Gly Phe Val His Tyr Leu Arg Phe Gln Arg Lys Ser  
305 310 315 320

Ile Asp Gly Ser Phe Gly Ser Asn Asp Gly Ser Gly Asn Met Val Ala  
325 330 335  
Ser His Pro Ile Ala Ala Ser Thr Pro Glu Gly Ser Asn Tyr Gly Thr  
340 345 350  
Ile Asp Glu Ser Ser Ser Pro Gly Arg Gln Met Ser Ser Ser Asp  
355 360 365  
Gly Gly Pro Pro Gly Gln Ser Asp Thr Asp Ser Ser Val Glu Glu Ser  
370 375 380  
Asp Phe Asp Thr Met Pro Asp Ile Glu Ser Asp Lys Asn Ile Ile Arg  
385 390 395 400  
Thr Lys Met Phe Leu Tyr Leu Ser Asp Leu Ser Arg Lys Asp Arg Arg  
405 410 415  
Ile Val Ser Lys Lys Tyr Lys Ile Tyr Phe Thr Asn Ile Ile Thr Ile  
420 425 430  
Ala Val Phe Tyr Ala Leu Pro Val Ile Gln Leu Val Ile Thr Tyr Gln  
435 440 445  
Thr Val Val Asn Val Thr Gly Asn Gln Asp Ile Cys Tyr Tyr Asn Phe  
450 455 460  
Leu Cys Ala His Pro Leu Gly Val Leu Ser Ala Phe Asn Asn Ile Leu  
465 470 475 480  
Ser Asn Leu Gly His Val Leu Leu Gly Phe Leu Phe Leu Ile Val  
485 490 495  
Leu Arg Arg Asp Ile Leu His Arg Arg Ala Leu Glu Ala Lys Asp Ile  
500 505 510  
Phe Ala Val Glu Tyr Gly Ile Pro Lys His Phe Gly Leu Phe Tyr Ala  
515 520 525  
Met Gly Ile Ala Leu Met Met Glu Gly Val Leu Ser Ala Cys Tyr His  
530 535 540  
Val Cys Pro Asn Tyr Ser Asn Phe Gln Phe Asp Thr Ser Phe Met Tyr  
545 550 555 560  
Met Ile Ala Gly Leu Cys Met Leu Lys Leu Tyr Gln Thr Arg His Pro  
565 570 575  
Asp Ile Asn Ala Ser Ala Tyr Ser Ala Tyr Ala Ser Phe Ala Val Val  
580 585 590  
Ile Met Val Thr Val Leu Gly Val Val Phe Gly Lys Asn Asp Val Trp  
595 600 605  
Phe Trp Val Ile Phe Ser Ala Ile His Val Leu Ala Ser Leu Ala Leu  
610 615 620  
Ser Thr Gln Ile Tyr Tyr Met Gly Arg Phe Lys Ile Asp Leu Gly Ile  
625 630 635 640  
Phe Arg Arg Ala Ala Met Val Phe Tyr Thr Asp Cys Ile Gln Gln Cys  
645 650 655



WO 01/3454 PCT/US00/34983

ccacatatac aacagcagattt gcaattatac atagctagctt aaaaagtgtg gctctgaga 1020 650  
 ttcaacttg tagattttaa aacaagtgc gtacgttaag agaagagcag atcatgtctat 1080  
 tctgacattt gcagagatat acacacactt tttgtacag aagaggttg tctgtgtgtg 1140  
 ggttcgattt atccctgcc accccatcc 1169

<210> 12  
 <211> 2936  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (135) ..(1601)

<400> 12  
 accgccttcg ccggagacct tcaagtcgcy cggctgctcc gagcggcggg ccgcagaggt 60  
 tcaagcagatt ctctctgttc agctctcgga gtactgtgga ttacaggcac gtgcacaac 120  
 accagccac caaa atg cca gaa gag atg gac aag cca ctg atc agc ctg 170  
 Met Pro Glu Glu Met Asp Lys Pro Leu Ile Ser Leu 10  
 1  
 cag ctg gtc gag agc gat agt agc ctt gcc aag gtc ccc gcc gat gag gcc 218  
 His Leu Val Asp Ser Asp Ser Ser Leu Ala Lys Val Pro Asp Glu Ala 15  
 15  
 ccc aaa gtc ggc atc ctg ggt agc ggg gac ttt gcc cgc tcc ctg gcc 266  
 Pro Lys Val Gly Ile Leu Gly Ser Gly Asp Phe Ala Arg Ser Leu Ala 30  
 35  
 40  
 acc cgc ctg gtc ggc tct ggc ttc aaa gtc gtc gtc ggc agc cgc aac 314  
 Thr Arg Leu Val Gly Ser Gly Phe Lys Val Val Val Gly Ser Arg Asn 45  
 50  
 55  
 ccc aaa cgc aca gcc agc ctg tat ccc tca gtc gcc caa gtc act ttc 362  
 Pro Lys Arg Thr Ala Arg Leu Tyr Pro Ser Ala Ala Gln Val Thr Phe 65  
 70  
 caa gag gag gca gtc agc tcc ccg gag gtc atc ttt gtc gct gtc ttc 410  
 Gln Glu Glu Ala Val Ser Ser Pro Glu Val Ile Phe Val Ala Val Phe 80  
 85  
 90  
 cgg gag cac tac tct tca ctg tgc agt ctg agt gac cag ctg ggc ggc 458  
 Arg Glu His Tyr Ser Ser Leu Cys Ser Leu Ser Asp Gln Leu Ala Gly 95  
 100  
 aag atc ctg gtc gat gtc agc aac cct aca gag caa gag cac ctt cag 506  
 Lys Ile Leu Val Asp Val Ser Asn Pro Thr Glu Gln Glu His Leu Gln 110  
 115  
 120  
 cat cgt gag tcc aat gct gag tac ctg gcc tcc ctg ttc ccc act tgc 554  
 His Arg Glu Ser Asn Ala Glu Tyr Leu Ala Ser Leu Phe Pro Thr Cys 125  
 130  
 135  
 aca gtc gtc aag gcc ttc aat gtc atc tct gcc tgg acc ctg cag gct 602  
 Thr Val Val Lys Ala Phe Asn Val Ile Ser Ala Trp Thr Leu Gln Ala 145  
 150  
 155

WO 01/3454 PCT/US00/34983

ggc cca agg gat ggt aac agg cag gtc ccc atc tgc ggt gac cag cca 650  
 Gly Pro Arg Asp Gly Asn Arg Gln Val Pro Ile Cys Gly Gln Pro 160  
 165  
 gaa gcc aag cgt gct gtc tgc gag atg ggc ctc gcc atg ggc ttc atg 698  
 Glu Ala Lys Arg Ala Val Ser Glu Met Ala Leu Ala Met Gly Phe Met 175  
 180  
 185  
 ccc gtc gac atg gga tcc ctg ggc tca gcc tgg gag gtc gag gcc atg 746  
 Pro Val Asp Met Gly Ser Leu Ala Ser Ala Trp Glu Val Glu Ala Met 190  
 195  
 200  
 ccc ctg cgc ctg ctc ccg gcc tgg aag gtc ccc acc ctg ctg gcc ctg 794  
 Pro Leu Arg Leu Leu Pro Ala Trp Lys Val Pro Thr Leu Leu Ala Leu 205  
 210  
 215  
 ggg ctg ttc gtc tgc ttc tat gcc tac aac ttc gtc cgg gag gtt ctg 842  
 Gly Leu Phe Val Cys Phe Tyr Ala Tyr Asn Phe Val Arg Asp Val Leu 225  
 230  
 235  
 cag ccc tat gtc cag gaa agc cag aac aag ttc ttc aag ctg ccc gtc 890  
 Gln Pro Tyr Val Gln Glu Ser Gln Asn Lys Phe Phe Lys Leu Pro Val 240  
 245  
 250  
 tcc gtc gtc aac acc aca ctg ccg gtc gtc gcc tac gtc ctg ctg tca 938  
 Ser Val Val Asn Thr Thr Leu Pro Cys Val Ala Tyr Val Leu Leu Ser 255  
 260  
 265  
 ctg gtc tac ttg ccc ggc gtc gtc gtc gtc gcc ctg cag ctg cgg cgc 986  
 Leu Val Tyr Leu Pro Gly Val Leu Ala Ala Leu Leu Gln Leu Arg Arg 270  
 275  
 280  
 ggc acc aag tac cag cgc ttc ccc gac tgg ctg gac cag tgg cta cag 1034  
 Gly Thr Lys Tyr Gln Arg Phe Pro Asp Trp Leu Asp His Trp Leu Gln 285  
 290  
 295  
 cag cgc aag cag atc ggg ctg ctc agc ttc ttc tgc gcc gcc ctg cag 1082  
 His Arg Lys Gln Ile Gly Leu Leu Ser Phe Phe Cys Ala Ala Leu His 305  
 310  
 315  
 gcc ctg tac agc ttc tgc ttg ccg ctg cgc cgc gcc cag tac gac 1130  
 Ala Leu Tyr Ser Phe Cys Leu Pro Leu Arg Arg Ala His Arg Tyr Asp 320  
 325  
 330  
 ctg gtc aac ctg gca gtc aag cag gtc ttg gcc aac aag agc cag ctc 1178  
 Leu Val Asn Leu Ala Val Lys Gln Val Leu Ala Asn Lys Ser His Leu 335  
 340  
 345  
 tgg gtc gag gag gtc tgg cgc atg gag atc tac ttc tcc ctg gga 1226  
 Trp Val Glu Glu Glu Val Trp Arg Met Glu Ile Tyr Leu Ser Leu Gly 350  
 355  
 360  
 gtc ctg gcc ctc ggc acg ttg tcc ctg ctg gcc ctg acc tca ctg cgg 1274  
 Val Leu Ala Leu Gly Thr Leu Ser Leu Leu Ala Val Thr Ser Leu Pro 365  
 370  
 375  
 tcc att gca aac tgc ctg ctc aac tgg agg gag ttc agc ttc gtt cag tcc 1322  
 Ser Ile Ala Asn Ser Leu Asn Trp Arg Glu Phe Ser Phe Val Gln Ser 385  
 390  
 395  
 tca ctg ggc ttt gtc gcc ctc gtc agc aca ctg cag acg ctg acc 1370  
 Ser Leu Gly Phe Val Ala Leu Val Leu Ser Thr Leu His Thr Leu Thr 400  
 405  
 410

tac ggc tgg acc cgc ttc gag gag agc cgc tac aag ttc tac ctg 1418  
 Tyr Gly Trp Thr Arg Ala Phe Glu Glu Ser Arg Tyr Lys Phe Tyr Leu 425  
 415 420  
 cct ccc acc ttc acg ctc acg ctg ctg gtc ccc tgc gtc gtc atc ctg 1466  
 Pro Pro Thr Phe Thr Leu Thr Leu Leu Val Pro Cys Val Val Ile Leu 440  
 430 435  
 gcc aaa gcc ctg ttt ctc ctg ccc tgc atc agc cgc aga ctc gcc agg 1514  
 Ala Lys Ala Leu Phe Leu Leu Pro Cys Ile Ser Arg Arg Leu Ala Arg 455  
 445 450  
 atc cgg aga ggc tgg gag agg gag agc acc atc aag ttc acg ctg ccc 1562  
 Ile Arg Arg Gly Trp Glu Arg Glu Ser Thr Ile Lys Phe Thr Leu Pro 470  
 465 475  
 aca gac cac gcc ctg gcc gag aag acg agc cac gta tga ggtgctgcc 1611  
 Thr Asp His Ala Leu Ala Glu Lys Thr Ser His Val 485  
 480  
 ctgggtctctg gaccocgggc acacgaggga cgtgcccctg agcccgcttag gttttctttt 1671  
 ctgtgtggtg caaagtggta taactgtgtg caaataggag gtttgaggtc caaattctctg 1731  
 ggaotcaaat gtatgcagta ctattcagaa tgatatcac acatatgtgt atattgtattt 1791  
 acatatatc cacatatata acaggaattg caattatata tagctagcta aaagtgtggg 1851  
 ttctgtgat ttcaacttgt agatttaaaa acaagtgcg tagcttaaga gaagagaga 1911  
 taatgtcatt gtgacattg cagagatata cacacacttt ttgtacagaa gaggtgtgtg 1971  
 ctgtgtgtggg ttcgatttat cctgcgccac cccatcccac caacttccct ttgtctactt 2031  
 ccccaaggct ctgacagagc tagggctctg aaggggaggg aaggcaacgg ctctgccacg 2091  
 agcatccctt ggagcatgtg agcagcggct ggtctctctc ctccacctgg ggcagcagea 2151  
 ggaggcctgg gggggaggaa aatcaggcag tgggctgga gtctgtcctt ggtcctttgc 2211  
 ccggtgttg gaggatggag ggaattgggt gaagctgtc cactctacc ttgtgtgagt 2271  
 ggggagacat ttctcctgaa agtcagaagt caccatagag cctgcacaatg gatctctctg 2331  
 tgagagtac gtcaactctt ttccagacc attagtgc ctggctggg aacaagtga 2391  
 atttccctcc ctctcttaac ctggcgatga ggtctcttta aaccactgtg ccttctcacc 2451  
 ctttccatct tcaatttgaa cgaactccag gaaggcctag agcagaccct ttagaatca 2511  
 gcccaagggg gagagcaaga gaaaactc tagggagtaa agctcccgg gcgtcagagt 2571  
 tgagcctgc ctgggctgaa ggaactctt caggaatca gtcttgagga aaatattgg 2631  
 ggactccaaa tgtctctctg cagaggacc agaaaaccac actggctcca acttctcct 2691  
 catggggcat kacactcaa aacagtggg agcaacttt ccaccaagc tacaacctta 2751  
 aaatgtctgt gcccaaacg acaagaggga agagaccgc cgggggcaca ggaagtctgt 2811  
 cctccagta caggccatcc ttgtgtctcc ctactgact tagttactt cccctgtgaa 2871

gaacaggtg ttctggctg agccccaac cctctgcaga accaggttga tctgccacag 2931  
 aaaaa 2936  
 <210> 13  
 <211> 488  
 <212> PRT  
 <213> Homo sapiens  
 <400> 13  
 Met Pro Glu Glu Met Asp Lys Pro Leu Ile Ser Leu His Leu Val Asp 15  
 1 5 10  
 Ser Asp Ser Ser Leu Ala Lys Val Pro Asp Glu Ala Pro Lys Val Gly 30  
 20 25  
 Ile Leu Gly Ser Gly Asp Phe Ala Arg Ser Leu Ala Thr Arg Leu Val 45  
 35 40  
 Gly Ser Gly Phe Lys Val Val Val Gly Ser Arg Asn Pro Lys Arg Thr 60  
 50 55  
 Ala Arg Leu Tyr Pro Ser Ala Ala Gln Val Thr Phe Gln Glu Glu Ala 80  
 65 70 75  
 Val Ser Ser Pro Glu Val Ile Phe Val Ala Val Phe Arg Glu His Tyr 95  
 85 90  
 Ser Ser Leu Cys Ser Leu Ser Asp Gln Leu Ala Gly Lys Ile Leu Val 110  
 100 105  
 Asp Val Ser Asn Pro Thr Glu Gln Glu His Leu Gln His Arg Glu Ser 125  
 115 120  
 Asn Ala Glu Tyr Leu Ala Ser Leu Phe Pro Thr Cys Thr Val Val Lys 140  
 130 135 140  
 Ala Phe Asn Val Ile Ser Ala Trp Thr Leu Gln Ala Gly Pro Arg Asp 160  
 145 150 155  
 Gly Asn Arg Gln Val Pro Ile Cys Gly Asp Gln Pro Glu Ala Lys Arg 175  
 165 170  
 Ala Val Ser Glu Met Ala Leu Ala Met Gly Phe Met Pro Val Asp Met 190  
 180 185  
 Gly Ser Leu Ala Ser Ala Trp Glu Val Glu Ala Met Pro Leu Arg Leu 205  
 195 200

Leu Pro Ala Trp Lys Val Pro Thr Leu Leu Ala Leu Gly Leu Phe Val  
210 215 220

Cys Phe Tyr Ala Tyr Asn Phe Val Arg Asp Val Leu Gln Pro Tyr Val  
225 230 235 240

Gln Glu Ser Gln Asn Lys Phe Lys Leu Pro Val Ser Val Val Asn  
245 250 255

Thr Thr Leu Pro Cys Val Ala Tyr Val Leu Leu Ser Leu Val Tyr Leu  
260 265 270

Pro Gly Val Leu Ala Ala Leu Gln Leu Arg Arg Gly Thr Lys Tyr  
275 280 285

Gln Arg Phe Pro Asp Trp Leu Asp His Trp Leu Gln His Arg Lys Gln  
290 295 300

Ile Gly Leu Leu Ser Phe Phe Cys Ala Ala Leu His Ala Leu Tyr Ser  
305 310 315 320

Phe Cys Leu Pro Leu Arg Arg Ala His Arg Tyr Asp Leu Val Asn Leu  
325 330 335

Ala Val Lys Gln Val Leu Ala Asn Lys Ser His Leu Trp Val Glu Glu  
340 345 350

Glu Val Trp Arg Met Glu Ile Tyr Leu Ser Leu Gly Val Leu Ala Leu  
355 360 365

Gly Thr Leu Ser Leu Leu Ala Val Thr Ser Leu Pro Ser Ile Ala Asn  
370 375 380

Ser Leu Asn Trp Arg Glu Phe Ser Phe Val Gln Ser Ser Leu Gly Phe  
385 390 395 400

Val Ala Leu Val Leu Ser Thr Leu His Thr Leu Thr Tyr Gly Trp Thr  
405 410 415

Arg Ala Phe Glu Glu Ser Arg Tyr Lys Phe Tyr Leu Pro Pro Thr Phe  
420 425 430

Thr Leu Thr Leu Leu Val Pro Cys Val Val Ile Leu Ala Lys Ala Leu  
435 440 445

Phe Leu Leu Pro Cys Ile Ser Arg Arg Leu Ala Arg Ile Arg Arg Gly  
450 455 460

Trp Glu Arg Glu Ser Thr Ile Lys Phe Thr Leu Pro Thr Asp His Ala  
465 470 475 480

Leu Ala Glu Lys Thr Ser His Val  
485

<210> 14

<211> 1467

<212> DNA

<213> Homo sapiens

<400> 14

atgccagaag agatggacaa gccactgac agcctccacc tggtagacag ctagtagc 60  
cttgcgaagg tccccgatga ggcctccaaa gtgggcatcc tgggtagcgg ggaattgcc 120  
cgtctccctgg ccacacgctt ggtgggtctt ggttcaaaag tgggtggtgg gagccgaac 180  
cccaaacgca cagccaggct gtaacctca ggcgcccaag tgactttcca agaggagca 240  
gtagactccc cggagggtcat ctttgtggt gtgttccggg agcaactactc ttaactgtgc 300  
agtctcaatg accagctggc gggcaagatc ctgggtggatg tagaacaacc tacagagcaa 360  
gagcaccttc agcatcgtga gtccaatgct gactactcgg cctccctctt cccacttgc 420  
acaagtggca aggccttcaa tgcatactct gcttggacc tgcaggctgg cccaaggat 480  
ggtaacaggc aggtgcccac ctgcggtgac cagccagaag ccaagcgtgo tctctcgag 540  
atggcgtcgt ccaatgggtt catgcctgt gacatgggt ccttgggtc agctgggag 600  
gtggaggcca tgcctctgc cctctcccg gcttggaaag tgcacacct gttggcctg 660  
gggtctctcg tctgtttcta tgctacaac ttcttccggg acgtttctga gccctatgtg 720  
caggaaagcc agaacaagtt cttcaagctg cctgtgtccg tggtaaacac cacactgctg 780  
tggtggcctt agtgtctgt gtaactcgtg taactgtccg gctgtctggc ggtgacctg 840  
cagctgcggc ggcggaccaa gtaccagcgc ttcccagact ggtggacca ctggctacag 900  
caccgcaagc agatcgggct gctcagcttc ttctgcgcg ccttgcacgc cctctacagc 960  
ttctgtctgc cgtgtgcgcg cgcacacgc tagacactgg tcaacctggc agtaagagag 1020  
gtcttggcca acaagagcca cctctgggtg gaggaggagg tctggcgat ggagatctac 1080  
ctctccctgg gagtgtctgg cctcggcacg ttgtcctcgc tggcctgac ctacactcgc 1140  
tcattgcaa actcgtcaa ctggaggagg ttacgtctcg ttacgtctc actgggtttt 1200  
gtggcctcgt tcttgagcac actgcacag ctcacctag gctggacctg cgccttgag 1260  
gagagcctt acaagtctta cctgcctccc accttcacgc taactgtct ggtgctctgc 1320  
gtcgtcatcc tggccaaagc cctgtttctc ctgccttcca tcaagcgcag actcgcagg 1380  
atccggagag gctgggagag ggagagcacc atcaagtcca cgtgcccc agaccagcc 1440

ctggccgaga agacagacca cgtatga

<210> 15

<211> 237

<212> PRT

<213> Homo sapiens

<400> 15

Val Pro Thr Leu Leu Ala Leu Gly Leu Phe Val Cys Phe Tyr Ala Tyr  
1 5 10 15

Asn Phe Val Arg Asp Val Leu Gln Pro Tyr Val Gln Glu Ser Gln Asn  
20 25 30

Lys Phe Phe Lys Leu Pro Val Ser Val Val Asn Thr Thr Leu Pro Cys  
35 40 45

Val Ala Tyr Val Leu Leu Ser Leu Val Tyr Leu Pro Gly Val Leu Ala  
50 55 60

Ala Ala Leu Gln Leu Arg Arg Gly Thr Lys Tyr Gln Arg Phe Pro Asp  
65 70 75 80

Trp Leu Asp His Trp Leu Gln His Arg Lys Gln Ile Gly Leu Leu Ser  
85 90 95

Phe Phe Cys Ala Ala Leu His Ala Leu Tyr Ser Phe Cys Leu Pro Leu  
100 105 110

Arg Arg Ala His Arg Tyr Asp Leu Val Asn Leu Ala Val Lys Gln Val  
115 120 125

Leu Ala Asn Lys Ser His Leu Trp Val Glu Glu Val Trp Arg Met  
130 135 140

Glu Ile Tyr Leu Ser Leu Gly Val Leu Ala Leu Gly Thr Leu Ser Leu  
145 150 155 160

Leu Ala Val Thr Ser Leu Pro Ser Ile Ala Asn Ser Leu Asn Trp Arg  
165 170 175

Glu Phe Ser Phe Val Gln Ser Ser Leu Gly Phe Val Ala Leu Val Leu  
180 185 190

Ser Thr Leu His Thr Leu Thr Tyr Gly Trp Thr Arg Ala Phe Glu Glu  
195 200 205

Ser Arg Tyr Lys Phe Tyr Leu Pro Pro Thr Phe Thr Leu Thr Leu Leu  
210 215 220

Val Pro Cys Val Val Ile Leu Ala Lys Ala Leu Phe Leu  
225 230 235

<210> 16

<211> 146

<212> PRT

<213> Homo sapiens

<400> 16

Ala Arg Gly Gly Arg Leu Arg Trp Arg Arg Leu Asp Cys Leu Ser  
1 5 10 15

Ala Ala Glu Ser Asp Thr Val Ala Tyr Glu Asp Leu Ser Glu Asp Tyr  
20 25 30

Thr Gln Lys Lys Trp Lys Gly Leu Ala Leu Ser Gln Arg Ala Leu His  
35 40 45

Trp Asn Met Met Leu Glu Asn Asp Arg Ser Met Ala Ser Leu Ala Gly  
50 55 60

Arg Asn Met Met Glu Ser Ser Glu Leu Thr Pro Lys Gln Glu Ile Phe  
65 70 75 80

Lys Gly Ser Glu Ser Ser Asn Ser Thr Ser Gly Gly Leu Phe Gly Val  
85 90 95

Val Pro Gly Gly Thr Glu Thr Gly Asp Val Cys Glu Asp Thr Phe Lys  
100 105 110

Glu Leu Glu Gly Gln Pro Ser Asn Glu Glu Gly Ser Arg Leu Glu Ser  
115 120 125

Asp Phe Leu Glu Ile Ile Asp Glu Asp Lys Lys Lys Ser Thr Lys Asp  
130 135 140

Arg Tyr

145

<210> 17

<211> 364

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1)..(364)

<223> n = A, T, G, or C

<400> 17

cggacgcgtg ggctcggggc aggtttccgg gagcagatgc caaaaagact ttctcataga  
60

gaagaggctt tctttgttag agacagaata aaaataatg gtatgtttct gttgttccc  
120

tccccctccc ccttgtgtga taccacatgt statagtatt taagtgaac tcaagccctc  
180

aaggcccaac ttctgtctct atattgtaat atagaatttc gaagagacat ttctactttt  
240

tacacattgg gcacaaagat aagctttgat taaagtagta agtaaaagc tacctaggaa  
300

atacttcagt gaattctaac mnnnnnnnnn mnnnnnnnn mnnnnnnnn mnnnnngaaa  
360

cagg

<210> 18

<211> 5773

<212> DNA

<213> Homo sapiens

<400> 18



ttctcgtatc atagctgggg gagctgagc gtgggagcgg tgcgcagct cctcctgaa 60  
 aacgcgaat gagctctgct tggctctccc tccactgggg gtgagagccc ctgccagga 120  
 ggccacgagc aaatggcccc atagtgaaa ctgggaagct tttaggcato tgcacagagc 180  
 gggagccagc cgggggagca cagtgctgga caggccnacc aactcaaat tgaagacatg 240  
 aaatccccc aaagaaccac ttgtgctc atgtttattg tgatttatto ttccaaagt 300  
 gcactgaact ggaattacga gtctactatt catcctttga gtctctatga acatgaacca 360  
 gctggtgaag aggcactgag gcaaaaacga gcggttgcca caaaaagtc taccgttgaa 420  
 gaatacactg ttaatatga gatcagtttt gaaaatgcac ccttcctgga tccatcaaaa 480  
 gctacttga acagccctcag ttctccaat catgggaata acactgacca aattactgac 540  
 attttgaca taaatgtgac aacagtctgc agactgctg gaaatgaaat ctggtgctcc 600  
 tgcgagacag gttatgggtg gcttcgggaa aggtgtcttc acaatctcat ttgtcaagag 660  
 cgtgactct tccctccagg gcaccattgc agtgcctta aagaactgccc tcccaatgga 720  
 ccttttgc tgcctcagga agatgtacc ctgaacatga gagtcagact aaatgtaggc 780  
 ttccaagaag accctcatgaa cactctccc gccctctata ggtctacaa gaccgacttg 840  
 gaaacagcgt tccgggaagg ttacggaatt ttaccggct tcaaggcgct gactgtgaca 900  
 ggggtcaggt ctggaagctg ggttgtaca tatgaatga agactacccc accatcactt 960  
 gagttaac ataaagccaa tgaacaagtt gtacagagcc tcaatcagac ctacaaaatg 1020  
 gactacaact ccttcaagc agtctatc atgaagaaga attctttgt cacacagaa 1080  
 atcatctttg aaggggacac agtcagctg gtgtgtgaaa aggaagtitt gtctccaat 1140  
 gtgtcttggc gctatgaaga acagcagttg gaaatccaga acagcagcag attctcgatt 1200  
 taacccgac ttctcaaaa catgacttgg gtgtccnagc tcaactacca caacatcact 1260  
 ccaggtgatg caggtgaata tgtttgcaa ctgatattag acatttttga atatgagtgc 1320  
 aagaagaaaa tagatgtat gccatccaa abtttgcaa atgaagaaat gaaggtgatg 1380  
 tgcgacaaac atcctgtatc tttagactgc tgcagtcagg gtaatgttaa ttggagcaaa 1440  
 gtagaatgga agcaggaagg aaaaataaat attccaggaa cccctgagac agacatagat 1500  
 tctagctgca gcagatacac cctcaaggct gatggaaacc agtgcccaag cgggtcgtct 1560  
 ggaacaacag tcaatcacac ttgtgagttc atcagtgcct atggagccag aggcagtga 1620  
 aacataaag tgaattcat ctctgtgccc aatcaacaa taaccocgga cccaatttct 1680  
 gttctgagg gacaaaact ttctataaaa tgcatacgtg atgtgagtaa ctatgatgag 1740  
 gtttatgga acactctgc tggaaataaa atataccaaa gattttatc cagcaggagg 1800  
 tatcttgatg gagcagaac agtctgaca gtcaagacct cgaccaggga gtggaatgga 1860  
 acctatcact gcatatttag atataagaat tcatcacgta ttgaaccaa agacgtcatt 1920

gtccaccgc tgcctctaaa gctgaacatc atggttgatc ctttgaagc tactgtttca 1980  
 tgcagtggtt cccatacat caagtctgc atagaggagg atggagacta caaagttaact 2040  
 ttccatatgg gtctctate cctctctgct gcaaaagag ttaacaaaa acaagtgtgc 2100  
 taacaaaca atttcaatgc aagctcagtt tctgttgtt caaaaactgt tgaatgtgtc 2160  
 tgtcaactta ccaatgctgc taataattca gtttggagcc catctatgaa gctgaatctg 2220  
 gtctctgggg aaaaatcac atgccaggat cccgtaatat gtgtcggaga gccggggaaa 2280  
 gtctccaga agctatgcgg gttctcaac gtccacaga gccctgagga gtcccatagg 2340  
 gggggaccat cacttcaaaa tgtgtaggct cccagtgggg ggtagaagag aaatgactgc 2400  
 atctctgcc caataaacag tctgtccag atggttaagg ctttgatcaa gagccctct 2460  
 caggatgaga tgcctccctac atacctgaag gatctttcta ttagcataga caaaggggaa 2520  
 catgaaata gctcttctcc tgggagctcg ggagccattt ttaacatcct tgatctgctc 2580  
 tcaacagttc caacccaagt aaatcagaa atgatgacgc acgtgctctc taogtttaat 2640  
 gtcatcttg gcaagccgct ctgaaacac tgggaagttt tacaacagca atggaccaat 2700  
 cagagttcac agtactaca ttcagtggaa agattttccc aagcattaca gtcaggagat 2760  
 agccctctt tgcctctctc ccaactaat gtgcagatga gcagcaggtt aatcaagtcc 2820  
 agccaccag aaactatca acagaggttt gtttcccat acttgacct ctggggcaat 2880  
 gtgtctatg caaagagcta cctagaanaa ttgcagtcgg attcgtctat tgcaccatg 2940  
 gctttccaa ctctccagg catctctgct caggatctcc aggaataaa cttgcagag 3000  
 agcttagtga tgaacacac tgcagccac aatacagcta tgcattcag gatttcaatg 3060  
 acttttaaga acaatagccc ttcagggggc gaaacgaagt gtgtctctcg gaacttcagg 3120  
 cttgccaca acacaggggg gtgggacagc agtgggtgct atgttgaga aggtgatggg 3180  
 gacaatgca cctgtatctg tgaccacta acatcattct ccatcctcat gtccctgac 3240  
 tcccagatc ctagtctct cctgggaata ctctggata ttatttcta tgttgggtg 3300  
 ggcttttcaa tcttgagctt ggcagcctgt ctagtgtgtg aagctgtgtg gtggaaatcg 3360  
 gtgaccaaga atcggaactt ttatatgac cacactgca tagtgaatat cgtgctctcc 3420  
 cttctgggtc gccaaacact ggttcattgg ggtcgtgccc atccaggaca atcgctacat 3480  
 actctgaag acagcctgtg tggctgccc cttctcact cactttctct acctcagcgt 3540  
 cttctctgg atgtgacac tgggcctcat gctgtctcat cgcctggttt tcatctgca 3600  
 tgaacaagc aggtccactc agaaagccat tgccttctgt cttggctatg gctgcccact 3660  
 tgcctatctg gtcatacgc tgggagccac ccagcccccg gaagtctata cggaggagaa 3720  
 tgtctgttg ctcaactggg aggcaccaa ggcctcgtg gctttcgcca tcccagcact 3780

WO 01/3444 PCT/US00/34983  
gacatctgtg gtgggtgaaca taaccatcac tatttgttc atcacaaga tccctgagcc 3840  
ttccattgga gacaagccat gcaagcagga gaagcagc cgtgttcaga tcagcaagag 3900  
cattgggttc ctacacacac tcttgggct cacttggggt ttgtgtctca ccaactgttt 3960  
cccaggacc aacctgtgt tccatcatc atttgccatc ctcaatgtct tccagggtatt 4020  
attcatatta ctctttggat gctctggga tctgaagga caggaaagctt tgcgaataa 4080  
gttttcattg tcgagatggt ctccacagca ctcaaagtc acaaccctgg gtccatccac 4140  
acctgtgttt tctatgagtt ctccaatc aaggagattt aacaatttgt ttggtaaac 4200  
aggaacgtat aatgtttcca ccccaagc aaccagtc tccctggaaa actcatccag 4260  
tgtctcttcg ttgtccaact aagaacagga taatccaacc tacgtgacct ccgaggga 4320  
gtgctgtgc ttttaaaag agatgcttgc aaagcaatgg ggaactgttt ctcgggcag 4380  
gtttccggga gcagatgcca aaaaagcttt tccatagaga agaggcttcc ttgttaag 4440  
acagaataaa aataatgttt atgttttgt ttgttccctc cccctccccc ttgtgtata 4500  
ccacatgtgt atagatttta agtgaactc aagccctcaa ggcaccaact ctctgtctat 4560  
attgtaatat agaatttcga agagacattt tcaacttita cacattgggc acaagataa 4620  
gctttgatta aagtagtaag taaaggcta cctaggaat acttcagtga atttaagaa 4680  
ggaaggagg aagaagaa ggaagaagg gaggaaca gggagaagg gaaagaag 4740  
aaaaagaa agatgaanaa aggaacaaat aagacaaac aacattaagg gccatttgt 4800  
aagatttcca tgttaagat ctaataaat cactcagtc aacattgaga atttttttt 4860  
aatggctcaa aaatggaac tgaagcaag tcatgggaa tgaatacttt ggcagatc 4920  
ttctctatgt ctctctagct agaggagga aaaaaggct gaaaaatag ggaagaaatt 4980  
cttccatcag aacgacttca agtgataac aatatttata agaatgaat ggaaggaaat 5040  
atgatoctcc tgagactaac ttgtgttgt aaggttttga ctaagtgaat gtaactgcag 5100  
aggaagtatt acaagatat gtcattagat ccaagtctg attaaatttt tatagtttat 5160  
cagaaaagcc ttatatitta gttgttcca cttttgaaa gcaaaaaata tatattgat 5220  
ataccctca attgccaaat ttgatatgtt gcactgaaga cagaccctgt catatattta 5280  
atggcttcaa gcaggtaact ctctgtcat tatagaatag attttaataa tcttatagca 5340  
ttgtatatta ttatttgtgt tgcactgtt attattattg tggatactgg cccctgtgtt 5400  
gtgtcatagc tccctatgta ttctctgttt ccaactttaa gttccagac caatacatc 5460  
taagagtttt gcattgtcta aatttgttt attccaacca cgtggaaagc tccctggaag 5520  
aaattttaca ttccgttgtt ctgtgtctct aatgacactt gacctgtgtt acaaatggc 5580  
agagcccttc ccaaggattt gattgttgtt gaattatctg catgtgtgtt ttttttgtt 5640  
gtgtatgtat ttcattaaaa aatataata cttatgaaa ttgcaogcat attagattta 5700

WO 01/3444 PCT/US00/34983  
accatgtact attgatacag caagctaca ttgcaataa aagtcgcgac ccaaaaggag 5760  
aatgagacaa aaa 5773

<210> 19  
<211> 5714  
<212> DNA  
<213> Homo sapiens  
<220>  
<221> CDS  
<222> (272) .. (4312)  
<400> 19  
ctgcacagacc ggtccgaat tcccggtcgc acgatttctg gatcatagct gggggagggt 60  
gagcgtggga gcggtgctgc cagtcctgcc tgaacgcgc aatgagctct tgettggttc 120  
tccctccact gggcgtgaga gccctgccc agggagccca ggacaaatgg ccccatagt 180  
gaaactggga agcttttagg catctgatca gagcgggagc cagccggggg accacagtgc 240  
tggaacggcc aaccaactca aactgaaga c atg aaa tcc oca agg aga acc 292  
Met Lys Ser Pro Arg Arg Thr 1 5  
act ttg tgc ctc atg ttt att gtg att tat tct tcc aaa gct gca ctg 340  
Thr Leu Cys Leu Met Phe Ile Val Ile Tyr Ser Ser Lys Ala Ala Leu 20  
aac tgg aat tac gag tct act att cat cct ctg agt ctt cat gaa cat 388  
Asn Trp Asn Tyr Glu Ser Thr Ile His Pro Leu Ser Leu His Glu His 35  
gaa cca gct ggt gaa gag gca ctg agg caa aaa cga gcc gtt gcc aca 436  
Glu Pro Ala Gly Glu Glu Ala Leu Arg Glu Lys Arg Ala Val Ala Thr 55  
aaa agt cct acg gct gaa gaa tac act gtt aat att gag atc agt ttt 484  
Lys Ser Pro Thr Thr Ala Glu Glu Tyr Thr Val Asn Ile Glu Ile Ser Phe 65  
gaa aat gca tcc ttc ctg gat cct atc aaa gcc tac ttg aac agc ctc 532  
Glu Asn Ala Ser Phe Leu Asn Pro Ile Lys Ala Tyr Leu Asn Ser Leu 85  
agt ttt cca att cat ggg aat aac act gac caa att act gac att ttg 580  
Ser Phe Pro Ile His Gly Asn Asn Thr Asp Glu Ile Thr Asp Ile Leu 95  
agc ata aat stg aca aca stc tgc tgc aga cct gct gga aat gaa atc tgg 628  
Ser Ile Asn Val Thr Thr Val Cys Arg Pro Ala Gly Asn Glu Ile Trp 115  
tgc tcc tgc gag aca ggt tat ggg tgg cct cgg gaa agg tgt ctt cac 676  
Cys Ser Cys Glu Thr Gly Tyr Gly Trp Pro Arg Glu Arg Cys Leu His 135  
aat ctc att tgt caa gag cgt gac stc ttc ctc cca ggg cac cat tgc 724  
Asn Leu Ile Cys Glu Glu Arg Asp Val Phe Leu Pro Gly His His Cys 145  
140 150

agt tgc ctt aaa gaa cag cct ccc aat gga cct ttt tgc ctg ctt cag 772  
 Ser Cys Leu Lys Glu Leu Pro Pro Asn Gly Pro Phe Cys Leu Leu Gln 165  
 155  
 gaa gat gtt acc ctg aac atg aga gtc aga cta aat gta ggc ttt caa 820  
 Glu Asp Val Thr Leu Asn Met Arg Val Arg Leu Asn Val Gly Phe Gln 180  
 170 175  
 gaa gac ctc atg aac act tcc tcc gcc ctc tat agg tcc tac aag acc 868  
 Glu Asp Leu Met Asn Thr Ser Ser Ala Leu Tyr Arg Ser Tyr Lys Thr 195  
 185 190  
 gac ttg gaa aca cgc ttc cgg aag ggt tac gga att tta cca ggc ttc 916  
 Asp Leu Glu Thr Ala Phe Arg Lys Gly Tyr Gly Ile Leu Pro Gly Phe 215  
 200 205 210  
 aag ggc gtg act gtg aca ggg ttc aag tct gga agt gtg gtt gtg aca 964  
 Lys Gly Val Thr Val Thr Gly Phe Lys Ser Gly Ser Val Val Thr 230  
 220 225  
 tat gaa gtc aag act aca cca tca cta ctt gag tta ata cat aaa gcc 1012  
 Tyr Glu Val Lys Thr Thr Pro Pro Ser Leu Glu Leu Ile His Lys Ala 245  
 235 240  
 aat gaa caa gtt gta cag agc ctc aat cag acc tac aaa atg gac tac 1060  
 Asn Glu Gln Val Val Gln Ser Leu Asn Gln Thr Tyr Lys Met Asp Tyr 260  
 250 255  
 aac tcc ttt caa gca gtt act aat gaa agc aat ttc ttt gtc aca 1108  
 Asn Ser Phe Gln Ala Val Thr Ile Asn Glu Ser Asn Phe Val Thr 275  
 265 270  
 cca gaa atc atc ttt gaa ggg gac aca gtc agt ctg gtg tgt gaa aag 1156  
 Pro Glu Ile Ile Phe Glu Gly Asp Thr Val Ser Leu Val Cys Glu Lys 295  
 280 285 290  
 gaa gtt ttg tcc tcc aat gtg tct tgg cgc tat gaa gaa cag cag ttg 1204  
 Glu Val Leu Ser Ser Asn Val Ser Trp Arg Tyr Glu Glu Gln Gln Leu 310  
 300 305  
 gaa atc cag aac agc agc aga ttc cag att tac acc gca ctt ttc aac 1252  
 Glu Ile Gln Asn Ser Ser Arg Phe Ser Ile Tyr Thr Ala Leu Phe Asn 325  
 315 320  
 aac atg act ctg gtg tcc aag ctc acc atc cca aac atc act cca ggt 1300  
 Asn Met Thr Ser Val Ser Lys Leu Thr Ile His Asn Ile Thr Pro Gly 340  
 330 335 340  
 gat gca ggt gaa tat gtt tgc aca ctg ata tta gac att ttt gaa tat 1348  
 Asp Ala Gly Glu Tyr Val Cys Lys Leu Ile Leu Asp Ile Phe Glu Tyr 355  
 345 350 355  
 gag tgc aag aag aaa ata gat gtt atg ccc atc caa att ttg gca aat 1396  
 Glu Cys Lys Lys Lys Ile Asp Val Met Pro Ile Gln Ile Leu Ala Asn 375  
 360 365 370  
 gaa gaa atg aag gtg atg tgc gac aac aat cct gta tct ttg aac tgc 1444  
 Glu Glu Met Lys Val Met Cys Asp Asn Pro Val Ser Leu Asn Cys 390  
 380 385 390  
 tgc agt cag ggt aat gtt aat tgg agc aaa gta gaa tgg aag cag gaa 1492  
 Cys Ser Gln Gly Asn Val Asn Trp Ser Lys Val Glu Trp Lys Gln Glu 400

gga aaa ata aat att cca gga acc cct gag aca gac ata gat tct agc 1540  
 Gly Lys Ile Asn Ile Pro Gly Thr Pro Glu Thr Asp Ile Asp Ser Ser 420  
 410 415  
 tgc agc aga tac acc ctc aag gct gat gga acc cag tgc cca agc ggg 1588  
 Cys Ser Arg Tyr Thr Leu Lys Ala Asp Gly Thr Gln Cys Pro Ser Gly 435  
 425 430 435  
 tgc tct gga aca aca gtc atc tac act tct gag ttc atc agt gcc tat 1636  
 Ser Ser Gly Thr Thr Val Ile Tyr Thr Cys Glu Phe Ile Ser Ala Tyr 455  
 440 445 450  
 gga gcc aga ggc agt gca aac ata aaa atg aca ttc atc tct gtg gcc 1684  
 Gly Ala Arg Gly Ser Ala Asn Ile Lys Val Thr Phe Ile Ser Val Ala 470  
 460 465 470  
 aat ata aca ata acc ccg gac cca att tct gtt tct gag gga caa aac 1732  
 Asn Leu Thr Ile Thr Pro Asp Pro Ile Ser Val Ser Glu Gly Gln Asn 485  
 475 480  
 ttt tct ata aaa tgc atc agt gat gtg agt aac tat gat gag gtt tat 1780  
 Phe Ser Ile Lys Cys Ile Ser Asp Val Ser Asn Tyr Asp Glu Val Tyr 500  
 490 495  
 tgg aac act tct gct gga att aaa ata tac caa aga ttt tat acc acg 1828  
 Trp Asn Thr Ser Ala Gly Ile Lys Ile Tyr Gln Arg Phe Tyr Thr Thr 515  
 505 510 515  
 agg agg tat ctt gat gga gca gaa tca gta ctg aca gtc aag acc tcy 1876  
 Arg Arg Tyr Leu Asp Gly Ala Glu Ser Val Leu Thr Val Lys Thr Ser 535  
 520 525 530  
 acc agg gag tgg aat gga acc tat cac tgc ata ttt aga tat aag aat 1924  
 Thr Arg Glu Trp Asn Gly Thr Tyr His Cys Ile Phe Arg Tyr Lys Asn 550  
 540 545 550  
 tca tac agt att gca acc aaa gac gtc att gtt cac ccg ctg cct cta 1972  
 Ser Tyr Ser Ile Ala Thr Lys Asp Val Ile Val His Pro Leu Pro Leu 565  
 555 560 565  
 aag ctg aac atc atg gtt gat cct ttg gaa gct act gtt tca tgc agt 2020  
 Lys Leu Asn Ile Met Val Asp Pro Leu Glu Ala Thr Val Ser Cys Ser 580  
 570 575 580  
 ggt tcc cat cac atc aag tgc tgc ata gag gag gat gga gac tac aaa 2068  
 Gly Ser His His Ile Lys Cys Cys Ile Glu Glu Asp Gly Asp Tyr Lys 595  
 585 590 595  
 gtt act ttc cat atg ggt tcc tca tcc ctt cct gct gca aaa gaa gtt 2116  
 Val Thr Phe His Met Gly Ser Ser Ser Leu Pro Ala Lys Glu Val 615  
 600 605 610 615  
 aac aaa aaa caa gty tgc tac aaa cac aat ttc aat gca agc tca gtt 2164  
 Asn Lys Lys Lys Gln Val Cys Tyr Lys His Asn Phe Asn Ala Ser Ser Val 630  
 620 625 630  
 tcc tgg tgt tca aaa act gtt gat gtg tgt tgt cac ttt acc aat gct 2212  
 Ser Trp Cys Ser Lys Thr Val Asp Val Cys Cys His Phe Thr Asn Ala 645  
 635 640 645  
 gct aat aat tca gtt tgg agc cca tct atg aag ctg aat ctg gtt cct 2260

Ala Asn Asn Ser Val Trp Ser Pro Ser Met Lys Leu Asn Leu Val Pro  
650 655 660

ggg gaa aac atc aca tgc cag gat ccc gta ata ggt gtc gga gag ccg  
Gly Glu Asn Ile Thr Cys Gln Asp Pro Val Ile Gly Val Gly Glu Pro  
665 670 675 2308

ggg aaa gtc atc cag aag cta tgc cgg ttc tca aac gtt ccc agc agc  
Gly Lys Val Ile Gln Lys Leu Cys Arg Phe Ser Asn Val Pro Ser Ser  
680 685 690 2356

ctt gag agt ccc att ggc ggg acc atc act tac aaa tgt gta ggc tcc  
Pro Glu Ser Pro Ile Gly Gly Thr Ile Thr Tyr Lys Cys Val Gly Ser  
700 705 710 2404

cgg tgg gag gag aag aga aat gac tgc atc tct gcc cca ata aac agt  
Gln Trp Glu Glu Lys Arg Asn Asp Cys Ile Ser Ala Pro Ile Asn Ser  
715 720 725 2452

ctg ctc cag atg gct aag gct ttg atc aag agc ccc tct cag gat gag  
Leu Leu Gln Met Ala Lys Ala Leu Ile Lys Ser Pro Ser Gln Asp Glu  
730 735 740 2500

atg ctc cct aca tac ctg aag gat ctt tct att agc ata gac aaa gcg  
Met Leu Pro Thr Tyr Leu Lys Asp Leu Ser Ile Ser Ile Asp Lys Ala  
745 750 755 2548

gaa cat gaa atc agc tct cct ggc agt ctg gga gcc att att aac  
Glu His Glu Ile Ser Ser Ser Pro Gly Ser Leu Gly Ala Ile Ile Asn  
760 765 770 2596

atc ctt gat ctg ctc tca aca gtt cca acc caa gta aat tca gaa atg  
Ile Leu Asp Leu Leu Ser Thr Val Pro Thr Gln Val Asn Ser Glu Met  
780 785 790 2644

atg acg cac gtg ctc tct acg gtt aat gtc atc ctt ggc aag ccc gtc  
Met Thr His Val Leu Ser Thr Val Asn Val Ile Leu Gly Lys Pro Val  
795 800 805 2692

ttg aac acc tgg aag gtt tta caa cag caa tgg acc aat cag agt tca  
Leu Asn Thr Trp Lys Val Leu Gln Gln Trp Thr Asn Gln Ser Ser  
810 815 820 2740

cag cta cta cat tca gtg gaa aga ttt tcc caa gca tta cag tca gga  
Gln Leu Leu His Ser Val Glu Arg Phe Ser Gln Ala Leu Gln Ser Gly  
825 830 835 2788

gat agc cct cct ttg tcc ttc cca act aat gtc cag atg agc agc  
Asp Ser Pro Pro Leu Ser Phe Ser Gln Thr Asn Val Gln Met Ser Ser  
840 845 850 2836

acg gta atc aag tcc agc cac cca gaa acc tat caa cag agg ttt gtt  
Thr Val Ile Lys Ser Ser His Pro Glu Thr Tyr Gln Gln Arg Phe Val  
860 865 870 2884

ttc cca tac ttt gac ctc tgg ggc aat gtg gtc att gac aag agc tac  
Phe Pro Tyr Phe Asp Leu Trp Gly Asn Val Val Ile Asp Lys Ser Tyr  
875 880 885 2932

cta gaa aac ttg cag tgg gat tgg tct att gtc acc atg gct ttc cca  
Leu Glu Asn Leu Gln Ser Asp Ser Ser Ile Val Thr Met Ala Phe Pro  
890 895 900 2980

act ctc caa gcc atc ctt gct cag gat atc cag gaa aat aac ttt gca  
Thr Leu Gln Ala Ile Leu Ala Gln Asp Ile Gln Glu Asn Asn Phe Ala  
905 910 915 3028

gag agc tta gtg atg aca acc act gtc agc cac aat acg act atg cca  
Glu Ser Leu Val Met Thr Thr Val Ser His Asn Thr Thr Met Pro  
920 925 930 3076

ttc agg att tca atg atc ttt aag aac aat agc cct tca ggc ggc gaa  
Phe Arg Ile Ser Met Thr Phe Lys Asn Asn Ser Pro Ser Gly Gly Glu  
940 945 950 3124

acg aag tgt gtc ttc tgg aac ttc agg ctt gcc aac aac aca ggg ggg  
Thr Lys Cys Val Phe Thr Asn Phe Arg Leu Ala Asn Asn Thr Gly Gly  
955 960 965 3172

tgg gac agc agt ggg tgc tat gtt gaa gaa ggt gat ggg gac aat gtc  
Trp Asp Ser Ser Gly Cys Tyr Val Glu Glu Gly Asp Gly Asp Asn Val  
970 975 980 3220

acc tgt atc tgt gac cac cta aca tca ttc tcc atc ctc atg tcc cct  
Thr Cys Ile Cys Asp His Leu Thr Ser Phe Ser Ile Leu Met Ser Pro  
985 990 995 3268

gac tcc cca gat cct agt tct ctc ctg gga ata ctc ctg gat att  
Asp Ser Pro Asp Pro Ser Ser Leu Leu Gly Ile Leu Leu Asp Ile  
1000 1005 1010 3313

att tct tat gtt ggg gtg ggc ttt tcc atc ttg agc ttg gca gcc  
Ile Ser Tyr Val Gly Val Gly Phe Ser Ile Leu Ser Leu Ala Ala  
1015 1020 1025 3358

tgt cta gtt gtg gaa gct gtg gtg tgg aaa tgg gtg acc aag aat  
Cys Leu Val Val Glu Ala Val Val Trp Lys Ser Val Thr Lys Asn  
1030 1035 1040 3403

cgg act tct tat atg cgc cac acc tgc ata gtg aat atc gct gcc  
Arg Thr Ser Tyr Met Arg His Thr Cys Ile Val Asn Ile Ala Ala  
1045 1050 1055 3448

tcc ctt ctg gtc gcc aac acc tgg ttc att gtg gtc gct gcc atc  
Ser Leu Leu Val Ala Asn Thr Trp Phe Ile Val Val Ala Ala Ile  
1060 1065 1070 3493

cag gac aat cgc tac ata ctc tgc aag aca gcc tgt gtg gct gcc  
Gln Asp Asn Arg Tyr Ile Leu Cys Lys Thr Ala Cys Val Ala Ala  
1075 1080 1085 3538

acc ttc ttc atc cac ttc ttc tac ctc agc gtc ttc ttc tgg atg  
Thr Phe Phe Ile His Phe Phe Tyr Leu Ser Val Phe Phe Trp Met  
1090 1095 1100 3583

ctg aca ctg ggc ctc atg ctg ttc tat tgc ctg gtt ttc att ctg  
Leu Thr Leu Gly Leu Met Leu Phe Tyr Arg Leu Val Phe Ile Leu  
1105 1110 1115 3628

cat gaa aca agc agg tcc act cag aaa gcc att gcc ttc tgt ctt  
His Glu Thr Ser Arg Ser Thr Gln Lys Ala Ile Ala Phe Cys Leu  
1120 1125 1130 3673

ggc tat ggc tgc cca ctt gcc atc tgg gtc atc acg ctg gga gcc  
Gly Tyr Gly Cys Pro Leu Ala Ile Ser Val Ile Thr Leu Gly Ala  
1135 1140 1145 3718

acc cag ccc cgg gaa gtc tat acg agg aag aat gtc tgt tgg ctc 3763  
 Thr Gln Pro Arg Glu Val Tyr Thr Arg Lys Asn Val Cys Trp Leu 1155 1160  
 aac tgg aag gac acc aag gcc ctg ctg gct ttc gcc atc cca gca 3808  
 Asn Trp Glu Asp Thr Lys Ala Leu Leu Ala Phe Ala Ile Pro Ala 1170 1175  
 ctg atc att gtg gtg gtg aac ata acc atc act att gtg gtc atc 3853  
 Leu Ile Ile Val Val Val Asn Ile Thr Ile Thr Ile Val Val Ile 1180 1190  
 acc aag atc ctg agg cct tcc att gga gac aag cca tgc aag cag 3898  
 Thr Lys Ile Leu Arg Pro Ser Ile Gly Asp Lys Pro Cys Lys Gln 1195 1200 1205  
 gag aag agc agc ctg ttt cag atc agc aag agc att ggg gtc ctc 3943  
 Glu Lys Ser Ser Leu Phe Gln Ile Ser Lys Ser Ile Gly Val Leu 1210 1215 1220  
 aca cca ctc ttg ggc ctc act tgg ggt ttt ggt ctc acc act gtg 3988  
 Thr Pro Leu Leu Gly Leu Thr Trp Gly Phe Gly Leu Thr Thr Val 1225 1230 1235  
 ttc cca ggg acc aac ctt gtg ttc cat atc ata ttt gcc atc ctc 4033  
 Phe Pro Gly Thr Asn Leu Val Phe His Ile Ile Phe Ala Ile Leu 1240 1245 1250  
 aat gtc ttc cag gga tta ttc att tta ctc ttt gga tgc ctc tgg 4078  
 Asn Val Phe Gln Gly Leu Phe Ile Leu Leu Phe Gly Cys Leu Trp 1255 1260 1265  
 gat ctg aag gta cag gaa gct ttg ctg aat aag ttt tca ttg tgg 4123  
 Asp Leu Lys Val Gln Glu Ala Leu Leu Asn Lys Phe Ser Leu Ser 1270 1275 1280  
 aga tgg tct tca cag cac tca aag tca aca tcc ctg ggt tca tcc 4168  
 Arg Trp Ser Ser Gln His Ser Lys Ser Thr Ser Leu Gly Ser Ser 1285 1290 1295  
 aca cct gtg ttt tct atg agt tct cca ata tca agg aga ttt aac 4213  
 Thr Pro Val Phe Ser Met Ser Ser Pro Ile Ser Arg Arg Phe Asn 1300 1305 1310  
 aat ttg ttt ggt aaa aca gga acg tat aat gtt tcc acc cca gaa 4258  
 Asn Leu Phe Gly Lys Thr Gly Thr Tyr Asn Val Ser Thr Pro Glu 1315 1320 1325  
 gca acc agc tca tcc ctg gaa aac tca tcc agt gct tct tgg ttg 4303  
 Ala Thr Ser Ser Ser Leu Glu Asn Ser Ser Ala Ser Ser Leu 1330 1335 1340  
 ctc aac taa gaacaggata atccaacta cgtgacctcc cggggacagt 4352  
 Leu Asn 1345  
 ggcgtgctt ttaaaagag atgcttgcaa agcaatgggg aacgtgttct cggggcaggt 4412  
 ttccggagc agatgccaaa aagactttt catagagaag aggttttctt ttgtaagac 4472  
 agaataaaaa taattgttat gttctgttt gtccctccc cctcccctt gtgtgatacc 4532

acatgtgtat agtatttaag tgaactcaa gccctcaagg cccaacttct ctgtctatat 4592  
 tgaataatag aatttogaag agacattttc actttttaca catggggcac aaagataaagc 4652  
 tttgatcaaa gtatgaagta aaaggttaac taggaataac ttcaagtgaat totaagaagg 4712  
 aaggaaaggaa gaaaggaagg aaagaaggga gggaaacagg gegaaggga aaaaagaa 4772  
 aaagagaag atgnaaatag gaacaataa agacaacaa cattaaagggc catattgtaa 4832  
 gatctcatg ttaatgatct aataataca ctcaagtcaa catgagaat tttttttaa 4892  
 tggctcaaaa atggaaaactg aaagcaagtc atgggggaatg aatactttgg gcagatctt 4952  
 cctcatgtct tottagttaa gagggaggaaa aaagggtga aaaaatagggg aggaatttc 5012  
 ttcatcgaa cgaacttcaag tggatacaaa tatttataag aaatgaatgg aaggaaatat 5072  
 gatctctcg agactaaact tgtatgttaa ggtttgaact aagtgaatgt atctgcagag 5132  
 gaagtattac aaagatatgt cattagatcc aagtctgat taaattttta tagtttatca 5192  
 gaaaagcctt atattttagt ttgtccaca ttgtgaagc aaaaaatata tatttgatat 5252  
 acccttcaat tgcctaaatt gatagtgc actgaagaca gccctgtca tatatttaat 5312  
 ggcctcaagc aggtacttct ctgtgcatta tagaatagat ttttaatac ttatagcatt 5372  
 gtattattatt atgtgtgtg tcaactgtat tattatgtg gatactggcc ctgtgtgtgt 5432  
 tgcatactc octatgtatt ctctgtttcc atctttaagt tcccagacca atatacatca 5492  
 agagtttgc atgtgtctaaa ttgtgtttat tccaaccag tggaaagctc ctgaaagaa 5552  
 attttacatt cgtgtgttct gtgtctctaa tgaacttga ccttgttgaa caaatggcag 5612  
 agccttccc aaggatttga ttgttttga attatctga tgtgtgcttt ttttgggtgt 5672  
 gtatttcatt aaaaaatata aatatttatg aaaaaaaaa aa 5714  
 <210> 20  
 <211> 1346  
 <212> PRT  
 <213> Homo sapiens  
 <400> 20  
 Met Lys Ser Pro Arg Arg Thr Thr Leu Cys Leu Met Phe Ile Val Ile 15  
 1 5 10  
 Tyr Ser Ser Lys Ala Ala Leu Asn Trp Asn Tyr Glu Ser Thr Ile His 30  
 20 25 30  
 Pro Leu Ser Leu His Glu His Glu Pro Ala Gly Glu Glu Ala Leu Arg 45  
 35 40  
 Gln Lys Arg Ala Val Ala Thr Lys Ser Pro Thr Ala Glu Glu Tyr Thr 60  
 50 55 60

Val Asn Ile Glu Ile Ser Phe Glu Asn Ala Ser Phe Leu Asp Pro Ile  
65 70 75 80

Lys Ala Tyr Leu Asn Ser Leu Ser Phe Pro Ile His Gly Asn Asn Thr  
85 90 95

Asp Gln Ile Thr Asp Ile Leu Ser Ile Asn Val Thr Thr Val Cys Arg  
100 105 110

Pro Ala Gly Asn Glu Ile Trp Cys Ser Cys Glu Thr Gly Tyr Gly Trp  
115 120 125

Pro Arg Glu Arg Cys Leu His Asn Leu Ile Cys Gln Glu Arg Asp Val  
130 135 140

Phe Leu Pro Gly His His Cys Ser Cys Leu Lys Glu Leu Pro Pro Asn  
145 150 155 160

Gly Pro Phe Cys Leu Leu Gln Glu Asp Val Thr Leu Asn Met Arg Val  
165 170 175

Arg Leu Asn Val Gly Phe Gln Glu Asp Leu Met Asn Thr Ser Ser Ala  
180 185 190

Leu Tyr Arg Ser Tyr Lys Thr Asp Leu Glu Thr Ala Phe Arg Lys Gly  
195 200 205

Tyr Gly Ile Leu Pro Gly Phe Lys Gly Val Thr Val Thr Gly Phe Lys  
210 215 220

Ser Gly Ser Val Val Thr Tyr Glu Val Lys Thr Thr Thr Pro Pro Ser  
225 230 235 240

Leu Glu Leu Ile His Lys Ala Asn Glu Gln Val Val Gln Ser Leu Asn  
245 250 255

Gln Thr Tyr Lys Met Asp Tyr Asn Ser Phe Gln Ala Val Thr Ile Asn  
260 265 270

Glu Ser Asn Phe Phe Val Thr Pro Glu Ile Ile Phe Glu Gly Asp Thr  
275 280 285

Val Ser Leu Val Cys Glu Lys Glu Val Leu Ser Ser Asn Val Ser Trp  
290 295 300

Arg Tyr Glu Glu Gln Gln Leu Glu Ile Gln Asn Ser Ser Arg Phe Ser  
305 310 315 320

Ile Tyr Thr Ala Leu Phe Asn Asn Met Thr Ser Val Ser Lys Leu Thr  
325 330 335

Ile His Asn Ile Thr Pro Gly Asp Ala Gly Glu Tyr Val Cys Lys Leu  
340 345 350

Ile Leu Asp Ile Phe Glu Tyr Glu Cys Lys Lys Ile Asp Val Met  
355 360 365

Pro Ile Gln Ile Leu Ala Asn Glu Glu Met Lys Val Met Cys Asp Asn  
370 375 380

Asn Pro Val Ser Leu Asn Cys Cys Ser Gln Gly Asn Val Asn Trp Ser  
385 390 395 400

Lys Val Glu Trp Lys Gln Glu Gly Lys Ile Asn Ile Pro Gly Thr Pro  
405 410 415

Glu Thr Asp Ile Asp Ser Ser Cys Ser Arg Tyr Thr Leu Lys Ala Asp  
420 425 430

Gly Thr Gln Cys Pro Ser Gly Ser Ser Gly Thr Thr Val Ile Tyr Thr  
435 440 445

Cys Glu Phe Ile Ser Ala Tyr Gly Ala Arg Gly Ser Ala Asn Ile Lys  
450 455 460

Val Thr Phe Ile Ser Val Ala Asn Leu Thr Ile Thr Pro Asp Pro Ile  
465 470 475 480

Ser Val Ser Glu Gly Gln Asn Phe Ser Ile Lys Cys Ile Ser Asp Val  
485 490 495

Ser Asn Tyr Asp Glu Val Tyr Trp Asn Thr Ser Ala Gly Ile Lys Ile  
500 505 510

Tyr Gln Arg Phe Tyr Thr Thr Arg Arg Tyr Leu Asp Gly Ala Glu Ser  
515 520 525

Val Leu Thr Val Lys Thr Ser Thr Arg Glu Trp Asn Gly Thr Tyr His  
530 535 540

Cys Ile Phe Arg Tyr Lys Asn Ser Tyr Ser Ile Ala Thr Lys Asp Val  
545 550 555 560

Ile Val His Pro Leu Pro Leu Lys Leu Asn Ile Met Val Asp Pro Leu  
35

565 570 575

Glu Ala Thr Val Ser Cys Ser Gly Ser His His Ile Lys Cys Cys Ile  
580 585 590

Glu Glu Asp Gly Asp Tyr Lys Val Thr Phe His Met Gly Ser Ser  
595 600 605

Leu Pro Ala Ala Lys Glu Val Asn Lys Lys Glu Val Cys Tyr Lys His  
610 615 620

Asn Phe Asn Ala Ser Ser Val Ser Trp Cys Ser Lys Thr Val Asp Val  
635 630 635 640

Cys Cys His Phe Thr Asn Ala Ala Asn Asn Ser Val Trp Ser Pro Ser  
645 650 655

Met Lys Leu Asn Leu Val Pro Gly Glu Asn Ile Thr Cys Glu Asp Pro  
660 665 670

Val Ile Gly Val Gly Glu Pro Gly Lys Val Ile Gln Lys Leu Cys Arg  
675 680 685

Phe Ser Asn Val Pro Ser Ser Pro Glu Ser Pro Ile Gly Thr Ile  
690 695 700

Thr Tyr Lys Cys Val Gly Ser Gln Trp Glu Lys Arg Asn Asp Cys  
705 710 715 720

Ile Ser Ala Pro Ile Asn Ser Leu Leu Gln Met Ala Lys Ala Leu Ile  
725 730 735

Lys Ser Pro Ser Gln Asp Glu Met Leu Pro Thr Tyr Leu Lys Asp Leu  
740 745 750

Ser Ile Ser Ile Asp Lys Ala Glu His Glu Ile Ser Ser Ser Pro Gly  
755 760 765

Ser Leu Gly Ala Ile Ile Asn Ile Leu Asp Leu Ser Thr Val Pro  
770 775 780

Thr Gln Val Asn Ser Glu Met Met Thr His Val Leu Ser Thr Val Asn  
785 790 795 800

Val Ile Leu Gly Lys Pro Val Leu Asn Thr Trp Lys Val Leu Gln Gln  
805 810 815

Gln Trp Thr Asn Gln Ser Ser Gln Leu Leu His Ser Val Glu Arg Phe  
820 825 830

Ser Gln Ala Leu Gln Ser Gly Asp Ser Pro Pro Leu Ser Phe Ser Gln  
835 840 845

Thr Asn Val Gln Met Ser Ser Thr Val Ile Lys Ser Ser His Pro Glu  
850 855 860

Thr Tyr Gln Gln Arg Phe Val Phe Pro Tyr Phe Asp Leu Trp Gly Asn  
865 870 875 880

Val Val Ile Asp Lys Ser Tyr Leu Glu Asn Leu Gln Ser Asp Ser Ser  
885 890 895

Ile Val Thr Met Ala Phe Pro Thr Leu Gln Ala Ile Leu Ala Gln Asp  
900 905 910

Ile Gln Glu Asn Asn Phe Ala Glu Ser Leu Val Met Thr Thr Thr Val  
915 920 925

Ser His Asn Thr Thr Met Pro Phe Arg Ile Ser Met Thr Phe Lys Asn  
930 935 940

Asn Ser Pro Ser Gly Gly Glu Thr Lys Cys Val Phe Trp Asn Phe Arg  
945 950 955 960

Leu Ala Asn Asn Thr Gly Gly Trp Asp Ser Ser Gly Cys Tyr Val Glu  
965 970 975

Glu Gly Asp Gly Asp Asn Val Thr Cys Ile Cys Asp His Leu Thr Ser  
980 985 990

Phe Ser Ile Leu Met Ser Ser Pro Asp Ser Pro Asp Pro Ser Ser Leu Leu  
995 1000 1005

Gly Ile Leu Leu Asp Ile Ile Ser Tyr Val Gly Val Gly Phe Ser  
1010 1015 1020

Ile Leu Ser Leu Ala Ala Cys Leu Val Val Glu Ala Val Val Trp  
1025 1030 1035

Lys Ser Val Thr Lys Asn Arg Thr Ser Tyr Met Arg His Thr Cys  
1040 1045 1050

Ile Val Asn Ile Ala Ala Ser Leu Leu Val Ala Asn Thr Trp Phe  
1055 1060 1065

Ile Val Val Ala Ala Ile Gln Asp Asn Arg Tyr Ile Leu Cys Lys  
1070 1075 1080

Thr Ala Cys Val Ala Ala Thr Phe Phe Ile His Phe Phe Tyr Leu  
1085 1090 1095

Ser Val Phe Phe Trp Met Leu Thr Leu Gly Leu Met Leu Phe Tyr  
1100 1105 1110

Arg Leu Val Phe Ile Leu His Gln Thr Ser Arg Ser Thr Gln Lys  
1115 1120 1125

Ala Ile Ala Phe Cys Leu Gly Tyr Gly Cys Pro Leu Ala Ile Ser  
1130 1135 1140

Val Ile Thr Leu Gly Ala Thr Gln Pro Arg Gln Val Tyr Thr Arg  
1145 1150 1155

Lys Asn Val Cys Trp Leu Asn Trp Glu Asp Thr Lys Ala Leu Leu  
1160 1165 1170

Ala Phe Ala Ile Pro Ala Leu Ile Ile Val Val Asn Ile Thr  
1175 1180 1185

Ile Thr Ile Val Ile Thr Lys Ile Leu Arg Pro Ser Ile Gly  
1190 1195 1200

Asp Lys Pro Cys Lys Gln Glu Lys Ser Ser Leu Phe Gln Ile Ser  
1205 1210 1215

Lys Ser Ile Gly Val Leu Thr Pro Leu Leu Gly Leu Thr Trp Gly  
1220 1225 1230

Phe Gly Leu Thr Thr Val Phe Pro Gly Thr Asn Leu Val Phe His  
1235 1240 1245

Ile Ile Phe Ala Ile Leu Asn Val Phe Gln Gly Leu Phe Ile Leu  
1250 1255 1260

Leu Phe Gly Cys Leu Trp Asp Leu Lys Val Gln Glu Ala Leu Leu  
1265 1270 1275

Asn Lys Phe Ser Leu Ser Arg Trp Ser Ser Gln His Ser Lys Ser  
1280 1285 1290

Thr Ser Leu Gly Ser Ser Thr Pro Val Phe Ser Met Ser Ser Pro  
1295 1300 1305

Ile Ser Arg Arg Phe Asn Asn Leu Phe Gly Lys Thr Gly Thr Tyr  
1310 1315 1320

Asn Val Ser Thr Pro Glu Ala Thr Ser Ser Ser Leu Glu Asn Ser  
1325 1330 1335

Ser Ser Ala Ser Ser Leu Leu Asn  
1340 1345

<210> 21

<211> 4041

<212> DNA

<213> Homo sapiens

<400> 21

atgaatccc caaggagaac caattttg ctcattgtta ttgtgattta ttcttccaaa 60

gttgcaatga actgggaatta cgagcttact attcatcctt tgagctttca tgaacatgaa 120

ccagctggty aaggggcaact gagggcaaaa cagagccgttg ccacaaaaag tcttaccgct 180

gaagaaatca ctgttaatat tgagatcagt ttgaaaaatg catcttctct ggatctctatc 240

aaagctact tgaacagcct cagttttcca attcatggga ataacactga ccaaatattact 300

gacattttga gcataaatgt gacaacagtc tgcagacctg ctggaaatga aatctggtgc 360

tctgtcgaga caggttatgg gtggcctcgg gaaggtgtgc ttcaaatct catittgtcaa 420

gagctgaag tcttctccc agggcaccat tgcagttgct ttaagaact gcttcccaat 480

ggactttttt gcctgcttca ggaagatgtt accctgaaca tgagagctcag actaaatgta 540

ggctttcaag aagacctcat gaacacttc tccgcctct ataggttcta caagaccgac 600

ttggaaacag cgttcctgaa ggtttacgga attttaccag gcttcaagggt cgtgactgtg 660

acagggttca agtctggaag tgtgtgtgtg acatatgaag tcaagactac accaccatca 720

cttgagttaa tacataaagc caatgaacaa gtgtacaga gcttcaatca gacctacaaa 780

atggactaca actcctttca agcagttact atcaatgaaa gcaattttct tgcacacca 840

gaaatcatct ttgaagggtga caccagtcagt ctgggtgtgt aaaaaggagt ttgtctctcc 900

aatgtctctt ggcgttatga agaacagcag ttggaaatcc agaacagcag cagattctcg 960

atttacacgg cacitttcaa caacatgact tgggtgtcca agctcaccat ccacaacatc 1020

actccaggtg atgcaggtga atattgttc aaactgatat tagacatttt tgaatatgag 1080

tgcaagaga aaatagatgt tatgccatc caaattttgg caaatgaga aatgaagggtg 1140

atgtgcgaca acaatcctgt atctttgaac tgcgtcagtc aggttaatgt taattggagc 1200

aaagtgaat ggaagcagga aggaanaata aatattccag gaacctctga gacagacata 1260

gattctagct gcagcagata caccctcaag gctgatggaa cccagtgccc aagcgggtcg 1320



tctggacaa cagtcatacta cacttgtag ttcatactg cctatggagc cagaggcagt 1380  
 gcaaacataa aagtgacatt catctctg gccaactaa caataacccc ggaaccaatt 1440  
 tctgttctg egggaacaaa cttttcata aaatgcatca gtgatgtgag taactatgat 1500  
 gaggtttatt ggaacacttc tgcctgaatt aaatatacc aaagatitita taccacaggg 1560  
 aggtatctg atggagcaga atcagtagt acagtaaga cctcgaccag ggaagtgaat 1620  
 ggaacctatc actgcatatt tagatataag aatcatata gtaattcaac caaagacgtc 1680  
 atgttcaacc cgtcgtctct aaagtgaac atcatgggtg atcctttgga agctactggt 1740  
 tcatgcaatg gttccatca catcaatg tgcataaggg aggatggaga ctacaagtt 1800  
 actttccata tgggttctct atccctctct gctgcaaaag aagttaacaa aaaaacagtg 1860  
 tgcataaac acaatttcaa tgaagctca gtttctggt gttcaaaac tgttgatgtg 1920  
 tgtgttact ttacaatgc tgcataaat tcaagttgga gccatctat gaagctgaat 1980  
 ctggttctct gggaaacat cacatgccag gatccgtaa taggtgtcgg agagcgggg 2040  
 aaagtcac agagctatg ccggttctca aacgttccca gcagcctga gagtccatt 2100  
 ggcgggaacca tcaattacaa atgtgtaggc tccagtggg tccagtgagg aatgactgc 2160  
 atctctgcc caataaacag tctgtctcag atggctaagg ctttgatcaa gagccctct 2220  
 caggatgaga tgcctctac atactgaag gatcttcta ttagcataga caaagoggaa 2280  
 catgaaatca gctctctcc tgggagctg ggaagcattta ttaactctc tgaatcgtc 2340  
 tcaacagtte caaccagtt aaattcagaa atgatgacgc acgtgctctc tacygttaat 2400  
 gtcactctg gcaagccctt ctggaacccc ctggaagttt tacaacagca atggaccaat 2460  
 cagagttcac agtactata ttcagtgaaa agattttccc aagcattaca gtcaggagat 2520  
 agccctctt tgcctctct ccaactaat gtgcagatga gcagacggt aatcaagtc 2580  
 agccaccag aaactatca acagagttt gtttcccat acttgacct ctggggcaat 2640  
 gtggtcatg acaagagcta cctagaanaa ttgcagtcgg atcgtctat tgtcaacctg 2700  
 gctttcccaa ctctcaagc catcttctc caggatatcc aggaataaa ctttgacagg 2760  
 agcttagtga tgacaaccac tgcagccac aatacgaacta tgcattcag gatttcaatg 2820  
 acttttaaga acaatagccc ttcagggcg gaaacgaagt gtgtctctg gaacttcagg 2880  
 ctgccaaca acacaggggg gtgggacagc agtgggtgct atgttgaaga agtgatggg 2940  
 gacaatgta cctgtatctg tgaccaccta acatcattct ccatctcat gtccctgac 3000  
 tccccagatc ctagtctct cctgggaata ctctggata ttattctta tgttgggtg 3060  
 ggtctttcca tcttgagctt ggcagcctgt ctagtgtggt aagctgtgt gtggaaatcg 3120  
 gtgaccaaga atcggacttc ttatgagc cacactgca tagtgaatat cgtgctctc 3180

cttctggtg ccaacactg gttcattgtg gtgcgtgca tccaggacaa togtacata 3240  
 cttctgaaga cagcctgtgt ggcgcacac ttctctatcc acttcttcta cctcagcgtc 3300  
 ttctcttga tgcagactt gggcctcatg ctgtctatc gccctggtttt cactctgcat 3360  
 gaaacaaga ggtccactca gaagccatt gccctctgtc ttggctatgg ctgcccactt 3420  
 gccatctcgg tcatcagctt gggagccacc cagcccgagg aagctctatc gaggaagant 3480  
 gtctgtggc tcaactggga ggaacacaa ggcctgtgtg ctttcgcat cccagcactg 3540  
 atcattgtgg tggtaacat aaccatcact atgtgtgtca tcaaccaagat cctgagcgct 3600  
 tccattggag acaagcccatg caagcaggag aagacagcc tgtttcagat cagcaagagc 3660  
 attgggtcc tcaacactt cttggcctc acttgggggt ttggtctcac cactgtgttc 3720  
 ccagggaaca acctgtgtt ccatatcata ttggcatcc tcaatgtctt ccaggggatta 3780  
 ttcattttac tcttggatg cctctggat ctgaaggtac aggaagcttt gctgaataag 3840  
 ttttcattgt cgaatgggtc ttcacagcac tcaaaagtc aaatcctctgg ttcacaca 3900  
 cctgtgttt ctatgagttc tcaatatca aggagattta acaatttgtt tggtaaaaca 3960  
 ggaagcata atgtttccac cccagaagca accagctcat ccttgaaaaa ctcactcagt 4020  
 gcttctctgt tgcataacta a

<210> 22  
 <211> 265  
 <212> PRI  
 <213> Homo sapiens  
 <400> 22

Gly Ile Leu Leu Asp Ile Ile Ser Tyr Val Gly Val Gly Phe Ser Ile  
 1 5 10 15  
 Leu Ser Leu Ala Ala Cys Leu Val Val Glu Ala Val Val Trp Lys Ser  
 20 25 30  
 Val Thr Lys Asn Arg Thr Ser Tyr Met Arg His Thr Cys Ile Val Asn  
 35 40 45  
 Ile Ala Ala Ser Leu Leu Val Ala Asn Thr Trp Phe Ile Val Val Ala  
 50 55 60  
 Ala Ile Gln Asp Asn Arg Tyr Ile Leu Cys Lys Thr Ala Cys Val Ala  
 65 70 75 80  
 Ala Thr Phe Phe Ile Ile His Phe Phe Tyr Leu Ser Val Phe Phe Trp Met  
 85 90 95  
 Leu Thr Leu Gly Leu Met Leu Phe Phe Tyr Arg Leu Val Phe Ile Leu His  
 100 105 110  
 Glu Thr Ser Arg Ser Thr Gln Lys Ala Ile Ala Phe Cys Leu Gly Tyr  
 115 120 125  
 Gly Cys Pro Leu Ala Ile Ser Val Ile Thr Leu Gly Ala Thr Gln Pro  
 41

Ile Trp Cys Ser Cys Glu Thr Gly Tyr Gly Trp Pro Arg Glu Arg Cys  
100 105 110  
Leu His Asn Leu Ile Cys Gln Glu Arg Asp Val Phe Leu Pro Gly His  
115 120 125  
His Cys Ser Cys Leu Lys Glu Leu Pro Pro Asn Gly Pro Phe Cys Leu  
130 135 140  
Leu Gln Glu Asp Val Thr Leu Asn Met Arg Val Arg Leu Asn Val Gly  
145 150 155 160  
Phe Gln Glu Asp Leu Met Asn Thr Ser Ser Ala Leu Tyr Arg Ser Tyr  
165 170 175  
Lys Thr Asp Leu Glu Thr Ala Phe Arg Lys Gly Tyr Gly Ile Leu Pro  
180 185 190  
Gly Phe Lys Gly Val Thr Val Thr Gly Phe Lys Ser Gly Ser Val Val  
195 200 205  
Val Thr Tyr Glu Val Lys Thr Thr Pro Pro Ser Leu Glu Leu Ile His  
210 215 220  
Lys Ala Asn Glu Gln Val Val Gln Ser Leu Asn Gln Thr Tyr Lys Met  
225 230 235 240  
Asp Tyr Asn Ser Phe Gln Ala Val Thr Ile Asn Glu Ser Asn Phe Phe  
245 250 255  
Val Thr Pro Glu Ile Ile Phe Glu Gly Asp Thr Val Ser Leu Val Cys  
260 265 270  
Glu Lys Glu Val Leu Ser Ser Asn Val Ser Trp Arg Tyr Glu Glu Gln  
275 280 285  
Gln Leu Glu Ile Gln Asn Ser Ser Arg Phe Ser Ile Tyr Thr Ala Leu  
290 295 300  
Phe Asn Asn Met Thr Ser Val Ser Lys Leu Thr Ile His Asn Ile Thr  
305 310 315 320  
Pro Gly Asp Ala Gly Glu Tyr Val Cys Lys Leu Ile Leu Asp Ile Phe  
325 330 335  
Glu Tyr Glu Cys Lys Lys Lys Ile Asp Val Met Pro Ile Gln Ile Leu  
340 345 350  
Ala Asn Glu Glu Met Lys Val Met Cys Asp Asn Asn Pro Val Ser Leu  
355 360 365  
Asn Cys Cys Ser Gln Gly Asn Val Asn Trp Ser Lys Val Glu Trp Lys  
370 375 380  
Gln Glu Gly Lys Ile Asn Ile Pro Gly Thr Pro Glu Thr Asp Ile Asp  
385 390 395 400  
Ser Ser Cys Ser Arg Tyr Thr Leu Lys Ala Asp Gly Thr Gln Cys Pro  
405 410 415  
Ser Gly Ser Ser Gly Thr Thr Val Ile Tyr Thr Cys Glu Phe Ile Ser  
420 425 430

Arg Glu Val Tyr Thr Arg Lys Asn Val Cys Trp Leu Asn Trp Glu Asp  
145 150 155 160  
Thr Lys Ala Leu Leu Ala Phe Ala Ile Pro Ala Leu Ile Ile Val Val  
165 170 175  
Val Asn Ile Thr Ile Thr Ile Val Val Ile Thr Lys Ile Leu Arg Pro  
180 185 190  
Ser Ile Gly Asp Lys Pro Cys Lys Gln Glu Lys Ser Ser Leu Phe Gln  
195 200 205  
Ile Ser Lys Ser Ile Gly Val Leu Thr Pro Leu Leu Gly Leu Thr Trp  
210 215 220  
Gly Phe Gly Leu Thr Thr Val Phe Pro Gly Thr Asn Leu Val Phe His  
225 230 235 240  
Ile Ile Phe Ala Ile Leu Asn Val Phe Gln Gly Leu Phe Ile Leu Leu  
245 250 255  
Phe Gly Cys Leu Trp Asp Leu Lys Val  
260 265  
<210> 23  
<211> 21  
<212> PRT  
<213> Homo sapiens  
<400> 23  
Met Lys Ser Pro Arg Arg Thr Thr Leu Cys Leu Met Phe Ile Val Ile  
1 5 10 15  
Tyr Ser Ser Lys Ala  
20  
<210> 24  
<211> 1325  
<212> PRT  
<213> Homo sapiens  
<400> 24  
Ala Leu Asn Trp Asn Tyr Glu Ser Thr Ile His Pro Leu Ser Leu His  
1 5 10 15  
Glu His Glu Pro Ala Gly Glu Glu Ala Leu Arg Gln Lys Arg Ala Val  
20 25 30  
Ala Thr Lys Ser Pro Thr Ala Glu Glu Tyr Thr Val Asn Ile Glu Ile  
35 40 45  
Ser Phe Glu Asn Ala Ser Phe Leu Asp Pro Ile Lys Ala Tyr Leu Asn  
50 55 60  
Ser Leu Ser Phe Pro Ile His Gly Asn Asn Thr Asp Gln Ile Thr Asp  
65 70 75 80  
Ile Leu Ser Ile Asn Val Thr Thr Val Cys Arg Pro Ala Gly Asn Glu  
85 90 95

Ala Tyr Gly Ala Arg Gly Ser Ala Asn Ile Lys Val Thr Phe Ile Ser  
435 440 445

Val Ala Asn Leu Thr Ile Thr Pro Asp Pro Ile Ser Val Ser Glu Gly  
450 455 460

Gln Asn Phe Ser Ile Lys Cys Ile Ser Asp Val Ser Asn Tyr Asp Glu  
465 470 475 480

Val Tyr Trp Asn Thr Ser Ala Gly Ile Lys Ile Tyr Gln Arg Phe Tyr  
495 490 495

Thr Thr Arg Arg Tyr Leu Asp Gly Ala Glu Ser Val Leu Thr Val Lys  
500 505 510

Thr Ser Thr Arg Glu Trp Asn Gly Thr Tyr His Cys Ile Phe Arg Tyr  
515 520 525

Lys Asn Ser Tyr Ser Ile Ala Thr Lys Asp Val Ile Val His Pro Leu  
530 535 540

Pro Leu Lys Leu Asn Ile Met Val Asp Pro Leu Glu Ala Thr Val Ser  
545 550 555 560

Cys Ser Gly Ser His His Ile Lys Cys Cys Ile Glu Glu Asp Gly Asp  
565 570 575

Tyr Lys Val Thr Phe His Met Gly Ser Ser Ser Leu Pro Ala Ala Lys  
580 585 590

Glu Val Asn Lys Lys Gln Val Cys Tyr Lys His Asn Phe Asn Ala Ser  
595 600 605

Ser Val Ser Trp Cys Ser Lys Thr Val Asp Val Cys Cys His Phe Thr  
610 615 620

Asn Ala Ala Asn Ser Val Trp Ser Pro Ser Met Lys Leu Asn Leu  
625 630 635 640

Val Pro Gly Glu Asn Ile Thr Cys Gln Asp Pro Val Ile Gly Val Gly  
645 650 655

Glu Pro Gly Lys Val Ile Gln Lys Leu Cys Arg Phe Ser Asn Val Pro  
660 665 670

Ser Ser Pro Glu Ser Pro Ile Gly Gly Thr Ile Thr Tyr Lys Cys Val  
675 680 685

Gly Ser Gln Trp Glu Glu Lys Arg Asn Asp Cys Ile Ser Ala Pro Ile  
690 695 700

Asn Ser Leu Leu Gln Met Ala Lys Ala Leu Ile Lys Ser Pro Ser Gln  
705 710 715 720

Asp Glu Met Leu Pro Thr Tyr Leu Lys Asp Leu Ser Ile Ser Ile Asp  
725 730 735

Lys Ala Glu His Glu Ile Ser Ser Pro Gly Ser Leu Gly Ala Ile  
740 745 750

Ile Asn Ile Leu Asp Leu Leu Ser Thr Val Pro Thr Gln Val Asn Ser  
755 760 765

Glu Met Met Thr His Val Leu Ser Thr Val Asn Val Ile Leu Gly Lys  
770 775 780

Pro Val Leu Asn Thr Trp Lys Val Leu Gln Gln Trp Thr Asn Gln  
785 790 795 800

Ser Ser Gln Leu Leu His Ser Val Glu Arg Phe Ser Gln Ala Leu Gln  
805 810 815

Ser Gly Asp Ser Pro Pro Leu Ser Phe Ser Gln Thr Asn Val Gln Met  
820 825 830

Ser Ser Thr Val Ile Lys Ser Ser His Pro Glu Thr Tyr Gln Gln Arg  
835 840 845

Phe Val Phe Pro Tyr Phe Asp Leu Trp Gly Asn Val Val Ile Asp Lys  
850 855 860

Ser Tyr Leu Glu Asn Leu Gln Ser Asp Ser Ser Ile Val Thr Met Ala  
865 870 875 880

Phe Pro Thr Leu Gln Ala Ile Leu Ala Gln Asp Ile Gln Glu Asn Asn  
885 890 895

Phe Ala Glu Ser Leu Val Met Thr Thr Val Ser His Asn Thr Thr  
900 905 910

Met Pro Phe Arg Ile Ser Met Thr Phe Lys Asn Asn Ser Pro Ser Gly  
915 920 925

Gly Glu Thr Lys Cys Val Phe Trp Asn Phe Arg Leu Ala Asn Asn Thr  
930 935 940

Gly Gly Trp Asp Ser Ser Gly Cys Tyr Val Glu Glu Gly Asp Gly Asp  
945 950 955 960

Asn Val Thr Cys Ile Cys Asp His Leu Thr Ser Phe Ser Ile Leu Met  
965 970 975

Ser Pro Asp Ser Pro Asp Pro Ser Ser Leu Leu Gly Ile Leu Leu Asp  
980 985 990

Ile Ile Ser Tyr Val Gly Val Gly Phe Ser Ile Leu Ser Leu Ala Ala  
995 1000 1005

Cys Leu Val Val Glu Ala Val Val Trp Lys Ser Val Thr Lys Asn  
1010 1015 1020

Arg Thr Ser Tyr Met Arg His Thr Cys Ile Val Asn Ile Ala Ala  
1025 1030 1035

Ser Leu Leu Val Ala Asn Thr Trp Phe Ile Val Val Ala Ala Ile  
1040 1045 1050

Gln Asp Asn Arg Tyr Ile Leu Cys Lys Thr Ala Cys Val Ala Ala  
1055 1060 1065

Thr Phe Phe Ile His Phe Phe Tyr Leu Ser Val Phe Phe Trp Met  
1070 1075 1080

Leu Thr Leu Gly Leu Met Leu Phe Tyr Arg Leu Val Phe Ile Leu  
1085 1090 1095

His Glu Thr Ser Arg Ser Thr Gln Lys Ala Ile Ala Phe Cys Leu 1100 1105 1110  
 Gly Tyr Gly Cys Pro Leu Ala Ile Ser Val Ile Thr Leu Gly Ala 1115 1120 1125  
 Thr Gln Pro Arg Glu Val Tyr Thr Arg Lys Asn Val Cys Trp Leu 1130 1135 1140  
 Asn Trp Glu Asp Thr Lys Ala Leu Leu Ala Phe Ala Ile Pro Ala 1145 1150 1155  
 Leu Ile Ile Val Val Val Asn Ile Thr Ile Thr Ile Val Val Ile 1160 1165 1170  
 Thr Lys Ile Leu Arg Pro Ser Ile Gly Asp Lys Pro Cys Lys Gln 1175 1180 1185  
 Glu Lys Ser Ser Leu Phe Gln Ile Ser Lys Ser Ile Gly Val Leu 1190 1195 1200  
 Thr Pro Leu Leu Gly Leu Thr Trp Gly Phe Gly Leu Thr Thr Val 1205 1210 1215  
 Phe Pro Gly Thr Asn Leu Val Phe His Ile Ile Phe Ala Ile Leu 1220 1225 1230  
 Asn Val Phe Gln Gly Leu Phe Ile Leu Leu Phe Gly Cys Leu Trp 1235 1240 1245  
 Asp Leu Lys Val Gln Glu Ala Leu Leu Asn Lys Phe Ser Leu Ser 1250 1255 1260  
 Arg Trp Ser Ser Gln His Ser Lys Ser Thr Ser Leu Gly Ser Ser 1265 1270 1275  
 Thr Pro Val Phe Ser Met Ser Ser Pro Ile Ser Arg Arg Phe Asn 1280 1285 1290  
 Asn Leu Phe Gly Lys Thr Gly Thr Tyr Asn Val Ser Thr Pro Glu 1295 1300 1305  
 Ala Thr Ser Ser Ser Leu Glu Asn Ser Ser Ser Ala Ser Ser Leu 1310 1315 1320  
 Leu Asn 1325  
 <210> 25  
 <211> 1371  
 <212> PRT  
 <213> Homo sapiens  
 <400> 25  
 Ala Ser Asp Gln Ser Gly Ser Gln Pro Gly Asp His Ser Ala Gly Gln 1 5 10 15  
 Ala Asn Gln Leu Lys Leu Glu Asp Met Lys Ser Pro Arg Arg Thr Thr 20 25 30  
 Leu Cys Leu Met Phe Ile Val Ile Tyr Ser Ser Lys Ala Ala Leu Asn 46

Trp Asn Tyr Glu Ser Thr Ile His Pro Leu Ser Leu His Glu His Glu 50 55 60  
 Pro Ala Gly Glu Glu Ala Leu Arg Gln Lys Arg Ala Val Ala Thr Lys 65 70 75 80  
 Ser Pro Thr Ala Glu Glu Tyr Thr Val Asn Ile Glu Ile Ser Phe Glu 85 90 95  
 Asn Ala Ser Phe Leu Asp Pro Ile Lys Ala Tyr Leu Asn Ser Leu Ser 100 105 110  
 Phe Pro Ile His Gly Asn Asn Thr Asp Gln Ile Thr Asp Ile Leu Ser 115 120 125  
 Ile Asn Val Thr Thr Val Cys Arg Pro Ala Gly Asn Glu Ile Trp Cys 130 135 140  
 Ser Cys Glu Thr Gly Tyr Gly Trp Pro Arg Glu Arg Cys Leu His Asn 145 150 155 160  
 Leu Ile Cys Gln Glu Arg Asp Val Phe Leu Pro Gly His His Cys Ser 165 170 175  
 Cys Leu Lys Glu Leu Pro Pro Asn Gly Pro Phe Cys Leu Leu Gln Glu 180 185 190  
 Asp Val Thr Leu Asn Met Arg Val Arg Leu Asn Val Gly Phe Gln Glu 195 200 205  
 Asp Leu Met Asn Thr Ser Ser Ala Leu Tyr Arg Ser Tyr Lys Thr Asp 210 215 220  
 Leu Glu Thr Ala Phe Arg Lys Gly Tyr Gly Ile Leu Pro Gly Phe Lys 225 230 235 240  
 Gly Val Thr Val Thr Gly Phe Lys Ser Gly Ser Val Val Val Thr Tyr 245 250 255  
 Glu Val Lys Thr Thr Pro Pro Ser Leu Glu Leu Ile His Lys Ala Asn 260 265 270  
 Glu Gln Val Val Gln Ser Leu Asn Gln Thr Tyr Lys Met Asp Tyr Asn 275 280 285  
 Ser Phe Gln Ala Val Thr Ile Asn Glu Ser Asn Phe Phe Val Thr Pro 290 295 300  
 Glu Ile Ile Phe Glu Gly Asp Thr Val Ser Leu Val Cys Glu Lys Glu 305 310 315 320  
 Val Leu Ser Ser Asn Val Ser Trp Arg Tyr Glu Glu Gln Gln Leu Glu 325 330 335  
 Ile Gln Asn Ser Ser Arg Phe Ser Ile Tyr Thr Ala Leu Phe Asn Asn 340 345 350  
 Met Thr Ser Val Ser Lys Leu Thr Ile His Asn Ile Thr Pro Gly Asp 355 360 365  
 Ala Gly Glu Tyr Val Cys Lys Lys Leu Ile Leu Asp Ile Phe Glu Tyr Glu 47

WO 01/53451

375

380

PCT/US00/34983

Cys Lys Lys Lys Ile Asp Val Met Pro Ile Gln Ile Leu Ala Asn Glu  
385 390 395 400

Glu Met Lys Val Met Cys Asp Asn Asn Pro Val Ser Leu Asn Cys Cys  
405 410 415 420 425 430

Ser Gln Gly Asn Val Asn Trp Ser Lys Val Glu Trp Lys Gln Glu Gly  
420 425 430 435 440 445

Lys Ile Asn Ile Pro Gly Thr Pro Glu Thr Asp Ile Asp Ser Ser Cys  
435 440 445 450 455 460

Ser Arg Tyr Thr Leu Lys Ala Asp Gly Thr Gln Cys Pro Ser Gly Ser  
450 455 460 465 470 475 480

Ser Gly Thr Thr Val Ile Tyr Thr Cys Glu Phe Ile Ser Ala Tyr Gly  
465 470 475 480 485 490 495

Ala Arg Gly Ser Ala Asn Ile Lys Val Thr Phe Ile Ser Val Ala Asn  
485 490 495 500 505 510 515

Leu Thr Ile Thr Pro Asp Pro Ile Ser Val Ser Glu Gly Gln Asn Phe  
500 505 510 515 520 525 530

Ser Ile Lys Cys Ile Ser Asp Val Ser Asn Tyr Asp Glu Val Tyr Trp  
515 520 525 530 535 540 545

Asn Thr Ser Ala Gly Ile Lys Ile Tyr Gln Arg Phe Tyr Thr Thr Arg  
530 535 540 545 550 555 560

Arg Tyr Leu Asp Gly Ala Glu Ser Val Leu Thr Val Lys Thr Ser Thr  
545 550 555 560 565 570 575

Arg Glu Trp Asn Gly Thr Tyr His Cys Ile Phe Arg Tyr Lys Asn Ser  
565 570 575 580 585 590 595

Tyr Ser Ile Ala Thr Lys Asp Val Ile Val His Pro Leu Pro Leu Lys  
580 585 590 595 600 605 610

Leu Asn Ile Met Val Asp Pro Leu Glu Ala Thr Val Ser Cys Ser Gly  
595 600 605 610 615 620 625

Ser His His Ile Lys Cys Cys Ile Glu Glu Asp Gly Asp Tyr Lys Val  
610 615 620 625 630 635 640

Thr Phe His Met Gly Ser Ser Ser Leu Pro Ala Ala Lys Glu Val Asn  
625 630 635 640 645 650 655

Lys Lys Gln Val Cys Tyr Lys His Asn Phe Asn Ala Ser Ser Val Ser  
645 650 655 660 665 670 675

Trp Cys Ser Lys Thr Val Asp Val Cys Cys His Phe Thr Asn Ala Ala  
660 665 670 675 680 685 690

Asn Asn Ser Val Trp Ser Pro Ser Met Lys Leu Asn Leu Val Pro Gly  
675 680 685 690 695 700 705

Glu Asn Ile Thr Cys Gln Asp Pro Val Ile Gly Val Gly Glu Pro Gly  
690 695 700 705 710 715 720

Lys Val Ile Gln Lys Leu Cys Arg Phe Ser Asn Val Pro Ser Ser Pro  
715 720 725 730 735 740 745

48

WO 01/53451

710

715

PCT/US00/34983

720

Glu Glu Ser Pro Leu Gly Gly Thr Ile Thr Tyr Lys Cys Val Gly Ser  
725 730 735 740 745 750 755

Gln Trp Gly Glu Lys Arg Asn Asp Cys Ile Ser Ala Pro Ile Asn Ser  
740 745 750 755 760 765 770

Leu Leu Gln Met Ala Lys Ala Leu Ile Lys Ser Pro Ser Gln Asp Glu  
755 760 765 770 775 780 785

Met Leu Pro Thr Tyr Leu Lys Asp Leu Ser Ile Ser Ile Asp Lys Ala  
770 775 780 785 790 795 800

Glu His Glu Ile Ser Ser Ser Pro Gly Ser Leu Gly Ala Ile Ile Asn  
785 790 795 800 805 810 815

Ile Leu Asp Leu Leu Ser Thr Val Pro Thr Gln Val Asn Ser Glu Met  
805 810 815 820 825 830 835

Met Thr His Val Leu Ser Thr Val Asn Val Ile Leu Gly Lys Pro Val  
820 825 830 835 840 845 850

Leu Asn Thr Trp Lys Val Leu Gln Gln Gln Trp Thr Asn Gln Ser Ser  
835 840 845 850 855 860 865

Gln Leu Leu His Ser Val Glu Arg Phe Ser Gln Ala Leu Gln Ser Gly  
850 855 860 865 870 875 880

Asp Ser Pro Pro Leu Ser Phe Ser Gln Thr Asn Val Gln Met Ser Ser  
865 870 875 880 885 890 895

Thr Val Ile Lys Ser Ser His Pro Glu Thr Tyr Gln Gln Arg Phe Val  
885 890 895 900 905 910 915

Phe Pro Tyr Phe Asp Leu Trp Gly Asn Val Val Ile Asp Lys Ser Tyr  
900 905 910 915 920 925 930

Leu Glu Asn Leu Gln Ser Asp Ser Ser Ile Val Thr Met Ala Phe Pro  
915 920 925 930 935 940 945

Thr Leu Gln Ala Ile Leu Ala Gln Asp Ile Gln Glu Asn Asn Phe Ala  
930 935 940 945 950 955 960

Glu Ser Leu Val Met Thr Thr Thr Val Ser His Asn Thr Thr Met Pro  
945 950 955 960 965 970 975

Phe Arg Ile Ser Met Thr Phe Lys Asn Asn Ser Pro Ser Gly Gly Glu  
965 970 975 980 985 990 995

Thr Lys Cys Val Phe Thr Asn Phe Arg Leu Ala Asn Asn Thr Gly Gly  
980 985 990 995 1000 1005 1010

Trp Asp Ser Ser Gly Cys Tyr Val Glu Glu Gly Asp Gly Asp Asn Val  
995 1000 1005 1010 1015 1020 1025

Thr Cys Ile Cys Asp His Leu Thr Ser Phe Ser Ile Leu Met Ser  
1010 1015 1020 1025 1030 1035 1040

Pro Asp Ser Pro Asp Pro Ser Ser Leu Leu Gly Ile Leu Leu Asp  
1025 1030 1035 1040 1045 1050 1055

Ile Ile Ser Tyr Val Gly Val Gly Phe Ser Ile Leu Ser Leu Ala  
1055 1060 1065 1070 1075 1080 1085

49

WO 01/34454	PCT/US00/34983	WO 01/34454	PCT/US00/34983
1040	1050	1355	1360
Ala Cys Leu Val Val Glu Ala Val Val Trp Lys Ser Val Thr Lys		Leu Leu Asn	1365
1055	1065	1370	
Asn Arg Thr Ser Tyr Met Arg His Thr Cys Ile Val Asn Ile Ala		<210> 26	
1070	1080	<211> 386	
Ala Ser Leu Leu Val Ala Asn Thr Trp Phe Ile Gly Val Ala Ala		<212> DNA	
1085	1095	<213> Homo sapiens	
Ile Gln Asp Asn Arg Tyr Ile Leu Cys Lys Thr Ala Cys Val Ala		<220>	
1100	1110	<221> misc_feature	
Ala Thr Phe Phe Ile His Phe Phe Tyr Leu Ser Val Phe Phe Trp		<222> (1) .. (386)	
1115	1125	<223> n = A, T, G, or C	
Met Leu Thr Leu Gly Leu Met Leu Phe Tyr Arg Leu Val Phe Ile		<400> 26	
1130	1140	ggagtaacc tgctggcct ggtcctgct ggcgggtgg tgctgcgct gccacctgct	60
Leu His Glu Thr Ser Arg Ser Thr Gln Lys Ala Ile Ala Phe Cys		gtgacctgg ggtgacagc tgcctacacc acctgtatg ccttgccttt ctctccgctc	120
1145	1155	tatgcacgc tctgctggt gcttctgtat gggcacaagc gtctcagcta tcaaacgggg	180
Leu Gly Tyr Gly Cys Pro Leu Ala Ile Ser Val Ile Thr Leu Gly		tctctggccc tctgtctgct ctggggcgcc ttgggtacca cctcttcttc ctctctcttc	240
1160	1170	cgagatactc ccgcgcgcaa ccgctggggg ccttgcctct tctggcttct ctactgtctc	300
Ala Thr Gln Pro Arg Glu Val Tyr Thr Arg Lys Asn Val Cys Trp		ccgctcgcc tgcagttctt cacttgacg cttatgaacc tctactttgc ccaggtgggtg	360
1175	1185	ttcaaggcca aagtgaagcg tctggcc	386
Leu Asn Trp Glu Asp Thr Lys Ala Leu Leu Ala Phe Ala Ile Pro		<210> 27	
1190	1200	<211> 1372	
Ala Leu Ile Ile Val Val Asn Ile Thr Ile Thr Ile Val Val		<212> DNA	
1205	1215	<213> Homo sapiens	
Ile Thr Lys Ile Leu Arg Pro Ser Ile Gly Asp Lys Pro Cys Lys		<400> 27	
1220	1230	gggcgcctg gacacacagg tcccttgcca caggtaacaa cagaggacc ttagggagag	60
Gln Glu Lys Ser Ser Leu Phe Gln Ile Ser Lys Ser Ile Gly Val		aaggtccca gcttctctc caccctgaga accgtgctg agagaccacaa aagcacactg	120
1235	1245	ggcatcagt ctgcagcct accttggcct ccaggtagat gctagtggag ggcgcgcgc	180
Leu Thr Pro Leu Leu Gly Leu Thr Trp Gly Phe Gly Leu Thr Thr		tggcgacaaa ggaagaggca ggcagcaaga gacagcgcg agatgacgaa cagggagctc	240
1250	1260	ctcacaggga cgcggacaag cagcagggcc cagggcgtg ccgcgcgcgc gcatgggaga	300
Val Phe Pro Gly Thr Asn Leu Val Phe His Ile Ile Phe Ala Ile		gcacagcaca cagcagcttc accagcagaa agagcagcga ggcgccacaa aagggccctc	360
1265	1275	ggacagggag caagcctcgg ctcatcttcg ggcgcgacgc ttcactttgg ccttgaacac	420
Leu Asn Val Phe Gln Gly Leu Phe Ile Leu Leu Phe Gly Cys Leu		cacctgggc aaagttagg ttcataaggc gtcaaggtga agaactcag gacagcggg	480
1280	1290	cagcagtaga gaagccagaa gggcaagggg ccccgaggtt gttggcgcg gggagtatct	540
Trp Asp Leu Lys Val Gln Glu Ala Leu Leu Asn Lys Phe Ser Leu		tcggaatta gaaggagaaa aaggttttta cgaaggggg cccagagc agacagaggg	600
1295	1305	ccaggaacac ccgtctgata agctgagacg cttgtgcccc atacagaagc accagccaga	660
Ser Arg Trp Ser Ser Gln His Ser Lys Ser Thr Ser Leu Gly Ser		gctgggcata gacggagaag aagagcaggg catcacaggt ggtgtaggca gctatcagcc	720
1310	1320	ccaggtgtac agcaggtggc agcgaggga ccaagcccg caagcaggca accaggccca	780

gacagggttac tctccatgtc aaggagagga ggcctgaatc caggagacag acaaggaggg 840  
 actgtcgccg ggaaccaggga aataaatag gaggggtggg aattgcaggg tgcagagagg 900  
 gagatggggc gctcaggagg aaagaagttg tggctgacag gcccccctcc tcaacagaga 960  
 gaccttgcc cttggggaac acaggagacc tggggcccaag gccctcaact caggggcatg 1020  
 gaagggaagt cctctttctc caggacagac cggatagggg ttgggggtgt ctctccctg 1080  
 ggcctacccc agagcaagg atggggggat gcagaggccc aaggagagga ctccctggcc 1140  
 tgggtgcttc atgtgctct gggtaggggg ggaagctctc tctctccaa taatgcgat 1200  
 ggggagggcc tctgctctct gctctcgtaa ttctgaaaag ctgccccgc cccacctcc 1260  
 cccaccgct ccacgctoca gctcaagg gagggagag gagggagcca ggtctcagg 1320  
 ccgtgatgg gagagccggc ttaggtgtg gggaccggg gctccagaga aa 1372

<210> 28  
 <211> 3996  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (52)..(3996)

<400> 28  
 gccgacagtc cctcttgctc tgtctccggg attcaggcct cctccctga c atg gag  
 Met Glu  
 1

agt aac ctg tct ggc ctg gtc cct gct gcc ggg ctg gtc cct gcg ctg 105  
 Ser Asn Leu Ser Gly Leu Val Pro Ala Ala Gly Leu Val Pro Ala Leu  
 5 10 15

cca cct gct gtc acc ctg ggg ctg aca gct gcc tac acc acc ctg tat 153  
 Pro Pro Ala Val Thr Leu Gly Leu Thr Ala Ala Tyr Thr Thr Leu Tyr  
 20 25 30

gcc ctg ctg ttc ttc ttc gtc tat gcc cag ctg tgg ctg gtc ctt ctg 201  
 Ala Leu Leu Phe Phe Ser Val Tyr Ala Gln Leu Thr Val Leu Leu  
 35 40 45 50

tat ggg cac aag cgt ctg agc tat cag acg gtc ttc ctg gcc ctg tgt 249  
 Tyr Gly His Lys Arg Leu Ser Tyr Gln Thr Val Phe Leu Ala Leu Cys  
 55 60 65

ctg ctg tgg gcc gcc tgg cgt acc acc ctg ttc ttc ttc ttc ttc cga 297  
 Leu Leu Trp Ala Ala Leu Arg Thr Thr Leu Phe Ser Phe Tyr Phe Arg  
 70 75 80

gat act ccc cgc gcc aac cgc ctg ggg ccc ttg ccc ttc tgg ctt ctg 345  
 Asp Thr Pro Arg Ala Asn Arg Leu Gly Pro Leu Pro Phe Trp Leu Leu  
 85 90 95

tac tgc tgc ccc gtc tgc ctg cag ttc ttc acc ttg acg ctt atg aac 393  
 Tyr Cys Cys Pro Val Cys Leu Gln Phe Phe Thr Leu Thr Leu Met Asn  
 100 105 110

ctc tac ttt gcc cag gtc gtc ttc aag gcc aag gtc aag cgt cgg cgg 441  
 Leu Tyr Phe Ala Gln Val Phe Lys Ala Lys Val Lys Arg Arg Pro  
 115 120 125

gag atg agc cga ggc ttg ctg gct gtc cga ggg gcc ttt gtc ggg gcc 489  
 Glu Met Ser Arg Gly Leu Leu Ala Val Arg Gly Ala Phe Val Gly Ala  
 135 140 145

tgg ctg ctg ttt ctg ctg gtc aac gtc gtc tgt gct gtc ctc tcc cat 537  
 Ser Leu Leu Phe Leu Leu Val Asn Val Leu Cys Ala Val Leu Ser His  
 150 155 160

cgg cgc cgg gca cag ccc tgg gcc ctg ctg ctt gtc cgc gtc ctg gtc 585  
 Arg Arg Arg Ala Gln Pro Trp Ala Leu Leu Leu Val Arg Val Leu Val  
 165 170 175

agg gac tcc ctg ttc gtc atc tgc ggc ctg tct ctt gct gcc tgc ctc 633  
 Ser Asp Ser Leu Phe Val Ile Cys Ala Leu Ser Leu Ala Ala Cys Leu  
 180 185 190

tgc ctg gtc gcc agg cgg ggc ccc tcc act agc atc tac ctg gag gcc 681  
 Cys Leu Val Ala Arg Arg Ala Pro Ser Thr Ser Ile Tyr Leu Glu Ala  
 195 200 205

aag ggg acc agt gtc tgc cag ggc gcc ggc atg ggt ggc gcc atg gtc 729  
 Lys Gly Thr Ser Val Cys Gln Ala Ala Ala Met Gly Gly Ala Met Val  
 215 220 225

ctg ctg tat gcc agc cgg gcc tgc tac aac ctg aca gca ctg gcc ttg 777  
 Leu Leu Tyr Ala Ser Arg Ala Cys Tyr Asn Leu Thr Ala Leu Ala Leu  
 230 235 240

gcc ccc cag agc cgg ctg gac acc ttc gat tac gac tgg tac aat gtc 825  
 Ala Pro Gln Ser Arg Leu Asp Thr Phe Asp Tyr Asp Trp Tyr Asn Val  
 245 250 255

tct gac cag ggc gac ctg gtc aat gac ctg ggg aac aaa ggc tac ctg 873  
 Ser Asp Gln Ala Asp Leu Val Asn Asp Leu Gly Asn Lys Gly Tyr Leu  
 260 265 270

gta ttt ggc ctg atc ctg ttc gtc tgg gag cta ctg ccc acc acc ctg 921  
 Val Phe Gly Leu Ile Leu Phe Val Trp Glu Leu Leu Pro Thr Thr Leu  
 275 280 285

ctg gtc ggc ttc ttc cgg gtc cag cgg ccc cca cag gac ctg agc acc 969  
 Leu Val Gly Phe Phe Arg Val His Arg Pro Pro Gln Asp Leu Ser Thr  
 295 300 305

agc cac atc ctg aat ggg cag gtc ttt gcc tct cgg tcc tac ttc ttt 1017  
 Ser His Ile Leu Asn Gly Gln Val Phe Ala Ser Arg Ser Tyr Phe Phe  
 310 315 320

gac cgg gct ggg cac tgt gaa gat gag ggc tgc tcc tgg gag oac agc 1065  
 Asp Arg Ala Gly His Cys Gln Asp Glu Gly Cys Ser Trp Glu His Ser  
 325 330 335

cgg ggt gag agc acc agc tct tgt gac tgt ggc cct ggt cac tgt cct 1113  
 Arg Gly Glu Ser Thr Ser Ser Cys Asp Cys Gly Pro Gly His Cys Pro  
 340 345 350

gag act gac cct gtt tct ctg ctg cag tat gtc ggg cag tct agg ctc 1161  
 Glu Thr Asp Pro Val Ser Leu Leu Gln Tyr Val Gly Gln Ser Arg Leu  
 355 360 365

WO 01/3454 PCT/US00/34983

355 360 365 370 375 380 385 390 395 400 405 410 415 420 425 430 435 440 445 450 455 460 465 470 475 480 485 490 495 500 505 510 515 520 525 530 535 540 545 550 555 560 565 570 575 580 585 590 600 605 610 615 620 625 630 635 640 645 650 655 660 665 670 675 680 685 690 695 700 705 710 715 720 725 730 735 740 745 750 755 760 765 770 775 780 785 790 795 800 805 810 815 820 825 830 835 840 845 850 855 860 865 870 875 880 885 890 895 900 905 910 915 920 925 930 935 940 945 950 955 960 965 970 975 980 985 990 995

gly ala asp pro glu thr asn ser thr ser asn ser ser his ser ala  
tgg gac ctg ggc agc gcc ttc ttt ttc tca ggg acc atc atc acc acc  
trp asp leu gly ser ala phe phe ser gly thr ile ile thr thr  
atc ggc tat ggc aat gtc ggc ctg cgc aca gat gcc ggg cgc ctc ttc  
ile gly tyr gly asn val ala leu arg thr asp ala gly arg leu phe  
tgc atc ttt tat ggc ctg ggg ggg att ccg ctg ttt ggg atc cta ctg  
cys ile phe tyr ala leu val gly ile pro leu phe gly ile leu leu  
gca ggg gtc ggg gac cgg ctg ggc tcc tcc ctg cgc cat ggc atc ggt  
ala gly val gly asp arg leu gly ser ser leu arg his gly ile gly  
cac att gaa gcc atc ttc ttc aag tgg cac gtc cca cgg gag cta gta  
his ile glu ala ile phe leu lys trp his val pro pro glu leu val  
aga gtc ctg tgg ggc atg ctt ttc ctg ctg atc ggc tgc ctg ctc ttt  
arg val leu ser ala met leu phe leu leu ile gly cys leu leu phe  
gtc ctc acg ccc acg ttc gtc ttc tgc tat atg gag gac tgg agc aag  
val leu thr pro thr phe val phe cys tyr met glu asp trp ser lys  
ctg gag gcc atc tac ttt gtc ata gtc acg ctt acc acc gtc ggc ttt  
leu glu ala ile tyr phe val ile val thr leu thr thr val gly phe  
ggc gac tat gtc ggc ggc gac ccc agg cag gac tcc cgg gcc tat  
gly asp tyr val ala gly ala asp pro arg gln asp ser pro ala tyr  
cag cgc ctg gtc tgg ttc tgg atc ctg ctc ggc ctg gct tac ttc gcc  
gln pro leu val trp phe trp ile leu leu gly leu ala tyr phe ala  
tca gtc ctc acc acc atc ggg aac tgg ctg cga gta gtc tcc cgc cgc  
ser val leu thr thr ile gly asn trp leu arg val val ser arg arg  
act cgg gca gag atg ggc ggc ctc acg gct cag gct gcc agc tgg act  
thr arg ala glu met gly gly leu thr ala ala gln ala ala ser trp thr  
ggc aca gtc aca ggc cgc gtc acc cag cga gcc ggg gcc gcc gcc cgc  
gly thr val thr ala arg val thr gln arg ala gly pro ala ala pro  
ccg cgc gag aag gag cag cca ctg ctg cct cca cgc ccc tgt cca cgc  
pro pro glu lys glu gln pro leu leu pro pro pro pro cys pro ala  
cag cgc ctg ggc agc ccc cga tcc cct tgc ccc ccc gag aag gct cag  
gln pro leu gly arg pro arg ser pro ser pro pro glu lys ala gln  
ggt ggc gac cca gaa acc aac tgc acc agc agc agc cca tca gcc



ctg cct tcc ccg ccc acc ggc tgc ggc ctc gat tat ccc agc gag aac Pro Pro Ser Pro Thr Ala Ser Ala Leu Asp Tyr Pro Ser Glu Asn 870 875 880 2697	ctg ggc ttc atc gac gag tcc tgc gat acg gag agc gag cgc ggc tgc Leu Ala Phe Ile Asp Glu Ser Ser Asp Thr Gln Ser Glu Arg Gly Cys 885 890 895 2745	ctg ctc ccc ccg cgc aga ggt cgc cgc cgc cca aat ccc ccc agg Pro Leu Pro Arg Ala Pro Arg Gly Arg Arg Arg Pro Asn Pro Pro Arg 900 905 910 2793	aag ccc gtc ccc ccg cgc ccc ggc ccc ggc cct ccc cga gac aaa ggc gtc Lys Pro Val Arg Pro Arg Gly Pro Gly Arg Pro Arg Asp Lys Gly Val 915 920 925 2841	ctg acc ccc cca agg ctt tct gtc tgc ctc ccc cgc ggc ggt gta tcc Pro Thr Pro Arg Leu Ser Val Ser Leu Pro Arg Ala Gly Val Ser 930 935 940 2889	ctc aca gca cct cac gac tgt gcc tca aag cct gca tca ata aat gaa Leu Thr Ala Pro His Asp Cys Ala Ser Lys Pro Ala Ser Ile Asn Glu 950 955 960 2937	aac ggt ctc cgc cgc tgc ggc cgt gac gct ccc gga cgc gag tgg gtc Asn Gly Leu His Arg Cys Gly Arg Asp Ala Pro Gly Arg Glu Trp Val 965 970 975 2985	tgg aat tgc ttt cct ccg ggc acc gtc ggc gca cct ctc ggc tcc cgt Trp Asn Cys Phe Pro Arg Ala Thr Val Gly Ala Pro Leu Ala Ser Arg 980 985 990 3033	gac ccc cag ggc gag ggt ccc cgg gca ccc agc ctt ggc tgc ccg Asp Pro Gln Ala Glu Gly Pro Arg Ala Pro Ser Leu Gly Cys Pro 995 1000 1005 3078	cag ccc cca ccc aac ccc acg ttc tac ggc atc ccc aac ccg ggc Gln Pro Pro Pro Asn Pro Thr Phe Tyr Gly Ile Pro Asn Pro Ala 1010 1015 1020 3123	cgg ctc agt tcc cca gcc cgc tot tcc ttc ccg ctc cag cca tcc Arg Leu Ser Ser Pro Ala Arg Ser Ser Phe Pro Leu Gln Pro Ser 1025 1030 1035 3168	gcg acc ctt ggc tcc ctc ctt gta tgt ggc cca cag gtc tgc ctc Ala Thr Leu Gly Ser Leu Leu Val Cys Gly Pro Gln Val Ser Leu 1040 1045 1050 3213	aag tct tcc gac cgc caa ggc tgc gag gag agc gtc cat agc Lys Ser Ser Asp Arg Gln Gly Ser Asp Glu Glu Ser Val His Ser 1055 1060 1065 3258	gac act cgg gac ctc tgg acc acg acc agc ctc tcc cag gca cag Asp Thr Arg Asp Leu Trp Thr Thr Thr Thr Leu Ser Gln Ala Gln 1070 1075 1080 3303	ctg aac atg ccg ctc tcc gag gtc tgc gag ggc ttc gac gat gag Leu Asn Met Pro Leu Ser Glu Val Cys Glu Gly Phe Asp Asp Glu 1085 1090 1095 3348	ggc cgc aac att agc aag acc cgc ggc tgg cgc ccc ccg ggc cgg Gly Arg Asn Ile Ser Lys Thr Arg Gly Trp His Ser Pro Gly Arg 1100 1105 1110 3393	ggc tgc ttg gac gag ggc tac aag gcc agc cac aag ccg gag gaa Gly Ser Leu Asp Glu Gly Tyr Lys Ala Ser His Lys Pro Glu Glu 1115 1120 1125 3438	ctg gac gag cac ggc ctc gtc gag ctc gag ttg cac cgc ggc agc Leu Asp Glu His Ala Leu Val Glu Leu Glu Leu His Arg Gly Ser 1130 1135 1140 3483	tcc atg gaa atc aat ctc ggc gag aag gac act gca tcc cag atc Ser Met Glu Ile Asn Leu Gly Glu Lys Asp Thr Ala Ser Gln Ile 1145 1150 1155 3528	gag gcc gaa aag tct tcc tca atg tca tca ctc aat att ccg aag Glu Ala Glu Lys Ser Ser Ser Met Ser Ser Leu Asn Ile Ala Lys 1160 1165 1170 3573	cac atg ccc cat cga gcc tac tgg gca gag cag cag agc agg gtt His Met Pro His Arg Ala Tyr Trp Ala Glu Gln Gln Ser Arg Val 1175 1180 1185 3618	gga ggc gct ggc gag act ggc cgt ttc ggt ggc ctc cca ctc ccc Gly Ala Gly Glu Thr Gly Arg Phe Gly Gly Leu Pro Leu Pro 1190 1195 1200 3663	ctg atg gaa ctc atg gag aat gaa gct ctc gaa atc ctc acc aaa Leu Met Glu Leu Met Glu Asn Glu Ala Leu Glu Ile Leu Thr Lys 1205 1210 1215 3708	gcc ctc cgg agt aag ctc ccc gcc aac ccc caa gaa ctc cca cga Ala Leu Arg Ser Lys Leu Pro Ala Asn Pro Gln Glu Leu Pro Arg 1220 1225 1230 3753	cag att ctc gtc gat ttt gca ggc ctc ggc ccc agg ggc aga tgc Gln Ile Leu Val Asp Phe Ala Gly Leu Gly Pro Arg Gly Arg Cys 1235 1240 1245 3798	aaa gtt ccc ccg gct aac aca gac ctc agt gcc ctc cgc tac tgc Lys Val Pro Gln Ala Asn Thr Asp Leu Ser Ala Leu Arg Tyr Cys 1250 1255 1260 3843	tac ctc gaa tca tct ggc gtt cct cga atc aca cat ggc ggc ccc Tyr Leu Glu Ser Ser Ala Val Pro Arg Ile Thr His Ala Ala Pro 1265 1270 1275 3888	cct ggc tac cag tta ggc atc ggc agc gac cac ttc ctc act aag Pro Gly Tyr Gln Leu Gly Ile Gly Arg Asp His Phe Leu Thr Lys 1280 1285 1290 3933	gag ctc cag cga tac atc gaa ggc ctc aag aag cgc cgc agc aag Glu Leu Gln Arg Tyr Ile Glu Gly Leu Lys Lys Arg Arg Ser Lys 1295 1300 1305 3978	agg ctc tac gtc aat taa Arg Leu Tyr Val Asn 1310 3996	<210> 29 <211> 1114 <212> PRT <213> Homo sapiens
--	--	--	--	--	--	--	--	--	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

WO 01/K3454	245	250	255
Met Glu Ser Asn Leu Ser Gly Leu Val Pro Ala Ala Gly Leu Val Pro			
1	5	10	15
Ala Leu Pro Pro Ala Val Thr Leu Gly Leu Thr Ala Ala Tyr Thr Thr			
20	25	30	
Leu Tyr Ala Leu Leu Phe Phe Ser Val Tyr Ala Gln Leu Trp Leu Val			
35	40	45	
Leu Leu Tyr Gly His Lys Arg Leu Ser Tyr Gln Thr Val Phe Leu Ala			
50	55	60	
Leu Cys Leu Leu Trp Ala Ala Leu Arg Thr Thr Leu Phe Ser Phe Tyr			
65	70	75	80
Phe Arg Asp Thr Pro Arg Ala Asn Arg Leu Gly Pro Leu Pro Phe Trp			
85	90	95	
Leu Leu Tyr Cys Cys Pro Val Cys Leu Gln Phe Phe Thr Leu Thr Leu			
100	105	110	
Met Asn Leu Tyr Phe Ala Gln Val Val Phe Lys Ala Lys Val Lys Arg			
115	120	125	
Arg Pro Glu Met Ser Arg Gly Leu Leu Ala Val Arg Gly Ala Phe Val			
130	135	140	
Gly Ala Ser Leu Leu Phe Leu Leu Val Asn Val Leu Cys Ala Val Leu			
145	150	155	160
Ser His Arg Arg Arg Ala Gln Pro Trp Ala Leu Leu Leu Val Arg Val			
165	170	175	
Leu Val Ser Asp Ser Leu Phe Val Ile Cys Ala Leu Ser Leu Ala Ala			
180	185	190	
Cys Leu Cys Leu Val Ala Arg Arg Ala Pro Ser Thr Ser Ile Tyr Leu			
195	200	205	
Glu Ala Lys Gly Thr Ser Val Cys Gln Ala Ala Met Gly Gly Ala			
210	215	220	
Met Val Leu Leu Tyr Ala Ser Arg Ala Cys Tyr Asn Leu Thr Ala Leu			
225	230	235	240
Ala Leu Ala Pro Gln Ser Arg Leu Asp Thr Phe Asp Tyr Asp Trp Tyr			

  

WO 01/K3454	245	250	255
Asn Val Ser Asp Gln Ala Asp Leu Val Asn Asp Leu Gly Asn Lys Gly			
260	265	270	
Tyr Leu Val Phe Gly Leu Ile Leu Phe Val Trp Glu Leu Leu Pro Thr			
275	280	285	
Thr Leu Leu Val Gly Phe Phe Arg Val His Arg Pro Pro Gln Asp Leu			
290	295	300	
Ser Thr Ser His Ile Leu Asn Gly Gln Val Phe Ala Ser Arg Ser Tyr			
305	310	315	320
Phe Phe Asp Arg Ala Gly His Cys Glu Asp Glu Gly Cys Ser Trp Glu			
325	330	335	
His Ser Arg Gly Glu Ser Thr Ser Ser Cys Asp Cys Gly Pro Gly His			
340	345	350	
Cys Pro Glu Thr Asp Pro Val Ser Leu Leu Gln Tyr Val Gly Gln Ser			
355	360	365	
Arg Leu Trp Glu Leu Asn Thr Gln Ala Pro Val Pro Leu Thr Leu Gly			
370	375	380	
Pro Cys Ala Lys Phe Val Cys Arg Phe Leu Pro Arg Ile Leu Gly Val			
385	390	395	400
Val Ala Thr Pro Ser Ser Gly Arg Leu Leu Ala Ala Pro Val Ile Asp			
405	410	415	
Ser Gly Ala Gly Thr Pro Gln Gly Arg Leu Ala Gly Arg Gly Ala His			
420	425	430	
Leu Ser Arg Val Gly Ala Ser Gly Ser Gly Val Ala Ala Gly Pro Ala			
435	440	445	
Ala Arg His Ala Pro Arg Arg Arg Cys Ala Asp Ala Gly Glu Ala Val			
450	455	460	
Gly Ala Ser Cys Gly Arg Cys Ala Val Ala Leu Leu Ser Gly Val Cys			
465	470	475	480
Thr Leu Val Ser Thr His Val Cys Val Gly Ser Gly Cys Pro Gly Ala			
485	490	495	

WO 01/4344

PCT/US00/34983

Ala Gly Thr Pro Met Gly Ala Gly Asp Ala Gly Ala Ser Ala Glu Ser  
500 505 510

Ala Val Thr Thr Ala Pro Gln Glu Pro Pro Ala Arg Pro Leu Gln Ala  
515 520 525

Gly Ser Gly Ala Gly Pro Ala Pro Gly Arg Ala Met Arg Ser Thr Thr  
530 535 540

Leu Leu Ala Leu Leu Ala Leu Val Leu Leu Tyr Leu Val Ser Gly Ala  
545 550 555

Leu Val Phe Arg Ala Leu Glu Gln Pro His Glu Gln Ala Gln Arg  
565 570 575

Glu Leu Gly Glu Val Arg Glu Lys Phe Leu Arg Ala His Pro Cys Val  
580 585 590

Ser Asp Gln Glu Leu Gly Leu Leu Ile Lys Glu Val Ala Asp Ala Leu  
595 600 605

Gly Gly Gly Ala Asp Pro Glu Thr Asn Ser Thr Ser Asn Ser Ser His  
610 615 620

Ser Ala Trp Asp Leu Gly Ser Ala Phe Phe Ser Gly Thr Ile Ile  
625 630 635 640

Thr Thr Ile Gly Tyr Gly Asn Val Ala Leu Arg Thr Asp Ala Gly Arg  
645 650 655

Leu Phe Cys Ile Phe Tyr Ala Leu Val Gly Ile Pro Leu Phe Gly Ile  
660 665 670

Leu Leu Ala Gly Val Gly Asp Arg Leu Gly Ser Ser Leu Arg His Gly  
675 680 685

Ile Gly His Ile Glu Ala Ile Phe Leu Lys Trp His Val Pro Pro Glu  
690 695 700

Leu Val Arg Val Leu Ser Ala Met Leu Phe Leu Leu Ile Gly Cys Leu  
705 710 715 720

Leu Phe Val Leu Thr Pro Thr Phe Val Phe Cys Tyr Met Glu Asp Trp  
725 730 735

Ser Lys Leu Glu Ala Ile Tyr Phe Val Ile Val Thr Leu Thr Thr Val  
740 745 750

60

WO 01/4344

PCT/US00/34983

Gly Phe Gly Asp Tyr Val Ala Gly Ala Asp Pro Arg Gln Asp Ser Pro  
755 760 765

Ala Tyr Gln Pro Leu Val Trp Phe Trp Ile Leu Leu Gly Leu Ala Tyr  
770 775 780

Phe Ala Ser Val Leu Thr Thr Thr Ile Gly Asn Trp Leu Arg Val Val Ser  
785 790 795 800

Arg Arg Thr Arg Ala Glu Met Gly Gly Leu Thr Ala Gln Ala Ala Ser  
805 810 815

Trp Thr Gly Thr Val Thr Ala Arg Val Thr Gln Arg Ala Gly Pro Ala  
820 825 830

Ala Pro Pro Pro Glu Lys Glu Gln Pro Leu Leu Pro Pro Pro Cys  
835 840 845

Pro Ala Gln Pro Leu Gly Arg Pro Arg Ser Pro Ser Pro Pro Glu Lys  
850 855 860

Ala Gln Pro Pro Ser Pro Pro Thr Ala Ser Ala Leu Asp Tyr Pro Ser  
865 870 875 880

Glu Asn Leu Ala Phe Ile Asp Glu Ser Ser Asp Thr Gln Ser Glu Arg  
885 890 895

Gly Cys Pro Leu Pro Arg Ala Pro Arg Gly Arg Arg Pro Asn Pro  
900 905 910

Pro Arg Lys Pro Val Arg Pro Arg Gly Pro Gly Arg Pro Arg Asp Lys  
915 920 925

Gly Val Pro Thr Pro Pro Arg Leu Ser Val Ser Leu Pro Arg Ala Gly  
930 935 940

Val Ser Leu Thr Ala Pro His Asp Cys Ala Ser Lys Pro Ala Ser Ile  
945 950 955 960

Asn Glu Asn Gly Leu His Arg Cys Gly Arg Asp Ala Pro Gly Arg Glu  
965 970 975

Trp Val Trp Asn Cys Phe Pro Arg Ala Thr Val Gly Ala Pro Leu Ala  
980 985 990

Ser Arg Asp Pro Gln Ala Glu Gly Pro Arg Ala Pro Ser Leu Gly Cys  
995 1000 1005

61

Pro Gln Pro Pro Pro Asn Pro Thr Phe Tyr Gly Ile Pro Asn Pro  
1010 1015 1020

Ala Arg Leu Ser Ser Pro Ala Arg Ser Ser Phe Pro Leu Gln Pro  
1025 1030 1035

Ser Ala Thr Leu Gly Ser Leu Leu Val Cys Gly Pro Gln Val Ser  
1040 1045 1050

Leu Lys Ser Ser Asp Arg Gln Gly Ser Asp Glu Glu Ser Val His  
1055 1060 1065

Ser Asp Thr Arg Asp Leu Trp Thr Thr Thr Thr Leu Ser Gln Ala  
1070 1075 1080

Gln Leu Asn Met Pro Leu Ser Glu Val Cys Glu Gly Phe Asp Asp  
1085 1090 1095

Glu Gly Arg Asn Ile Ser Lys Thr Arg Gly Trp His Ser Pro Gly  
1100 1105 1110

Arg Gly Ser Leu Asp Glu Gly Tyr Lys Ala Ser His Lys Pro Glu  
1115 1120 1125

Glu Leu Asp Glu His Ala Leu Val Glu Leu Glu Leu His Arg Gly  
1130 1135 1140

Ser Ser Met Glu Ile Asn Leu Gly Glu Lys Asp Thr Ala Ser Gln  
1145 1150 1155

Ile Glu Ala Glu Lys Ser Ser Ser Met Ser Ser Leu Asn Ile Ala  
1160 1165 1170

Lys His Met Pro His Arg Ala Tyr Trp Ala Glu Gln Gln Ser Arg  
1175 1180 1185

Val Gly Gly Ala Gly Glu Thr Gly Arg Phe Gly Gly Leu Pro Leu  
1190 1195 1200

Pro Leu Met Glu Leu Met Glu Asn Glu Ala Leu Glu Ile Leu Thr  
1205 1210 1215

Lys Ala Leu Arg Ser Lys Leu Pro Ala Asn Pro Gln Glu Leu Pro  
1220 1225 1230

Arg Gln Ile Leu Val Asp Phe Ala Gly Leu Gly Pro Arg Gly Arg  
1240 1245 1250

1235 1240 1245

Cys Lys Val Pro Gln Ala Asn Thr Asp Leu Ser Ala Leu Arg Tyr  
1250 1255 1260

Cys Tyr Leu Glu Ser Ser Ala Val Pro Arg Ile Thr His Ala Ala  
1265 1270 1275

Pro Pro Gly Tyr Gln Leu Gly Ile Gly Arg Asp His Phe Leu Thr  
1280 1285 1290

Lys Glu Leu Gln Arg Tyr Ile Glu Gly Leu Lys Lys Arg Arg Ser  
1295 1300 1305

Lys Arg Leu Tyr Val Asn  
1310

<210> 30

<211> 3945

<212> DNA

<213> Homo sapiens

<400> 30

atggagagta acctgtctgg cctggtgacct gctccgggc tgggtcctgc gctgcaacct 60  
gctgtgacct tggggtgac agctgcttac accacctgt atgacctgt cttctctcc 120  
gtctatgcc agctctgggt ggtgctcttg tatggcaca agagtctcag ctatcagag 180  
gtgtctcgg cctctgtct gctctggcc gcttgctga ccaacctctt ctcctctac 240  
ttccgagata tccctgggc caaccgctg gggcccttg cctctgggt tctctactgc 300  
tgccctgtct gctgcagtt ctccacctg acgcttatga acctctactt tgcaccagg 360  
gtgtctcagg ccaaggtgaa gcgtcggcg gagatgagcc gaggttgct cgtgtcaga 420  
ggggccttg tggggcctc gctgtcttt ctgctggtga acgtgtgtg tgtgtgctc 480  
tccctatggc gccgggcaca gccctgggc ctgctgcttg tccggtctct ggtgagcag 540  
tccctgtctg tcatctggc gctgtctctt gctgctgct tctgctcgt cgcacggcg 600  
gcgctctcca ctagcatcta cctggagacc aagggaacca gtgtgtgcca ggcggcgcg 660  
atgggtggcg ccattgtctt gctctatgcc agcggggcct gctacaacct gacagcactg 720  
gcctggccc ccacagagcg gctggacacc ttcgattacg actggtacaa tgtgtctgac 780  
caggcgacc tgggtaatga cctggggaac aaaggctacc tggatttgg cctcctctc 840  
ttcgtgtgg agctactgcc caccacctg ctggtgggt tcttcgggtt gacggcgcc 900  
ccacaggacc ttagcaccag ccacatctc aatgggcagg tcttgcctc tgggtctac 960  
ttcttgacc gggctggga ctgtgaagat gagggtgctt cctgggagca cagcgggggt 1020

gagagcaccg gctcttctgga ctgtatgacct ggtactatgct ctgagactga cctctttctt 1080  
 ctgtatgacct atgtctgggca gttataggctc tggagactga ataccacagg cccagtcacc 1140  
 ctacacctag gccctctgct caagtcttct tgcctcttct tgcctaggat cctgtggggtc 1200  
 gtggttacc cctctctctg cggctctctt gctgtctctg tcatagactc aggggacagg 1260  
 agggccacgg ggcgactggc cgggggggggt gccacctct cgcgctggg cgcctccggg 1320  
 agtgtgtgg cgcgcgggtcc cgcgcgcgc cagctctcga ggcgtctctg tgcggacgcg 1380  
 ggggagggcg tgggagcgag ctgcggggcg tgcgcgggtg cctgtctgtc tggcgtgtgc 1440  
 agctatgtt ccaacacagt gtcgtggggc tctgggtgccc ctggcgccgc cggcacgcgc 1500  
 atggggggcg gggatgcggg ggcgtctctg gagagtgcag tgaagacagc tccccaggag 1560  
 ccccccgcgc ggcctctcca ggcgggcagt ggagctggcc cgggcctctg ggcgcctatg 1620  
 cgcagcacc cgtctctggc cctgtctggc ctggtcttgc tttacttgtt gctgtgtccc 1680  
 ctggtgtccc gggccctgga gcagccccc gagcagcagg cccagaggga gctgggggag 1740  
 gtcgagaga agttcctgag gcccctccg tgtgtgagcg accaggagct gggcctctct 1800  
 atcaaggagg tggctgtatg cctggggagg ggtgcggacc cagaaaccaa ctgcaccagc 1860  
 acagcgacc actcagctg ggacctgggc agcgcttct tttctagg gacctatc 1920  
 accacctcg gctatggcaa tgtggcctg cgcacagatg ccgggcgcct ctcttgatc 1980  
 tttatgccc tgggtgggat tccgtgttt gggatctctac tggcagggt cggggaccgg 2040  
 ctgggctcct cctgcgcga tggcataggt cacattgaag ccatctctt gaagtgcac 2100  
 gtgcacccg agtagtaag agtctgtctg gcgagtcttt tctgtctgat cggctgctg 2160  
 ctctttctc taagccac gttctgtctc tctatattg aggaactggag caagctggag 2220  
 gccatctact ttgtctatgt gacgttacc accgtgggt tggcgacta tgtggccggc 2280  
 gggagcccca ggcaggactc cccggctat cagccgtctg tgtgttctg gatctgtctc 2340  
 ggcctggctt acttcgctc agtctcacc accatcgga actggtctg agtagtctc 2400  
 cgcgcactc gggcagagat gggcggtctc acggctcagg ctgcagctg gactggcaca 2460  
 gtgcagcgc gctgaccca gggagccggg cccgcgcgc cgcgcgcga gaaggagcag 2520  
 ccaatgtctc ctccacgcgc ctgtccagcg cagcgtctg gcaggcccg atccctctg 2580  
 ccccccga aggtctagcc gcttccccc cccacggct cggccctgga ttatccagc 2640  
 gagaacctg ccttctcga cagctctcg gatacgaga gcgagcgcg ctgcccctg 2700  
 ccccgcgcg cagaggtctg ccgcgcacca aatccccc aatccccc ggaagccgt gcggcccg 2760  
 gggcccgcc gtccccgaga caaggcgtg ccgaccccc caagcttct tgtgtctg 2820  
 ccccgggcg ggtatccct cacagacct cagactgtg cctcaagcc tgcataata 2880  
 aatgaaacg gctgcaccg ctgcggcggt gacgtcccg gacgtcgat ggtgtggaat 2940

tgccttctc gggccaccgt gggggcact ctggcctccc gtgaccccca ggcgaggggt 3000  
 cccgggac ccagcttgg ctgcgcgag ccccccacca accccacct ctacgggac 3060  
 cccaaaccg ccggctcag tcccacgc cgtcttctt tcccgctcca gccatcccg 3120  
 acccttgct cctctctgt atgtggcca caggtgtcgc tcaagtcttc cgcgcgcaa 3180  
 ggcggagc aggagagcgt gcatagagac actgggacc tctgggacac gaccacgctg 3240  
 tccaggcag agctgaacat gccgtgtcc gaggctctgc agggcttcca cgaaggggc 3300  
 cgcacatta gcaagaccgc cgggtggcac agcccggggc ggggtctgctt ggcaggggg 3360  
 tacaaggcca gccacaagcc ggaggaaactg gacgagcac gcctgggtgga gctgaggtt 3420  
 caccggcca gctccatgga aatcaatctg ggggagaagg aactgcatc cccagatcg 3480  
 gccgaaagt ctctctaat gtcactatc aatattgca agcacatgcc ccatcgagcc 3540  
 tactggcag agcagcagag cagggttggga ggggtgggg agactggggc ttcgggtgg 3600  
 ctgcactgc cctgatgga actcatggag aatgaactc tggaaactct caccaaagcc 3660  
 ctccggagta agctcccgcc caaccccaa gaactccac gacagattct ggtggtttt 3720  
 gcagggtgg ggcacagggg gagatgcaa gttccacgg ctaacacaga cctgagtgc 3780  
 ctgcgtact gctactcga atcatctcg gttctcgaa tcaacatgc ggcgcacct 3840  
 ggtaccagt tagggatcgg cagggaccac ttcctgacta aggagctgca ggcatacatc 3900  
 gaaggctca agaagcccg gacgaagg ctgactgta attaa 3945

<210> 31  
 <211> 178  
 <212> PRT  
 <213> Homo sapiens  
 <400> 31

Leu Ser Tyr Gln Thr Val Phe Leu Ala Leu Cys Leu Leu Trp Ala Ala  
 1 5 10 15  
 Leu Arg Thr Thr Leu Phe Ser Phe Tyr Phe Arg Asp Thr Pro Arg Ala  
 20 25 30  
 Asn Arg Leu Gly Pro Leu Pro Phe Thr Leu Tyr Cys Cys Pro Val  
 35 40 45  
 Cys Leu Gln Phe Thr Leu Thr Leu Met Asn Leu Tyr Phe Ala Gln  
 50 55 60  
 Val Val Phe Lys Ala Lys Val Lys Arg Arg Pro Gln Met Ser Arg Gly  
 65 70 75 80  
 Leu Leu Ala Val Arg Gly Ala Phe Val Gly Ala Ser Leu Leu Phe Leu  
 85 90 95  
 Leu Val Asn Val Leu Cys Ala Val Leu Ser His Arg Arg Ala Gln  
 100 105 110

Pro Trp Ala Leu Leu Val Arg Val Leu Val Ser Asp Ser Leu Phe  
115 120 125  
Val Ile Cys Ala Leu Ser Leu Ala Ala Cys Leu Cys Leu Val Ala Arg  
130 135 140  
Arg Ala Pro Ser Thr Ser Ile Tyr Leu Glu Ala Lys Gly Thr Ser Val  
145 150 155 160  
Cys Gln Ala Ala Met Gly Ala Met Val Leu Leu Tyr Ala Ser  
165 170 175  
Arg Ala

<210> 32

<211> 334

<212> PRT

<213> Homo sapiens

<220> misc feature

<222> (1) .. (334)

<223> X = any amino acid or a stop codon

<400> 32

Val Arg Gly Leu Gly Pro Arg Leu Pro Val Phe Pro Lys Gly Lys Gly  
1 5 10 15  
Leu Ser Val Glu Glu Gly Leu Ser Ala Thr Thr Ser Phe Leu Leu  
20 25 30

Ser Ala Pro Ser Pro Ser Leu His Pro Ala Ile Pro Thr Pro Arg Ile  
35 40 45

Tyr Phe Pro Gly Pro Ala Asp Ser Pro Ser Leu Ser Val Ser Arg Asp  
50 55 60

Ser Gly Leu Pro Pro Leu Thr Trp Arg Val Thr Cys Leu Gly Leu Val  
65 70 75 80

Ala Cys Leu Pro Gly Leu Val Pro Ala Leu Pro Pro Ala Val Thr Leu  
85 90 95

Gly Leu Thr Ala Ala Tyr Thr Thr Leu Tyr Ala Leu Leu Phe Phe Ser  
100 105 110

Val Tyr Ala Gln Leu Trp Leu Val Leu Arg Met Gly His Lys Arg Leu  
115 120 125

Ser Tyr Gln Thr Val Phe Leu Ala Leu Cys Leu Phe Thr Ala Pro Leu  
130 135 140

Arg Thr Thr Phe Phe Ser Phe Xaa Phe Pro Lys Ile Leu Pro Ala Pro  
145 150 155 160

Asn Asn Ser Trp Gly Pro Leu Pro Phe Thr Leu Leu Tyr Cys Pro  
165 170 175

Val Cys Leu Gln Phe Phe Thr Leu Thr Leu Met Asn Leu Tyr Phe Ala  
180

Gln Val Val Phe Lys Ala Lys Ser Glu Ala Ser Gly Pro Lys Met Ser  
185 190 200 205  
Arg Gly Leu Leu Ala Val Arg Gly Ala Phe Val Gly Ala Ser Leu Leu  
210 215 220  
Phe Leu Leu Val Asn Val Leu Cys Ala Val Leu Val Pro Cys Gly Ala  
225 230 235 240  
Ala Ala Gln Pro Trp Ala Leu Leu Val Arg Val Leu Val Ser Asp  
245 250 255  
Ser Leu Phe Val Ile Cys Ala Leu Ser Leu Ala Ala Cys Leu Phe Leu  
260 265 270  
Cys Arg Gln Ala Gly Ala Leu His Xaa His Leu Pro Gly Gly Gln Gly  
275 280 285  
Arg Ala Ala Ala Leu Met Pro Arg Cys Leu Leu Gly Leu Ser Ala Ala  
290 295 300  
Val Leu Arg Val Xaa Arg Thr Ala Ala Glu Arg Pro Lys Arg His Leu  
305 310 315 320  
Gly Ile Ser Ala Ala Leu Pro Trp Pro Pro Gly Arg Cys  
325 330

<210> 33

<211> 443

<212> DNA

<213> Homo sapiens

<400> 33

cccacggctc cgatgaatc agagcaagag gggctaatta gagattttac tagctagggc 60  
tattaccac aaccatctat ccagactgta gacatagaat caccagatgg ctaggatcct 120  
ggcgacgctg gtccctctt ttcttacc tgaatgcat taaggatgca ttgccaaatg 180  
ctgccctctt ggccctgatga ctacactcca tattgggtcac ctgctcttctt ttctctacag 240  
tctctctcca gacaggcagc ccatacaacc gactacactt gggctcactg aatgaatcac 300  
attctctgpg tgtgctctcc agagatttca tcaagcacc cgcagtgccc ttctggaggc 360  
tctcaaaact ctccactcaa tggttctctga aagtgctggg ttctcactct ctagattgtt 420  
attctcatca gttacatgtg ggg 443

<210> 34

<211> 2735

<212> DNA

<213> Homo sapiens

<400> 34

ttagatttcag ctttgggaat ttgttcagg aactgcatca tccagcccaa aagcaaaaga 60  
ggaaaaatgca taagccaggga taactaaatg ttctttatt taaacttgaa aatgtgactg 120  
caaaaggagt caagaagttt agtaactaac acattctctc ttgttagaaa taactgctca 180

agaagataaa ttgttacata acagttgatt gatgaatatg aaatcagagc aagaggggct 240  
 aattagagaa attactagct agggctattt acccaacaa tctatccaga cgtgagacat 300  
 agaataacca gatggctagg atccgggtgc agctgggttcc ccttttctt taccttgaat 360  
 gtcataaagg atgcatttgc aaatctgcc cctctggcct gatgactaca cctcatattg 420  
 gtcaactgcc ttctttttcc tacagtcttc ctccagacag gcacggcata caaccgacta 480  
 cacttgggct cactgaatga atcaattct tctgtgtgc ctcccagaga ttctatcaaa 540  
 gcaaccggag tggccttctg gaggtctctc aaactctoca ctcaatgttt cctgaagtg 600  
 tegtgttct cactctaga ttgttattct catcagtttac atgtgggttt cacaatttta 660  
 ttcttcagaa tgcagatctg tctttatat cctcgggaaa cacaccttta tatccagtt 720  
 agtactgaca aaaattaaac tgggactgg ccnaaaacag tgcctttct cactttaatc 780  
 tcaataagt agataagact caagtattt ttgtcttgca atggcattga caaatgtttg 840  
 caccaaaaac catgtgaag ttcataaagg aaactgtgat ccaagatoca aggtcaaaaa 900  
 acaaaattca tcaattcagc acaccaccaa ctcaaggct aagcatctta ctgtaattc 960  
 atgtatctg ccatttgcga agtgccaaat tgaattattg atttgtaat aatttccttc 1020  
 cgttggttac ttatataga tattgcaatt ctgtgtgtgc aagtcagcta cactttttct 1080  
 atttgaaaaa caatttctg catttggat ttcaagtata gtgattgtta caaatatgaa 1140  
 ggacttgaat taacagcaag ttttcaagta aaactttact tatgtataac tgaatgagtt 1200  
 cttaagaca ttactaaca atttccaca aactaaaaat ttataaaca ataaataaaa 1260  
 tagactttaa aaaaagggt gtcaacagc tcttggtttt ttgttggttt ctgttgtgt 1320  
 ttttttagt tgaatgggtg aaaaatcaga caatgggtcac agagtctctc ctactgggat 1380  
 ttctctggg cccaaggatt cagatgtctc tcttggggt cttctcctg ttctatgtct 1440  
 tcaacctgct ggggaatggg accatctgg ggctcatctc actggactcc agactccaca 1500  
 ccccatgta cttctctc tcacacctgg ccgtctgcaa catgccttat gcctgcaaca 1560  
 cagtgcacca gatctgggtg aacctctgc atccagccaa gccatctcc ttgtctgggt 1620  
 gcataacata gacctttctc tttttagatt ttgcacatac tgaatgcctc ctgttgggtc 1680  
 tgatgtctca cagatgggtac gtggcatctc gccacctctc ccgatatttc atcatatga 1740  
 cctggaaaagt ctgcactact ctggccatca cttctctggac atgtggctcc cttctggcta 1800  
 tgggtccatgt gagcctcatc ctaagactgc ccttttgggt gccctgtgaa atcaaccact 1860  
 tctctgtga aatctgtctc gtctcaggc tggcctgtgc tgatacttg gccaaaccagg 1920  
 tgggtcatct ttgcagctgc atgttcatcc tgggtggacc actctgctg gtgctgtct 1980  
 cctactcaaa catctctggc gccatctga ggatccagtc tggggagggc cgcgaaggg 2040

cttctcac ctgctctctc cactctctgc tagtgggaact cttcttggc agcgcaatcg 2100  
 tcatgtacat ggcctctaag tcccgcatc ctgaggagca gcagaaggct cttttctctat 2160  
 ttacagttc tttaacccg atgctaaacc cctgattta caactgagg aatgtagagg 2220  
 tcaaggtgc cctgaggaga gcactgtgca aggaagtca ttctaaagag gtgtgacatt 2280  
 tgaactgcca gctcagttg tcaagtgagc tcttgatgcc caattattgc ctcaatccag 2340  
 aaaaattac ttctcttat ctgtgcttta ctgacagaag ggcaagtctt cttctgtttt 2400  
 ttgcagataa aattttagat gtgttgcaat catgtgggttt ctatgagatg tggttttatc 2460  
 agacaatttt ttcttttatt tcacaattac tttaatatct ttgaccagca aatactcca 2520  
 cctctcaggg agaggtatga gctttaggg aaacatctt ggagagggtc ctgtctctcc 2580  
 ctgaggtggg ctttgaatcc agcactcttc ccttggagg gtcaacttggg accagctaac 2640  
 ttctcagggt cctttctctc cagttctgct catcatctg tcatgtaaca ctttagtgtt 2700  
 ctatatttgc atagtgtat ccttccatta gtctg 2735  
 <210> 35  
 <211> 1788  
 <212> DNA  
 <213> Homo sapiens  
 <220>  
 <221> CDS  
 <222> (833) .. (1417)  
 <400> 35  
 atttggcctt cagagccaag aattcggcac gaggatcca aggtcaaaaa acaaaatcaa 60  
 tcaattcagc acaccaccaa ctcaaggct aagcatctta ctgtaatttc attgatgtcg 120  
 ccatttgcga agtgccaaat tgaattattg atttgcatt aattctcttc cgttgggtac 180  
 ttatatagta tattgcaatt ctgtgtgtcg aagtcagcta cactttttct atttgaaaaa 240  
 caatttcttg catttgggat ttcaaggtata gtgattgtta caaatatgaa ggacttgaat 300  
 taacagcaag ttttcaagta aaactttact tatgtataac tgaatgagtt ctttaagaca 360  
 ttactaaca attttcaca aactaaaaat ttataaaca ataaataaaa tagactttta 420  
 aaaaaagcgt gtcaacagc tgttggtttt ttgttggttt ctttggttgt ttttttagtag 480  
 tgaatagggtg aaaaatcaga caatgggtcac agagtctctc ctactgggat ttctctggg 540  
 ccaaggatt cagatgtctc ttttgggtt cttctcctg ttctatgtct tcaacctgtct 600  
 ggggaatggg accatctgg ggtcatctc actggactcc agactccca ccccatgta 660  
 cttctctc tcacacctgg ccgtctcaa catgcctat gccctgaaca cagtgcacca 720  
 gatgtgggtg aactctctgc atccagccaa gccatctcc ttgtctgggt gcatgaata 780  
 gacctttctc tttttagtt ttgcacatac tgaatggctc ctgttgggtc tg atg tcc 838  
 Met Ser

tac gat cgg tac gtc gcc atc tgc cac cct ctc cga tat ttc atc atc  
Tyr Asp Arg Tyr Val Ala Ile Cys His Pro Leu Arg Tyr Phe Ile Ile  
5 10 15 886

atg acc tgg aaa gtc tgc atc act ctg gcc atc act tcc tgg aca tgt  
Met Thr Trp Lys Val Cys Ile Thr Leu Ala Ile Thr Ser Trp Thr Cys  
20 25 30 934

ggc tcc ctc ctg gct atg gct cat gtc agc ctc atc cta aga ctg ccc  
Gly Ser Leu Leu Ala Met Val His Val Ser Leu Ile Leu Arg Leu Pro  
35 40 45 982

ttt tgt ggg cct cgt gaa atc aac cac ttc ttc tgt gaa atc ctg tct  
Phe Cys Gly Pro Arg Glu Ile Asn His Phe Phe Cys Glu Ile Leu Ser  
55 60 65 1030

gtc ctc agg ctg gcc tgt gct gat acc tgg ctc aac cag gtg gtc atc  
Val Leu Arg Leu Ala Cys Ala Asp Thr Trp Leu Asn Gln Val Val Ile  
70 75 80 1078

ttt gca gcc tgc atg ttc atc ctg gtg gga cca ctc tgc ctg gtg ctg  
Phe Ala Ala Cys Met Phe Ile Leu Val Gly Pro Leu Cys Leu Val Leu  
85 90 95 1126

gtc tcc tac tca cac atc ctg ggg gcc atc ctg agg atc cag tct ggg  
Val Ser Tyr Ser His Ile Leu Ala Ala Ile Leu Arg Ile Gln Ser Gly  
100 105 110 1174

ggg ggc cgc aga agc gcc ttc tcc acc tgc tcc tcc cac ctc tgc gta  
Glu Gly Arg Arg Lys Ala Phe Ser Thr Cys Ser Ser His Leu Cys Val  
115 120 125 1222

gtg gga ctc ttc ttt ggc agc gcc atc gtc atg tac atg gcc cct aag  
Val Gly Leu Phe Phe Gly Ser Ala Ile Val Met Tyr Met Ala Pro Lys  
135 140 145 1270

tcc cgc cat cct gag gag cag cag aag gtc ctt ttt cta ttt tac agt  
Ser Arg His Pro Glu Glu Gln Gln Lys Val Leu Phe Leu Phe Tyr Ser  
150 155 160 1318

tct ttc aac ccg atg cta aac ccc ctg att tac aac ctg agg aat gta  
Ser Phe Asn Pro Met Leu Asn Pro Leu Ile Tyr Asn Leu Arg Asn Val  
165 170 175 1366

gag gtc aag ggt gcc ctg agg aga gca ctg tgc aag gaa agt cat tcc  
Glu Val Lys Gly Ala Leu Arg Arg Ala Leu Cys Lys Glu Ser His Ser  
180 185 190 1414

taa gaggtgtgac atttgaactg ccggcctcag ttgtcacgtg gactcttgat  
1467

gccaattat tgcctcaatc cagaaaagt tactctctt tactgtgct ttactgacag  
1527

aagggaagt cttctctgt tttttgcga taaaattta gatgtgtgc attcatggg  
1587

tttctatgag atgtggtttt atcagacaat tttttctttt attcacaat tactttaata  
1647

tctgtaaaat aaagaattat ttttaattcat tttccagtc ccaaaagtta aatacagcc  
1707

acttactct ttaaccaaat gatatagttt ggctctgtgt ccccccacaa atctcatgtc  
1767

aaattgtaat cccgcgatgt g  
1788

<210> 36  
<211> 194  
<212> PRT  
<213> Homo sapiens  
<400> 36

Met Ser Tyr Asp Arg Tyr Val Ala Ile Cys His Pro Leu Arg Tyr Phe  
1 5 10 15

Ile Ile Met Thr Trp Lys Val Cys Ile Thr Leu Ala Ile Thr Ser Trp  
20 25 30

Thr Cys Gly Ser Leu Leu Ala Met Val His Val Ser Leu Ile Leu Arg  
35 40 45

Leu Pro Phe Cys Gly Pro Arg Glu Ile Asn His Phe Phe Cys Glu Ile  
50 55 60

Leu Ser Val Leu Arg Leu Ala Cys Ala Asp Thr Trp Leu Asn Gln Val  
65 70 75 80

Val Ile Phe Ala Ala Cys Met Phe Ile Leu Val Gly Pro Leu Cys Leu  
85 90 95

Val Leu Val Ser Tyr Ser His Ile Leu Ala Ala Ile Leu Arg Ile Gln  
100 105 110

Ser Gly Glu Gly Arg Arg Lys Ala Phe Ser Thr Cys Ser Ser His Leu  
115 120 125

Cys Val Val Gly Leu Phe Phe Gly Ser Ala Ile Val Met Tyr Met Ala  
130 135 140

Pro Lys Ser Arg His Pro Glu Glu Gln Gln Lys Val Leu Phe Leu Phe  
145 150 155 160

Tyr Ser Ser Phe Asn Pro Met Leu Asn Pro Leu Ile Tyr Asn Leu Arg  
165 170 175

Asn Val Glu Val Lys Gly Ala Leu Arg Arg Ala Leu Cys Lys Glu Ser  
180 185 190

His Ser

<210> 37  
<211> 585



<212> DNA  
<213> Homo sapiens

<400> 37  
atgctctacg atcggtacgt ggccatctgc caccctctcc gatatttcat catcatgacc 60  
tggaaagtct gcatactctt ggcatacact tcttggaact gtggctccct cctggctatg 120  
gtccatgtga gctctcactt agactgccc ttttgggc ctctggaat caaccacttc 180  
tctgtgaaa tctgtctgtt cctcaggtg gctgtgctg atacctggct caaccaggtg 240  
gtcatctttg cagctgcgat gttctctctg gtggaccac tctgctcggg gtgggtctcc 300  
tactacaca tctgtggcgc catctgagg atccagtctg gggaggggcc cagaaggcc 360  
ttctccact gctctccca cctctgcta gtggactct tcttggcag cgcctatgct 420  
atgtacatgg cccttaagtc cgcctactct gaggagcagc agaaggctct tttctcttt 480  
taccgttctt tcaaccgat gctaaacccc ctgatttaca acctgaggaa tctagaggtc 540  
aagggtgccc tgaggagagc actgtgcaag gaaatcatt cctaa 585

<210> 38  
<211> 173  
<212> PRT  
<213> Homo sapiens  
<400> 38

Met Ser Tyr Asp Arg Tyr Val Ala Ile Cys His Pro Leu Arg Tyr Phe 15  
1 5 10  
Ile Ile Met Thr Trp Lys Val Cys Ile Thr Leu Ala Ile Thr Ser Trp 30  
20 25  
Thr Cys Gly Ser Leu Leu Ala Met Val His Val Ser Leu Ile Leu Arg 45  
35 40  
Leu Pro Phe Cys Gly Pro Arg Glu Ile Asn His Phe Phe Cys Glu Ile 60  
50 55  
Leu Ser Val Leu Arg Leu Ala Cys Ala Asp Thr Trp Leu Asn Gln Val 80  
65 70 75  
Val Ile Phe Ala Ala Cys Met Phe Ile Leu Val Gly Pro Leu Cys Leu 95  
85 90  
Val Leu Val Ser Tyr Ser His Ile Leu Ala Ala Ile Leu Arg Ile Gln 110  
100 105  
Ser Gly Glu Gly Arg Arg Lys Ala Phe Ser Thr Cys Ser Ser His Leu 125  
115 120  
Cys Val Val Gly Leu Phe Phe Gly Ser Ala Ile Val Met Tyr Met Ala 140  
130 135  
Pro Lys Ser Arg His Pro Glu Gln Gln Lys Val Leu Phe Leu Phe 160  
145 150 155  
Tyr Ser Ser Phe Asn Pro Met Leu Asn Pro Leu Ile Tyr

<210> 39  
<211> 35  
<212> PRT  
<213> Homo sapiens  
<400> 39  
Met Ser Tyr Asp Arg Tyr Val Ala Ile Cys His Pro Leu Arg Tyr Phe 15  
1 5 10  
Ile Ile Met Thr Trp Lys Val Cys Ile Thr Leu Ala Ile Thr Ser Trp 30  
20 25  
Thr Cys Gly 35  
<210> 40  
<211> 159  
<212> PRT  
<213> Homo sapiens  
<400> 40

Ser Leu Leu Ala Met Val His Val Ser Leu Ile Leu Arg Leu Pro Phe 15  
1 5 10  
Cys Gly Pro Arg Glu Ile Asn His Phe Phe Cys Glu Ile Leu Ser Val 30  
20 25  
Leu Arg Leu Ala Cys Ala Asp Thr Trp Leu Asn Gln Val Val Ile Phe 45  
35 40  
Ala Ala Cys Met Phe Ile Leu Val Gly Pro Leu Cys Leu Val Leu Val 60  
50 55  
Ser Tyr Ser His Ile Leu Ala Ala Ile Leu Arg Ile Gln Ser Gly Glu 80  
65 70 75  
Gly Arg Arg Lys Ala Phe Ser Thr Cys Ser Ser His Leu Cys Val Val 95  
85 90  
Gly Leu Phe Phe Gly Ser Ala Ile Val Met Tyr Met Ala Pro Lys Ser 110  
100 105  
Arg His Pro Glu Glu Gln Gln Lys Val Leu Phe Phe Tyr Ser Ser 125  
115 120  
Phe Asn Pro Met Leu Asn Pro Leu Ile Tyr Asn Leu Arg Asn Val Glu 140  
130 135  
Val Lys Gly Ala Leu Arg Arg Ala Leu Cys Lys Glu Ser His Ser 155  
145 150  
<210> 41  
<211> 1782  
<212> DNA  
<213> Homo sapiens  
<220>  
<221> CDS  
<222> (485)...(1411)

<400> 41  
 atttggccct cgaggccaag aattggccac gaggatcca aggtcaaaaa aacaaattca 60  
 toaatcagc acacaccacaa ctacaggct aagcatctta ctgtaattc attgatgctg 120  
 ccatttgca agtgcacaat tgaattattg atttgcaat aatttccttc ggttggttac 180  
 ttatatgta tattgcaatt ctgttgctg aagtcagcta cacittttct atttgaaaaa 240  
 caatttttg catttggat ttccaggata gtgattgta caaatatgaa ggaattgaat 300  
 taacagcaag ttctcaagta aaactttact tatgtataac tgaatgagtt cttaagaca 360  
 ttactaaca attttccaca aactaaaaat ttataaaca atataaaaa tagactttaa 420  
 aaaaaagct gtcacacagc tgcctgtttt ttgtttttt ctgttttgt tttttatgag 480  
 tgaatg gtc aaa aat cag aca atg gtc aca gag ttc ctc cta ctg gga 529  
 Met Val Lys Asn Gln Thr Met Val Thr Glu Phe Leu Leu Leu Gly 15  
 1 5 10  
 ttt ctc ctg ggc cca agg att cag atg ctc ctc ttt ggg ctc ttc ttc 577  
 Phe Leu Leu Gly Pro Arg Ile Gln Met Leu Leu Phe Gly Leu Phe Ser 30  
 20 25  
 ctg ttc tat atc ttc acc ctg ctg ggg aac ggg gcc atc ctg ggg ctc 625  
 Leu Phe Tyr Ile Phe Thr Leu Leu Gly Asn Gly Ala Ile Leu Gly Leu 45  
 35 40  
 atc tca ctg gac tcc aga ctc cat acc ccc atg tac ttc ttc ctc tca 673  
 Ile Ser Leu Asp Ser Arg Leu His Thr Pro Met Tyr Phe Phe Leu Ser 60  
 50 55  
 cac ctg gct gtc gac atc gcc tac acc cgc aac acg ctg ccc cag 721  
 His Leu Ala Val Asp Ile Ala Tyr Thr Arg Asn Thr Val Pro Gln 75  
 65 70  
 atg ctg gcg aac ctc ctg cat cca gcc aag ccc atc tcc ttt gct ggt 769  
 Met Leu Ala Asn Leu Leu His Pro Ala Lys Pro Ile Ser Phe Ala Gly 95  
 80 85 90  
 tgc atg acg cag acc ttt ctc tgt tgg agt ttt gga cac agc gaa tgt 817  
 Cys Met Thr Gln Thr Phe Leu Cys Leu Ser Phe Gly His Ser Glu Cys 110  
 100 105  
 ctc ctg ctg gtc atg tcc tac gat cgt tac ctg gcc atc tgc cac 865  
 Leu Leu Leu Val Leu Met Ser Tyr Asp Arg Tyr Val Ala Ile Cys His 120  
 115 125  
 cct ctc cga tac tcc gtc atc atg acc tgc tgc atc act ctg gcc atc 913  
 Pro Leu Arg Tyr Ser Val Ile Met Thr Cys Cys Ile Thr Leu Ala Ile 140  
 130 135  
 act tcc tgg aca tgt ggc tcc ctc ctg gct atg gtc cat gtc agc ctc 961  
 Thr Ser Trp Thr Cys Gly Ser Leu Leu Ala Met Val His Val Ser Leu 155  
 145 150  
 atc cta aga ctg ccc ttt tgt ggg cct cgt gaa atc aac cac ttc ttc 1009  
 Ile Leu Arg Leu Pro Phe Cys Gly Pro Arg Glu Ile Asn His Phe Phe 175  
 160 165 170  
 tgt gaa atc ctg tct gtc ctc agg ctg gcc tgt gct gat acc tgg ctc 1057  
 74

Cys Glu Ile Leu Ser Val Leu Arg Leu Ala Cys Ala Asp Thr Trp Leu 180  
 185  
 aac cag gtc gtc atc ttt gca gcc tgc atg ttc atc ctg gtc gga cca 1105  
 Asn Gln Val Val Ile Phe Ala Ala Cys Met Phe Ile Leu Val Gly Pro 205  
 195 200  
 ctc tgc ctg gtc ctg gtc tcc tca cac atc atc ctg ggc gcc atc ctg 1153  
 Leu Cys Leu Val Leu Val Ser Tyr Ser His Ile Leu Ala Ala Ile Leu 220  
 210 215  
 agg atc cag tct ggg gag ggc cgc aga aag gcc ttc tcc acc tgc tcc 1201  
 Arg Ile Gln Ser Gly Glu Gly Arg Arg Lys Ala Phe Ser Thr Cys Ser 235  
 225 230  
 tcc cac ctc tgc gta gtc gga ctc ttc ttt ggc agc gcc atc gtc atg 1249  
 Ser His Leu Cys Val Val Gly Leu Phe Gly Ser Ala Ile Val Met 255  
 240 245  
 tac atg gcc cct aag tcc cgc cat cct gag gag cag cag aag gtc ctt 1297  
 Tyr Met Ala Pro Lys Ser Arg His Pro Glu Glu Gln Lys Val Leu 265  
 260  
 ttt cta ttt tac agt tct ttc aac cgc atg cta aac ccc ctg att tac 1345  
 Phe Leu Phe Tyr Ser Ser Phe Asn Pro Met Leu Asn Pro Leu Ile Tyr 285  
 275 280  
 aac ctg agg aat gta gag gtc aag ggt gcc ctg agg aga gca ctg tgc 1393  
 Asn Leu Arg Asn Val Glu Val Lys Gly Ala Leu Arg Arg Ala Leu Cys 300  
 290 295  
 aag gaa agt cat tcc taa gaggtgac attgaaactg ccagctcag 1441  
 Lys Glu Ser His Ser 305  
 ttgtcaagtg gactcttgat gcccaattat tgcctcaate cagaaaagtt tactctctt 1501  
 tatctgtgct ttactgacag aagggaagt ctctctcgt ttttgcaga taaaatttta 1561  
 gatgtgtgc attcattggg ttctatgag atgtgtttt atcagacaat tttttcttt 1621  
 atttcacaat tactttaata tctgtaaat aagaattat ttttaattcat tttccagtc 1681  
 ccaaaagtta aatacagccc acttacttct ttaaccaaat gatatagttt ggctctgtg 1741  
 cccaccccaa atctcatgct aaattgaat cccgcatgt g 1782  
 <210> 42  
 <211> 308  
 <212> PRT  
 <213> Homo sapiens  
 <400> 42  
 Met Val Lys Asn Gln Thr Met Val Thr Glu Phe Leu Leu Gly Phe 15  
 1 5 10  
 Leu Leu Gly Pro Arg Ile Gln Met Leu Leu Phe Gly Leu Phe Ser Leu 30  
 20 25

Phe Tyr Ile Phe Thr Leu Leu Gly Asn Gly Ala Ile Leu Gly Leu Ile  
35 40 45

Ser Leu Asp Ser Arg Leu His Thr Pro Met Tyr Phe Phe Leu Ser His  
50 55 60

Leu Ala Val Val Asp Ile Ala Tyr Thr Arg Asn Thr Val Pro Gln Met  
65 70 75

Leu Ala Asn Leu Leu His Pro Ala Lys Pro Ile Ser Phe Ala Gly Cys  
85 90 95

Met Thr Gln Thr Phe Leu Cys Leu Ser Phe Gly His Ser Glu Cys Leu  
100 105 110

Leu Leu Val Leu Met Ser Tyr Asp Arg Tyr Val Ala Ile Cys His Pro  
115 120 125

Leu Arg Tyr Ser Val Ile Met Thr Cys Cys Ile Thr Leu Ala Ile Thr  
130 135 140

Ser Trp Thr Cys Gly Ser Leu Leu Ala Met Val His Val Ser Leu Ile  
145 150 155

Leu Arg Leu Pro Phe Cys Gly Pro Arg Glu Ile Asn His Phe Phe Cys  
165 170 175

Glu Ile Leu Ser Val Leu Arg Leu Ala Cys Ala Asp Thr Trp Leu Asn  
180 185 190

Gln Val Val Ile Phe Ala Ala Cys Met Phe Ile Leu Val Gly Pro Leu  
195 200 205

Cys Leu Val Leu Val Ser Tyr Ser His Ile Leu Ala Ala Ile Leu Arg  
210 215 220

Ile Gln Ser Gly Glu Gly Arg Arg Lys Ala Phe Ser Thr Cys Ser Ser  
225 230 235 240

His Leu Cys Val Val Gly Leu Phe Phe Gly Ser Ala Ile Val Met Tyr  
245 250 255

Met Ala Pro Lys Ser Arg His Pro Glu Glu Gln Lys Val Leu Phe  
260 265 270

Leu Phe Tyr Ser Ser Phe Asn Pro Met Leu Asn Pro Leu Ile Tyr Asn  
275 280 285

Leu Arg Asn Val Glu Val Lys Gly Ala Leu Arg Arg Ala Leu Cys Lys  
290 295 300

Glu Ser His Ser  
305

<210> 43

<211> 927

<212> DNA

<213> Homo sapiens

<400> 43

atggtgaata atcagacaat ggtcacagag ttctctctac tgggattttt cctggggccca 60

aggattcaga tgcctctctt tgggctcttc tccctgttct atattctac cctgctgggg 120

aacggggcca tcttgggggt catctactg gactccagac tccatacccc catgtacttc 180

ttctctcac acctggctgt cgtcgacatc gctacaccc gcaacacggt gccccagatg 240

ctggcgacc tctgcatcc agccaagccc atctctttg.ctgggttgcac gacgcagacc 300

ttctctgtt tgaattttgg acaacgcaa tgtctctgc tgggtcatgt gctctacgat 360

cgttaagtgg ccattcgcca cctctccga tactccgtca tcattgacctg.ctgcatcact 420

ctggccatca ctctctggac atgtggctcc ctctggcta tgggtccatgt gagctcacc 480

ctaagactgc ccttttgggt gctctgtgaa atcaacact tctctgtga aatcctgtct 540

gtctcaggc tggcctgtgc tgatactgg ctaaacagg tgggtcatctt tgcagcctgc 600

atgtctacc tgggggacc actctgctg gtctgtgtct cctactcaca catcctgggg 660

gccatctga ggaaccagtc tggggagggc cgcagaaagg cctctccac ctgctcctcc 720

cacctctgg tagtgggact ctctttggc agcgccatcg tcattacat ggcctctaag 780

tccgcctac ctgaggagca gcagaaggtc cttttctat ttacagttc ttcaaccgg 840

atgttaacc ccttgattta caacctgagg aatgtagagg tcaagggtgc cctgaggaga 900

gcactgtca aggaagtca ttcctaa 927

<210> 44

<211> 248

<212> PRT

<213> Homo sapiens

<400> 44

Gly Asn Gly Ala Ile Leu Gly Leu Ile Ser Leu Asp Ser Arg Leu His  
1 5 10 15

Thr Pro Met Tyr Phe Phe Leu Ser His Leu Ala Val Val Asp Ile Ala  
20 25 30

Tyr Thr Arg Asn Thr Val Pro Gln Met Leu Ala Asn Leu Leu His Pro  
35 40 45

Ala Lys Pro Ile Ser Phe Ala Gly Cys Met Thr Gln Thr Phe Leu Cys  
 50 55 60  
 Leu Ser Phe Gly His Ser Glu Cys Leu Leu Val Leu Met Ser Tyr  
 65 70 75 80  
 Asp Arg Tyr Val Ala Ile Cys His Pro Leu Arg Tyr Ser Val Ile Met  
 85 90 95  
 Thr Cys Cys Ile Thr Ser Thr Leu Ala Ile Thr Ser Trp Thr Cys Gly Ser Leu  
 100 105 110  
 Leu Ala Met Val His Val Ser Leu Ile Leu Arg Leu Pro Phe Cys Gly  
 115 120 125  
 Pro Arg Glu Ile Asn His Phe Phe Cys Glu Ile Leu Ser Val Leu Arg  
 130 135 140  
 Leu Ala Cys Ala Asp Thr Trp Leu Asn Gln Val Val Ile Phe Ala Ala  
 145 150 155 160  
 Cys Met Phe Ile Leu Val Gly Pro Leu Cys Leu Val Leu Val Ser Tyr  
 165 170 175  
 Ser His Ile Leu Ala Ala Ile Leu Arg Ile Gln Ser Gly Glu Gly Arg  
 180 185 190  
 Arg Lys Ala Phe Ser Thr Cys Ser Ser His Leu Cys Val Val Gly Leu  
 195 200 205  
 Phe Phe Gly Ser Ala Ile Val Met Tyr Met Ala Pro Lys Ser Arg His  
 210 215 220  
 Pro Glu Glu Gln Gln Lys Val Leu Phe Leu Phe Tyr Ser Ser Phe Asn  
 225 230 235  
 Pro Met Leu Asn Pro Leu Ile Tyr  
 245  
 <210> 45  
 <211> 42  
 <212> PRT  
 <213> Homo sapiens  
 <400> 45  
 Met Val Lys Asn Gln Thr Met Val Thr Glu Phe Leu Leu Gly Phe  
 1 5 10 15  
 Leu Leu Gly Pro Arg Ile Gln Met Leu Leu Phe Gly Leu Phe Ser Leu  
 20 25 30  
 Phe Tyr Ile Phe Thr Leu Leu Gly Asn Gly  
 35 40  
 <210> 46  
 <211> 266  
 <212> PRT  
 <213> Homo sapiens  
 <400> 46

Ala Ile Leu Gly Leu Ile Ser Leu Asp Ser Arg Leu His Thr Pro Met  
 1 5 10 15  
 Tyr Phe Phe Leu Ser His Leu Ala Val Val Asp Ile Ala Tyr Thr Arg  
 20 25 30  
 Asn Thr Val Pro Gln Met Leu Ala Asn Leu Leu His Pro Ala Lys Pro  
 35 40 45  
 Ile Ser Phe Ala Gly Cys Met Thr Gln Thr Phe Leu Cys Leu Ser Phe  
 50 55 60  
 Gly His Ser Glu Cys Leu Leu Val Leu Met Ser Tyr Asp Arg Tyr  
 65 70 75 80  
 Val Ala Ile Cys His Pro Leu Arg Tyr Ser Val Ile Met Thr Cys Cys  
 85 90 95  
 Ile Thr Leu Ala Ile Thr Ser Trp Thr Cys Gly Ser Leu Leu Ala Met  
 100 105 110  
 Val His Val Ser Leu Ile Leu Arg Leu Pro Phe Cys Gly Pro Arg Glu  
 115 120 125  
 Ile Asn His Phe Phe Cys Glu Ile Leu Ser Val Leu Arg Leu Ala Cys  
 130 135 140  
 Ala Asp Thr Trp Leu Asn Gln Val Val Ile Phe Ala Ala Cys Met Phe  
 145 150 155 160  
 Ile Leu Val Gly Pro Leu Cys Leu Val Leu Val Ser Tyr Ser His Ile  
 165 170 175  
 Leu Ala Ala Ile Leu Arg Ile Gln Ser Gly Glu Gly Arg Arg Lys Ala  
 180 185 190  
 Phe Ser Thr Cys Ser Ser His Leu Cys Val Val Gly Leu Phe Gly  
 195 200 205  
 Ser Ala Ile Val Met Tyr Met Ala Pro Lys Ser Arg His Pro Glu Glu  
 210 215 220  
 Gln Gln Lys Val Leu Phe Leu Phe Tyr Ser Ser Phe Asn Pro Met Leu  
 225 230 235 240  
 Asn Pro Leu Ile Tyr Asn Leu Arg Asn Val Glu Val Lys Gly Ala Leu  
 245 250 255  
 Arg Arg Ala Leu Cys Lys Glu Ser His Ser  
 260 265  
 <210> 47  
 <211> 353  
 <212> PRT  
 <213> Homo sapiens  
 <220>  
 <221> misc feature  
 <222> (1) .. (353)  
 <223> X = any amino acid or a stop codon.  
 <400> 47

Arg His Leu Leu Thr Ile Phe His Lys Leu Lys Ile Tyr Lys Thr Ile 15  
 1 5 10  
 Asn Lys Ile Asp Phe Lys Lys Lys Arg Val Thr Gln Leu Leu Val Phe 30  
 20 25  
 Cys Leu Phe Leu Cys Leu Phe Phe Ser Ser Glu Met Val Lys Asn Gln 45  
 35 40  
 Thr Met Val Thr Glu Phe Leu Leu Leu Gly Phe Leu Leu Gly Pro Arg 60  
 50 55  
 Ile Gln Met Leu Leu Phe Gly Leu Phe Ser Leu Phe Tyr Val Phe Thr 80  
 65 70 75  
 Leu Leu Gly Asn Gly Thr Ile Leu Gly Leu Ile Ser Leu Asp Ser Arg 95  
 85 90  
 Leu His Thr Pro Met Tyr Phe Phe Leu Ser His Leu Ala Val Val Asn 110  
 100 105  
 Ile Ala Tyr Ala Cys Asn Thr Val Pro Gln Met Leu Val Asn Leu Leu 125  
 115 120  
 His Pro Ala Lys Pro Ile Ser Phe Ala Gly Cys Met Thr Xaa Thr Phe 140  
 130 135  
 Leu Phe Leu Ser Phe Ala His Thr Glu Cys Leu Leu Leu Val Leu Met 160  
 145 150 155  
 Ser Tyr Asp Arg Tyr Val Ala Ile Cys His Pro Leu Arg Tyr Phe Ile 175  
 165 170  
 Ile Met Thr Trp Lys Val Cys Ile Thr Leu Ala Ile Thr Ser Trp Thr 190  
 180 185  
 Cys Gly Ser Leu Leu Ala Met Val His Val Ser Leu Ile Leu Arg Leu 205  
 195 200  
 Pro Phe Cys Gly Pro Arg Glu Ile Asn His Phe Phe Cys Glu Ile Leu 220  
 210 215  
 Ser Val Leu Arg Leu Ala Cys Ala Asp Thr Trp Leu Asn Gln Val Val 240  
 225 230 235  
 Ile Phe Ala Ala Cys Met Phe Ile Leu Val Gly Pro Leu Cys Leu Val 255  
 245 250  
 Leu Val Ser Tyr Ser His Ile Leu Ala Ala Ile Leu Arg Ile Gln Ser 270  
 260 265  
 Gly Glu Gly Arg Arg Lys Ala Phe Ser Thr Cys Ser Ser His Leu Cys 285  
 275 280  
 Val Val Gly Leu Phe Phe Gly Ser Ala Ile Val Met Tyr Met Ala Pro 300  
 290 295  
 Lys Ser Arg His Pro Glu Glu Gln Lys Val Leu Phe Leu Phe Tyr 320  
 305 310 315  
 Ser Ser Phe Asn Pro Met Leu Asn Pro Leu Ile Tyr Asn Leu Arg Asn 335  
 325 330  
 80

Val Glu Val Lys Gly Ala Leu Arg Arg Ala Leu Cys Lys Glu Ser His 345  
 340 350  
 Ser  
 <210> 48  
 <211> 517  
 <212> PRT  
 <213> Homo sapiens  
 <400> 48  
 Glu Asn Trp Arg Gln Lys Lys Lys Thr Leu Leu Val Ala Ile Asp Arg 15  
 1 5 10  
 Ala Cys Pro Glu Ser Gly His Pro Arg Val Leu Ala Asp Ser Phe Pro 30  
 20 25  
 Gly Ser Ser Pro Tyr Glu Gly Tyr Asn Tyr Gly Ser Phe Glu Asn Val 45  
 35 40  
 Ser Gly Ser Thr Asp Gly Leu Val Asp Ser Ala Gly Thr Gly Asp Leu 60  
 50 55  
 Ser Tyr Gly Tyr Gln Gly His Asp Gln Phe Lys Arg Arg Leu Pro Ser 80  
 65 70 75  
 Gly Gln Met Arg Gln Leu Cys Ile Ala Met Gly Arg Ser Phe Glu Pro 95  
 85 90  
 Val Gly Thr Arg Pro Arg Val Asp Ser Met Ser Val Glu Glu Asp 110  
 100 105  
 Asp Tyr Asp Thr Leu Thr Asp Ile Asp Ser Asp Lys Asn Val Ile Arg 125  
 115 120  
 Thr Lys Gln Tyr Leu Tyr Val Ala Asp Leu Ala Arg Lys Asp Lys Arg 140  
 130 135  
 Val Leu Arg Lys Lys Tyr Gln Ile Tyr Phe Thr Asn Ile Ala Thr Ile 160  
 145 150 155  
 Ala Val Phe Tyr Ala Leu Pro Val Val Gln Leu Val Ile Thr Tyr Gln 175  
 165 170  
 Thr Val Val Asn Val Thr Gly Asn Gln Asp Ile Cys Tyr Tyr Asn Phe 190  
 180 185  
 Leu Cys Ala His Pro Leu Gly Asn Leu Ser Ala Phe Asn Asn Ile Leu 205  
 195 200  
 Ser Asn Leu Gly Tyr Ile Leu Leu Gly Leu Leu Phe Leu Leu Ile Ile 220  
 210 215  
 Leu Gln Arg Glu Ile Asn His Asn Arg Ala Leu Leu Arg Asn Asp Leu 240  
 225 230 235  
 Cys Ala Leu Glu Cys Gly Ile Pro Lys His Phe Gly Leu Phe Tyr Ala 255  
 245 250  
 Met Gly Thr Ala Leu Met Met Glu Gly Leu Leu Ser Ala Cys Tyr His 81

WO 01/3444 260 265 270 PCT/US00/34983

Lys Met Phe Leu Tyr Leu Ser Asp Leu Ser Arg Lys Asp Arg Arg Ile  
35 40 45

Val Ser Lys Lys Tyr Lys Ile Tyr Phe Trp Asn Ile Ile Thr Ile Ala  
50 55 60

Val Phe Tyr Ala Leu Pro Val Ile Gln Leu Val Ile Thr Tyr Gln Thr  
65 70 75 80

Val Val Asn Val Thr Gly Asn Gln Asp Ile Cys Tyr Tyr Asn Phe Leu  
85 90 95

Cys Ala His Pro Leu Gly Val Leu Ser Ala Phe Asn Asn Ile Leu Ser  
100 105 110

Asn Leu Gly His Val Leu Leu Gly Phe Leu Phe Leu Ile Val Leu  
115 120 125

Arg Arg Asp Ile Leu His Arg Arg Ala Leu Glu Ala Lys Asp Ile Phe  
130 135 140

Ala Val Glu Tyr Gly Ile Pro Lys His Phe Gly Leu Phe Tyr Ala Met  
145 150 155 160

Gly Ile Ala Leu Met Met Glu Gly Val Leu Ser Ala Cys Tyr His Val  
165 170 175

Cys Pro Asn Tyr Ser Asn Phe Gln Phe Asp Thr Ser Phe Met Tyr Met  
180 185 190

Ile Ala Gly Leu Cys Met Leu Lys Leu Tyr Gln Thr Arg His Pro Asp  
195 200 205

Ile Asn Ala Ser Ala Tyr Ser Ala Tyr Ala Ser Phe Ala Val Ile  
210 215 220

Met Val Thr Val Leu Gly Val Val Phe Gly Lys Asn Asp Val Trp Phe  
225 230 235 240

Trp Val Ile Phe Ser Ala Ile His Val Leu Ala Ser Leu Ala Leu Ser  
245 250 255

Thr Gln Ile Tyr Tyr Met Gly Arg Phe Lys Ile Asp Val Ser Asp Thr  
260 265 270

Asp Leu Gly Ile Phe Arg Arg Ala Ala Met Val Phe Tyr Thr Asp Cys  
275 280 285

Ile Gln Gln Cys Ser Arg Pro Leu Tyr Met Asp Arg Met Val Leu Leu  
290 295 300

Val Val Gly Asn Leu Val Asn Trp Ser Phe Ala Leu Phe Gly Leu Ile  
305 310 315 320

Tyr Arg Pro Arg Asp Phe Ala Ser Tyr Met Leu Gly Ile Phe Ile Cys  
325 330 335

Asn Leu Leu Leu Tyr Leu Ala Phe Tyr Ile Ile Met Lys Leu Arg Ser  
340 345 350

Ser Glu Lys Val Leu Pro Val Pro Leu Phe Cys Ile Val Ala Thr Ala  
355 360 365

WO 01/3444 260 265 270 PCT/US00/34983

Val Cys Pro Asn Tyr Thr Asn Phe Gln Phe Asp Thr Ser Phe Met Tyr  
275 280 285

Met Ile Ala Gly Leu Cys Met Leu Lys Leu Tyr Gln Lys Arg His Pro  
290 295 300

Asp Ile Asn Ala Ser Ala Tyr Ser Ala Tyr Ala Cys Leu Ala Ile Val  
305 310 315 320

Ile Phe Phe Ser Val Leu Gly Val Val Phe Gly Lys Gly Asn Thr Ala  
325 330 335

Phe Trp Ile Val Phe Ser Ile Ile His Ile Ile Ala Thr Leu Leu Leu  
340 345 350

Ser Thr Gln Leu Tyr Tyr Met Gly Arg Trp Lys Leu Asp Ser Gly Ile  
355 360 365

Phe Arg Arg Ile Leu His Val Leu Tyr Thr Asp Cys Ile Arg Gln Cys  
370 375 380

Ser Gly Pro Leu Tyr Val Asp Arg Met Val Leu Leu Val Met Gly Asn  
385 390 395 400

Val Ile Asn Trp Ser Leu Ala Ala Tyr Gly Leu Ile Met Arg Pro Asn  
405 410 415

Asp Phe Ala Ser Tyr Leu Leu Ala Ile Gly Ile Cys Asn Leu Leu Leu  
420 425 430

Tyr Phe Ala Phe Tyr Ile Ile Met Lys Leu Arg Ser Gly Glu Arg Ile  
435 440 445

Lys Leu Ile Pro Leu Leu Cys Ile Val Cys Thr Ser Val Val Trp Gly  
450 455 460

Phe Ala Leu Phe Phe Phe Gln Gly Leu Ser Thr Trp Gln Lys Thr  
465 470 475 480

Pro Ala Glu Ser Arg Glu His Asn Arg Asp Cys Ile Leu Leu Asp Phe  
485 490 495

Phe Asp Asp His Asp Ile Trp His Phe Leu Ser Ser Ile Ala Met Phe  
500 505 510

Gly Ser Phe Leu Val  
515

<210> 49  
<211> 444  
<212> PRT  
<213> Homo sapiens

<400> 49

Gly His Arg Ala Ser Gln Thr Gln Thr Ala Pro Val Glu Glu Ser Asp  
1 5 10 15

Phe Asp Thr Met Pro Asp Ile Glu Ser Asp Lys Asn Ile Ile Arg Thr  
20 25 30

WO 01/3454 PCT/US00/34983

Val Met Trp Ala Ala Leu Tyr Phe Phe Gln Asn Leu Ser Ser 370 375 380

Trp Glu Gly Thr Pro Ala Glu Ser Arg Glu Lys Asn Arg Glu Cys Ile 385 390 395 400

Leu Leu Asp Phe Phe Asp Asp His Asp Ile Trp His Phe Leu Ser Ala 405 410 415

Thr Ala Leu Phe Phe Ser Phe Leu Val Leu Leu Thr Leu Asp Asp 420 425 430

Leu Asp Val Val Arg Arg Asp Gln Ile Pro Val Phe 435 440

<210> 50  
<211> 267  
<212> PRT  
<213> Homo sapiens

<400> 50

Leu Phe Pro Gln Trp His Leu Pro Ile Lys Ile Ala Ala Ile Ile Ala 1 5 10 15

Ser Leu Thr Phe Leu Tyr Thr Leu Leu Arg Glu Val Ile His Pro Leu 20 25 30

Ala Thr Ser His Gln Gln Tyr Phe Tyr Lys Ile Pro Ile Leu Val Ile 35 40 45

Asn Lys Val Leu Pro Met Val Ser Ile Thr Leu Leu Ala Leu Val Tyr 50 55 60

Leu Pro Gly Val Ile Ala Ala Ile Val Gln Leu His Asn Gly Thr Lys 65 70 75 80

Tyr Lys Lys Phe Pro His Trp Leu Asp Lys Trp Met Leu Thr Arg Lys 85 90 95

Gln Phe Gly Leu Leu Ser Phe Phe Ala Val Leu His Ala Ile Tyr 100 105 110

Ser Leu Ser Tyr Pro Met Arg Ser Tyr Arg Tyr Lys Leu Leu Asn 115 120 125

Trp Ala Tyr Gln Gln Val Gln Gln Asn Lys Glu Asp Ala Trp Ile Glu 130 135 140

His Asp Val Trp Arg Met Glu Ile Tyr Val Ser Leu Gly Ile Val Gly 145 150 155 160

Leu Ala Ile Leu Ala Leu Leu Ala Val Thr Ser Ile Pro Ser Val Ser 165 170 175

Asp Ser Leu Thr Trp Arg Glu Phe His Tyr Ile Gln Ser Lys Leu Gly 180 185 190

Ile Val Ser Leu Leu Leu Gly Thr Ile His Ala Leu Ile Phe Ala Trp 195 200 205

Asn Lys Trp Ile Asp Ile Lys Gln Phe Val Trp Tyr Thr Pro Thr 210 215 220

Phe Met Ile Ala Val Phe Leu Pro Ile Val Val Leu Ile Phe Lys Ser 225 230 235 240

Ile Leu Phe Leu Pro Cys Leu Arg Lys Lys Ile Leu Lys Ile Arg His 250 255

84

WO 01/3454 PCT/US00/34983

Phe Met Ile Ala Val Phe Leu Pro Ile Val Leu Ile Phe Lys Ser 225 230 235 240

Ile Leu Phe Leu Pro Cys Leu Arg Lys Lys Ile Leu Lys Ile Arg His 250 255

Gly Trp Glu Asp Val Thr Lys Ile Asn Lys Thr 260 265

<210> 51  
<211> 267  
<212> PRT  
<213> Homo sapiens

<400> 51

Leu Phe Pro Gln Trp His Leu Pro Ile Lys Ile Ala Ala Ile Ile Ala 1 5 10 15

Ser Leu Thr Phe Leu Tyr Thr Leu Leu Arg Glu Val Ile His Pro Leu 20 25 30

Ala Thr Ser His Gln Gln Tyr Phe Tyr Lys Ile Pro Ile Leu Val Ile 35 40 45

Asn Lys Val Leu Pro Met Val Ser Ile Thr Leu Leu Ala Leu Val Tyr 50 55 60

Leu Pro Gly Val Ile Ala Ala Ile Val Gln Leu His Asn Gly Thr Lys 65 70 75 80

Tyr Lys Lys Phe Pro His Trp Leu Asp Lys Trp Met Leu Thr Arg Lys 85 90 95

Gln Phe Gly Leu Leu Ser Phe Phe Ala Val Leu His Ala Ile Tyr 100 105 110

Ser Leu Ser Tyr Pro Met Arg Ser Tyr Arg Tyr Lys Leu Leu Asn 115 120 125

Trp Ala Tyr Gln Gln Val Gln Gln Asn Lys Glu Asp Ala Trp Ile Glu 130 135 140

His Asp Val Trp Arg Met Glu Ile Tyr Val Ser Leu Gly Ile Val Gly 145 150 155 160

Leu Ala Ile Leu Ala Leu Leu Ala Val Thr Ser Ile Pro Ser Val Ser 165 170 175

Asp Ser Leu Thr Trp Arg Glu Phe His Tyr Ile Gln Ser Lys Leu Gly 180 185 190

Ile Val Ser Leu Leu Leu Gly Thr Ile His Ala Leu Ile Phe Ala Trp 195 200 205

Asn Lys Trp Ile Asp Ile Lys Gln Phe Val Trp Tyr Thr Pro Thr 210 215 220

Phe Met Ile Ala Val Phe Leu Pro Ile Val Val Leu Ile Phe Lys Ser 225 230 235 240

Ile Leu Phe Leu Pro Cys Leu Arg Lys Lys Ile Leu Lys Ile Arg His 250 255

85

245 250 255

Gly Trp Glu Asp Val Thr Lys Ile Asn Lys Thr  
260 265

<210> 52  
<211> 1349  
<212> PRT  
<213> Rattus norvegicus  
<400> 52

Met Lys Ser Ser Arg Thr Val Thr Leu Tyr Phe Val Leu Ile Val Ile  
1 5 10 15

Cys Ser Ser Glu Ala Thr Trp Ser Arg Pro Ala Glu Pro Ile Val His  
20 25 30

Pro Leu Ile Leu Gln Glu His Glu Leu Ala Gly Glu Glu Leu Leu Arg  
35 40 45

Pro Lys Arg Ala Val Ala Val Gly Gly Pro Val Ala Glu Glu Tyr Thr  
50 55 60

Val Asp Val Glu Ile Ser Phe Glu Asn Val Ser Phe Leu Glu Ser Ile  
65 70 75 80

Arg Ala His Leu Asn Ser Leu Arg Phe Pro Val Gln Gly Asn Gly Thr  
85 90 95

Asp Ile Leu Ser Met Ala Met Thr Thr Val Cys Thr Pro Thr Gly Asn  
100 105 110

Asp Leu Leu Cys Phe Cys Glu Lys Gly Tyr Gln Trp Pro Glu Glu Arg  
115 120 125

Cys Leu Ser Ser Leu Thr Cys Gln Glu His Asp Ser Ala Leu Pro Gly  
130 135 140

Arg Tyr Cys Asn Cys Leu Lys Glu Leu Pro Pro Gln Gly Pro Phe Cys  
145 150 155 160

Gln Leu Pro Glu Thr Tyr Ile Thr Leu Lys Ile Lys Val Arg Leu Asn  
165 170 175

Ile Gly Phe Gln Glu Asp Leu Glu Asn Thr Ser Ser Ala Leu Tyr Arg  
180 185 190

Ser Tyr Lys Thr Asp Leu Glu Arg Ala Phe Arg Ala Gly Tyr Arg Thr  
195 200 205

Leu Pro Gly Phe Arg Ser Val Thr Val Thr Gln Phe Thr Lys Gly Ser  
210 215 220

Val Val Val Asp Tyr Ile Val Glu Val Ala Ser Ala Pro Leu Pro Gly  
225 230 235 240

Ser Ile His Lys Lys Ala Asn Glu Gln Val Ile Gln Asn Leu Asn Gln Thr  
245 250 255

Tyr Lys Met Asp Tyr Asn Ser Phe Gln Gly Thr Pro Ser Asn Glu Thr  
260 265 270

Lys Phe Thr Val Thr Pro Glu Phe Ile Phe Glu Gly Asp Asn Val Thr  
275 280 285

Leu Glu Cys Glu Ser Glu Phe Val Ser Ser Asn Thr Ser Trp Phe Tyr  
290 295 300

Gly Glu Lys Arg Ser Asp Ile Gln Asn Ser Asp Lys Phe Ser Ile His  
305 310 315 320

Thr Ser Ile Ile Asn Asn Ile Ser Leu Val Thr Arg Leu Thr Ile Phe  
325 330 335

Asn Phe Thr Gln His Asp Ala Gly Leu Tyr Gly Cys Asn Val Thr Leu  
340 345 350

Asp Ile Phe Glu Tyr Gly Thr Val Arg Lys Leu Asp Val Thr Pro Ile  
355 360 365

Arg Ile Leu Ala Lys Glu Glu Arg Lys Val Val Cys Asp Asn Asn Pro  
370 375 380

Ile Ser Leu Asn Cys Cys Ser Glu Asn Ile Ala Asn Trp Ser Arg Ile  
385 390 395 400

Glu Trp Lys Gln Glu Gly Lys Ile Asn Ile Glu Gly Thr Pro Glu Thr  
405 410 415

Asp Leu Glu Ser Ser Cys Ser Thr Tyr Thr Leu Lys Ala Asp Gly Thr  
420 425 430

Gln Cys Pro Ser Gly Ser Ser Gly Thr Thr Val Ile Tyr Thr Cys Glu  
435 440 445

Phe Val Ser Val Tyr Gly Ala Lys Gly Ser Lys Asn Ile Ala Val Thr  
450 455 460

Phe Thr Ser Val Ala Asn Leu Thr Ile Thr Pro Asp Pro Ile Ser Val  
465 470 475 480

Ser Glu Gly Gln Ser Phe Ser Ile Thr Cys Leu Ser Asp Val Ser Ser  
485 490 495

Phe Asp Glu Val Tyr Trp Asn Thr Ser Ala Gly Ile Lys Ile His Pro  
500 505 510

Arg Phe Tyr Thr Met Arg Arg Tyr Arg Asp Gly Ala Glu Ser Val Leu  
515 520 525

Thr Val Lys Thr Ser Thr Arg Glu Trp Asn Gly Thr Tyr His Cys Ile  
530 535 540

Phe Arg Tyr Lys Asn Ser Tyr Ser Ile Ala Thr Lys Asp Val Thr Val  
545 550 555 560

His Pro Leu Pro Leu Glu Ser Asp Ile Met Met Asp Pro Leu Glu Ala  
565 570 575

Ser Gly Leu Cys Thr Ser Ser His Gln Phe Lys Cys Cys Ile Glu Glu  
580 585 590

Asn Asp Gly Glu Glu Tyr Ile Val Thr Phe His Val Asp Ser Ser Ser  
595 600 605



WO 01/K3454 PCT/US00/34983

Ann His Arg Ser Gly Gly Lys Pro Gln Cys Val Phe Trp Asn Phe Ser 945 950 955 960

Leu Ala Asn Asn Thr Gly Gly Trp Asp Ser Ser Gly Cys Thr Val Glu 965 970 975

Asp Asp Gly Arg Asp Asn Arg Asp Arg Val Phe Cys Lys Cys Asn His 980 985 990

Leu Thr Ser Phe Ser Ile Leu Met Ser Pro Asp Ser Pro Asp Pro Gly 995 1000 1005

Ser Leu Leu Lys Ile Leu Leu Asp Ile Ile Ser Tyr Ile Gly Leu 1010 1015 1020

Gly Phe Ser Ile Val Ser Leu Ala Ala Cys Leu Val Val Glu Ala 1025 1030 1035

Met Val Trp Lys Ser Val Thr Lys Asn Arg Thr Ser Tyr Met Arg 1040 1045 1050

His Ile Cys Ile Val Asn Ile Ala Leu Cys Leu Leu Ile Ala Asp 1055 1060 1065

Ile Trp Phe Ile Val Ala Gly Ala Ile His Asp Gly His Tyr Pro 1070 1075 1080

Leu Asn Glu Thr Ala Cys Val Ala Ala Thr Phe Phe Ile His Phe 1085 1090 1095

Phe Tyr Leu Ser Val Phe Phe Trp Met Leu Thr Leu Gly Leu Met 1100 1105 1110

Leu Phe Tyr Arg Leu Ile Phe Ile Leu His Asp Ala Ser Lys Ser 1115 1120 1125

Thr Gln Lys Ala Ile Ala Phe Ser Leu Gly Tyr Gly Cys Pro Leu 1130 1135 1140

Ile Ile Ser Ser Ile Thr Val Gly Val Thr Gln Pro Gln Glu Val 1145 1150 1155

Tyr Met Arg Lys Asn Ala Cys Trp Leu Asn Trp Glu Asp Thr Arg 1160 1165 1170

Ala Leu Leu Ala Phe Ala Ile Pro Ala Leu Ile Ile Val Val Val 1175 1180 1185

Asn Val Ser Ile Thr Val Val Val Ile Thr Lys Ile Leu Arg Pro 1190 1195 1200

Ser Val Gly Asp Lys Pro Gly Lys Gln Glu Lys Ser Ser Leu Phe 1205 1210 1215

Gln Ile Ser Lys Ser Ile Gly Val Leu Thr Pro Leu Leu Gly Leu 1220 1225 1230

Thr Trp Gly Phe Gly Leu Ala Thr Val Ile Gln Gly Ser Asn Ala 1235 1240 1245

Val Phe His Ile Ile Phe Thr Leu Leu Asn Ala Phe Gln Gly Leu 1250 1255 1260

WO 01/K3454 PCT/US00/34983

Phe Pro Ala Glu Arg Glu Val Ile Gly Lys Gln Ala Cys Tyr Thr Tyr 610 615 620

Ser Leu Pro Gly Lys Leu Pro Ser Arg Cys Pro Lys Asp Ile Asp Val 625 630 635 640

Phe Cys His Phe Thr Asn Ala Ala Asn Ser Ser Val Arg Ser Pro Ser 645 650 655

Met Lys Leu Thr Leu Val Pro Gly Lys Asn Ile Thr Cys Gln Asp Pro 660 665 670

Ile Ile Gly Ile Gly Glu Pro Gly Lys Val Ile Gln Lys Leu Cys Gln 675 680 685

Phe Ala Gly Val Ser Arg Ser Pro Gly Gln Thr Ile Gly Gly Thr Val 690 695 700

Thr Tyr Lys Cys Val Gly Ser Gln Trp Lys Glu Glu Thr Arg Ala Cys 705 710 715 720

Ile Ser Ala Pro Ile Asn Gly Leu Leu Gln Leu Ala Lys Ala Leu Ile 725 730 735

Lys Ser Pro Ser Gln Asp Gln Lys Leu Pro Lys Tyr Leu Arg Asp Leu 740 745 750

Ser Val Ser Thr Gly Lys Glu Glu Gln Asp Ile Arg Ser Ser Pro Gly 755 760 765

Ser Leu Gly Ala Ile Ile Ser Ile Leu Asp Leu Leu Ser Thr Val Pro 770 775 780

Thr Gln Val Asn Ser Glu Met Met Arg Asp Ile Leu Ala Thr Ile Asn 785 790 795 800

Val Ile Leu Asp Lys Ser Thr Leu Asn Ser Thr Glu Lys Leu Leu Gln 805 810 815

Gln Gln Ser Asn Gln Ser Ser Gln Phe Leu Gln Ser Val Glu Arg Phe 820 825 830

Ser Lys Ala Leu Glu Leu Gly Asp Ser Thr Pro Phe Leu Phe His 835 840 845

Pro Asn Val Gln Met Lys Ser Met Val Ile Lys Arg Gly His Ala Gln 850 855 860

Met Tyr Gln Gln Lys Phe Val Phe Thr Asp Ser Asp Leu Trp Gly Asp 865 870 875 880

Val Ala Ile Asp Glu Cys Gln Leu Gly Ser Leu Gln Pro Asp Ser Ser 885 890 895

Ile Val Thr Val Ala Phe Pro Thr Leu Lys Ala Ile Leu Ala Gln Asp 900 905 910

Gly Gln Arg Lys Thr Pro Ser Asn Ser Leu Val Met Thr Thr Thr Val 915 920 925

Ser His Asn Ile Val Lys Pro Phe Arg Ile Ser Met Thr Phe Lys Asn 930 935 940

Leu Asn Ile Met Val Asp Pro Leu Glu Ala Thr Val Ser Cys Ser Gly  
210 215 220  
Ser His His Ile Lys Cys Ile Glu Glu Asp Gly Asp Tyr Lys Val  
225 230 235 240  
Thr Phe His Met Gly Ser Ser Ser Leu Pro Ala Ala Lys Glu Val Asn  
245 250 255  
Lys Lys Gln Val Cys Tyr Lys His Asn Phe Asn Ala Ser Ser Val Ser  
260 265 270  
Trp Cys Ser Lys Thr Val Asp Val Cys Cys His Phe Thr Asn Ala Ala  
275 280 285  
Asn Asn Ser Val Trp Ser Pro Ser Met Lys Leu Asn Leu Val Pro Gly  
290 295 300  
Glu Asn Ile Thr Cys Gln Asp Pro Val Ile Gly Val Gly Glu Pro Gly  
305 310 315 320  
Lys Val Ile Gln Lys Leu Cys Arg Phe Ser Asn Val Pro Ser Ser Pro  
325 330 335  
Glu Ser Pro Ile Gly Gly Thr Ile Thr Tyr Lys Cys Val Gly Ser Gln  
340 345 350  
Trp Glu Glu Lys Arg Asn Asp Cys Ile Ser Ala Pro Ile Asn Ser Leu  
355 360 365  
Leu Gln Met Ala Lys Ala Leu Ile Lys Ser Pro Ser Gln Asp Glu Met  
370 375 380  
Leu Pro Thr Tyr Leu Lys Asp Leu Ser Ile Ser Ile Asp Lys Ala Glu  
385 390 395 400  
His Glu Ile Ser Ser Ser Pro Gly Ser Leu Gly Ala Ile Ile Asn Ile  
405 410 415  
Leu Asp Leu Leu Ser Thr Val Pro Thr Gln Val Asn Ser Glu Met Met  
420 425 430  
Thr His Val Leu Ser Thr Val Asn Val Ile Leu Gly Lys Pro Val Leu  
435 440 445  
Asn Thr Trp Lys Val Leu Gln Gln Gln Trp Thr Asn Gln Ser Ser Gln  
450 455 460  
Leu Leu His Ser Val Glu Arg Phe Ser Gln Ala Leu Gln Ser Gly Asp  
465 470 475 480  
Ser Pro Pro Leu Ser Phe Ser Gln Thr Asn Val Gln Met Ser Ser Thr  
485 490 495  
Val Ile Lys Ser Ser His Pro Glu Thr Tyr Gln Gln Arg Phe Val Phe  
500 505 510  
Pro Tyr Phe Asp Leu Trp Gly Asn Val Val Ile Asp Lys Ser Tyr Leu  
515 520 525  
Glu Asn Leu Gln Ser Asp Ser Ser Ile Val Thr Met Ala Phe Pro Thr  
530 535 540

phe Ile Leu Leu Phe Gly Cys Leu Trp Asp Gln Lys Val Gln Glu  
1265 1270 1275  
Ala Leu Leu His Lys Phe Ser Leu Ser Arg Trp Ser Ser Gln His  
1280 1285 1290  
Ser Lys Ser Thr Ser Leu Gly Ser Ser Thr Pro Val Phe Ser Met  
1295 1300 1305  
Ser Ser Pro Ile Ser Arg Arg Phe Asn Asn Leu Phe Gly Lys Thr  
1310 1315 1320  
Gly Thr Tyr Asn Val Ser Thr Pro Glu Thr Thr Ser Ser Ser Val  
1325 1330 1335  
Glu Asn Ser Ser Ser Ala Tyr Ser Leu Leu Asn  
1340 1345  
<210> 53  
<211> 986  
<212> PRT  
<213> Homo sapiens  
<400> 53  
Cys Lys Lys Lys Ile Asp Val Met Pro Ile Gln Ile Leu Ala Asn Glu  
1 5 10 15  
Glu Met Lys Val Met Cys Asp Asn Asn Pro Val Ser Leu Asn Cys Cys  
20 25 30  
Ser Gln Gly Asn Val Asn Trp Ser Lys Val Glu Trp Lys Gln Glu Gly  
35 40 45  
Lys Ile Asn Ile Pro Gly Thr Pro Glu Thr Asp Ile Asp Ser Ser Cys  
50 55 60  
Ser Arg Tyr Thr Leu Lys Ala Asp Gly Thr Gln Cys Pro Ser Gly Ser  
65 70 75 80  
Ser Gly Thr Thr Val Ile Tyr Thr Cys Glu Phe Ile Ser Ala Tyr Gly  
85 90 95  
Ala Arg Gly Ser Ala Asn Ile Lys Val Thr Phe Ile Ser Val Ala Asn  
100 105 110  
Leu Thr Ile Thr Pro Asp Pro Ile Ser Val Ser Glu Gly Gln Asn Phe  
115 120 125  
Ser Ile Lys Cys Ile Ser Asp Val Ser Asn Tyr Asp Glu Val Tyr Trp  
130 135 140  
Asn Thr Ser Ala Gly Ile Lys Ile Tyr Gln Arg Phe Tyr Thr Thr Arg  
145 150 155 160  
Arg Tyr Leu Asp Gly Ala Glu Ser Val Leu Thr Val Lys Thr Ser Thr  
165 170 175  
Arg Glu Trp Asn Gly Thr Tyr His Cys Ile Phe Arg Tyr Lys Asn Ser  
180 185 190  
Tyr Ser Ile Ala Thr Lys Asp Val Ile Val His Pro Leu Pro Leu Lys  
195 200 205

Leu Gln Ala Ile Leu Ala Gln Asp Ile Gln Glu Asn Asn Phe Ala Glu 545 550 555 560  
 Ser Leu Val Met Thr Thr Thr Val Ser His Asn Thr Thr Met Pro Phe 565 570 575  
 Arg Ile Ser Met Thr Phe Lys Asn Asn Ser Pro Ser Gly Gly Glu Thr 580 585 590 595  
 Lys Cys Val Phe Thr Asn Phe Arg Leu Ala Asn Asn Thr Gly Gly Trp 595 600 605 610  
 Asp Ser Ser Gly Cys Tyr Val Glu Glu Gly Asp Gly Asp Asn Val Thr 610 615 620  
 Cys Ile Cys Asp His Leu Thr Ser Phe Ser Ile Leu Met Ser Pro Asp 625 630 635 640  
 Ser Pro Asp Pro Ser Ser Leu Leu Gly Ile Leu Leu Asp Ile Ile Ser 645 650 655 660  
 Tyr Val Gly Val Gly Phe Ser Ile Leu Ser Leu Ala Ala Cys Leu Val 660 665 670  
 Val Glu Ala Val Val Trp Lys Ser Val Thr Lys Asn Arg Thr Ser Tyr 675 680 685  
 Met Arg His Thr Cys Ile Val Asn Ile Ala Ala Ser Leu Leu Val Ala 690 695 700  
 Asn Thr Trp Phe Ile Val Val Ala Ala Ile Gln Asp Asn Arg Tyr Ile 705 710 715 720  
 Leu Cys Lys Thr Ala Cys Val Ala Ala Thr Phe Phe Ile His Phe Phe 725 730 735  
 Tyr Leu Ser Val Phe Phe Thr Met Leu Thr Leu Gly Leu Met Leu Phe 740 745 750  
 Tyr Arg Leu Val Phe Ile Leu His Glu Thr Ser Arg Ser Thr Gln Lys 755 760 765  
 Ala Ile Ala Phe Cys Leu Gly Tyr Gly Cys Pro Leu Ala Ile Ser Val 770 775 780  
 Ile Thr Leu Gly Ala Thr Gln Pro Arg Glu Val Tyr Thr Arg Lys Asn 785 790 795 800  
 Val Cys Trp Leu Asn Trp Glu Asp Thr Lys Ala Leu Leu Ala Phe Ala 805 810 815  
 Ile Pro Ala Leu Ile Ile Val Val Val Asn Ile Thr Ile Thr Ile Val 820 825 830  
 Val Ile Thr Lys Ile Leu Arg Pro Ser Ile Gly Asp Lys Pro Cys Lys 835 840 845  
 Gln Glu Lys Ser Ser Leu Phe Gln Ile Ser Lys Ser Ile Gly Val Leu 850 855 860  
 Thr Pro Leu Leu Gly Leu Thr Trp Gly Phe Gly Leu Thr Thr Val Phe 865 870 875 880

Pro Gly Thr Asn Leu Val Phe His Ile Ile Phe Ala Ile Leu Asn Val 885 890 895  
 Phe Gln Gly Leu Phe Ile Leu Leu Phe Gly Cys Leu Trp Asp Leu Lys 900 905 910  
 Val Gln Glu Ala Leu Leu Asn Lys Phe Ser Leu Ser Arg Trp Ser Ser 915 920 925 930  
 Gln His Ser Lys Ser Thr Ser Leu Gly Ser Ser Thr Pro Val Phe Ser 930 935 940  
 Met Ser Ser Pro Ile Ser Arg Arg Phe Asn Asn Leu Phe Gly Lys Thr 945 950 955 960  
 Gly Thr Tyr Asn Val Ser Thr Pro Glu Ala Thr Ser Ser Ser Leu Glu 965 970 975  
 Asn Ser Ser Ser Ala Ser Ser Leu Asn 980 985  
 <210> 54  
 <211> 322  
 <212> PRT  
 <213> Mus musculus  
 <400> 54  
 Leu Ser Pro Ala Val Pro Pro Tyr Val Lys Leu Gly Leu Thr Ala Val 1 5 10 15  
 Tyr Thr Val Phe Tyr Ala Leu Leu Phe Val Phe Ile Tyr Ala Gln Leu 20 25 30  
 Trp Leu Val Leu Arg Tyr Arg His Lys Arg Leu Ser Tyr Gln Ser Val 35 40 45  
 Phe Leu Phe Leu Cys Leu Phe Thr Ala Ser Leu Arg Thr Val Leu Phe 50 55 60  
 Ser Phe Tyr Phe Arg Asp Phe Val Ala Ala Asn Ser Phe Ser Pro Phe 65 70 75 80  
 Val Phe Trp Leu Leu Tyr Cys Phe Pro Val Cys Leu Gln Phe Phe Thr 85 90 95  
 Leu Thr Leu Met Asn Leu Tyr Phe Thr Gln Val Ile Phe Lys Ala Lys 100 105 110  
 Ser Lys Tyr Ser Pro Glu Leu Leu Lys Tyr Arg Leu Pro Leu Tyr Leu 115 120 125  
 Ala Ser Leu Phe Ile Ser Leu Val Phe Leu Leu Val Asn Leu Thr Cys 130 135 140  
 Ala Val Leu Val Lys Thr Gly Asp Trp Asp Arg Lys Val Ile Val Ser 145 150 155 160  
 Val Arg Val Ala Ile Asn Asp Thr Leu Phe Val Leu Cys Ala Ile Ser 165 170 175  
 Leu Ser Ile Cys Leu Tyr Lys Ile Ser Lys Met Ser Leu Ala Asn Ile 180 185 190

WO 01/31454 PCT/US00/34983

Ser Leu Arg His Gly Ile Gly His Ile Glu Ala Ile Phe Leu Lys Trp  
145 150 155 160

His Val Pro Pro Glu Leu Val Arg Val Leu Ser Ala Met Leu Phe Leu  
165 170 175

Leu Ile Gly Cys Leu Leu Phe Val Leu Thr Pro Thr Phe Val Phe Cys  
180 185 190

Tyr Met Glu Asp Trp Ser Lys Leu Glu Ala Ile Tyr Phe Val Ile Val  
195 200 205

Thr Leu Thr Thr Val Gly Phe Gly Asp Tyr Val Ala Gly Ala Asp Pro  
210 215 220

Arg Gln Asp Ser Pro Ala Tyr Gln Pro Leu Val Trp Phe Trp Ile Leu  
225 230 235 240

Leu Gly Leu Ala Tyr Phe Ala Ser Val Leu Thr Thr Ile Gly Asn Trp  
245 250 255

Leu Arg Val Val Ser Arg Arg Thr Arg Ala Glu Met Gly Gly Leu Thr  
260 265 270

Ala Gln Ala Ala Ser Trp Thr Gly Thr Val Thr Ala Arg Val Thr Gln  
275 280 285

Arg Ala Gly Pro Ala Ala Pro Pro Glu Lys Glu Gln Pro Leu Leu  
290 295 300

Pro Pro Pro Pro Cys Pro Ala Gln Pro Leu Gly Arg Pro Arg Ser Pro  
305 310 315 320

Ser Pro Pro Glu Lys Ala Gln Pro Pro Ser Pro Pro Thr Ala Ser Ala  
325 330 335

Leu Asp Tyr Pro Ser Glu Asn Leu Ala Phe Ile Asp Glu Ser Ser Asp  
340 345 350

Thr Gln Ser Glu Arg Gly Cys Pro Leu Pro Arg Ala Pro Arg Gly Arg  
355 360 365

Arg Arg Pro Asn Pro Pro Arg Lys Pro Val Arg Pro Arg Gly Pro Gly  
370 375 380

Arg Pro Arg Asp Lys Gly Val Pro  
385 390

<210> 56  
<211> 166  
<212> PRT  
<213> Homo sapiens

<400> 56

Met Ser Tyr Asp Arg Tyr Met Ala Ile Cys His Pro Leu Gln Tyr Ser  
1 5 10 15

Val Ile Met Arg Trp Gly Val Cys Thr Val Leu Ala Val Thr Ser Trp  
20 25 30

Ala Cys Gly Ser Leu Leu Ala Leu Val His Val Val Leu Ile Leu Arg  
35 40 45

95

WO 01/31454 PCT/US00/34983

Tyr Leu Glu Ser Lys Gly Ser Ser Val Cys Gln Val Thr Ala Ile Gly  
180 185 190 195 200 205

Val Thr Val Ile Leu Leu Tyr Ala Ser Arg Ala Cys Tyr Asn Leu Phe  
210 215 220

Ile Leu Ser Phe Ser Gln Ile Lys Asn Val His Ser Phe Asp Tyr Asp  
225 230 235 240

Trp Tyr Asn Val Ser Asp Gln Ala Asp Leu Lys Ser Gln Leu Gly Asp  
245 250 255

Ala Gly Tyr Val Val Phe Gly Val Val Leu Phe Val Trp Glu Leu Leu  
260 265 270

Pro Thr Thr Leu Val Val Tyr Phe Phe Arg Val Arg Asn Pro Thr Lys  
275 280 285

Asp Leu Thr Asn Pro Gly Met Val Pro Ser His Gly Phe Ser Pro Arg  
290 295 300

Ser Tyr Phe Phe Asp Asn Pro Arg Arg Tyr Asp Ser Asp Asp Leu  
305 310 315 320

Ala Trp

<210> 55  
<211> 392  
<212> PRT  
<213> Homo sapiens

<400> 55

Met Arg Ser Thr Thr Leu Leu Ala Leu Leu Ala Leu Val Leu Leu Tyr  
1 5 10 15

Leu Val Ser Gly Ala Leu Val Phe Arg Ala Leu Glu Gln Pro His Glu  
20 25 30

Gln Gln Ala Gln Arg Glu Leu Gly Glu Val Arg Glu Lys Phe Leu Arg  
35 40 45

Ala His Pro Cys Val Ser Asp Gln Glu Leu Gly Leu Leu Ile Lys Glu  
50 55 60

Val Ala Asp Ala Leu Gly Gly Ala Asp Pro Glu Thr Asn Ser Thr  
65 70 75 80

Ser Asn Ser Ser His Ser Ala Trp Asp Leu Gly Ser Ala Phe Phe  
85 90 95

Ser Gly Thr Ile Ile Thr Thr Ile Gly Tyr Gly Asn Val Ala Leu Arg  
100 105 110

Thr Asp Ala Gly Arg Leu Phe Cys Ile Phe Tyr Ala Leu Val Gly Ile  
115 120 125

Pro Leu Phe Gly Ile Leu Leu Ala Gly Val Gly Asp Arg Leu Gly Ser  
130 135 140

94

Leu Pro Phe Cys Gly Pro His Glu Ile Asn His Phe Phe Cys Glu Ile  
50 55 60

Leu Ser Val Leu Lys Leu Ala Cys Ala Asp Thr Trp Leu Asn Gln Val  
65 70 75 80

Val Ile Phe Ala Ala Ser Val Phe Ile Leu Val Gly Pro Leu Cys Leu  
85 90 95

Val Leu Val Ser Tyr Ser Arg Ile Leu Ala Ala Ile Leu Arg Ile Gln  
100 105 110

Ser Gly Glu Gly Arg Arg Lys Ala Phe Ser Thr Cys Ser Ser His Leu  
115 120 125

Cys Met Val Gly Leu Phe Phe Gly Ser Ala Ile Val Met Tyr Met Ala  
130 135 140

Pro Lys Ser Arg His Pro Glu Glu Gln Lys Val Leu Ser Leu Phe  
145 150 155 160

Tyr Ser Leu Phe Asn Pro  
165

<210> 57

<211> 171

<212> PRT

<213> Homo sapiens

<400> 57

Met Ser Tyr Asp Arg Tyr Val Ala Ile Cys His Pro Leu Arg Tyr Phe  
1 5 10 15

Ile Ile Met Thr Trp Lys Val Cys Ile Thr Leu Gly Ile Thr Ser Trp  
20 25 30

Thr Cys Gly Ser Leu Leu Ala Met Val His Val Ser Leu Ile Leu Arg  
35 40 45

Leu Pro Phe Cys Gly Pro Arg Glu Ile Asn His Phe Phe Cys Glu Ile  
50 55 60

Leu Ser Val Leu Arg Leu Ala Cys Ala Asp Thr Trp Leu Asn Gln Val  
65 70 75 80

Val Ile Phe Glu Ala Cys Met Phe Ile Leu Val Gly Pro Leu Cys Leu  
85 90 95

Val Leu Val Ser Tyr Ser His Ile Leu Gly Gly Ile Leu Arg Ile Gln  
100 105 110

Ser Gly Glu Gly Arg Arg Lys Ala Phe Ser Thr Cys Ser Ser His Leu  
115 120 125

Cys Val Val Gly Leu Phe Phe Gly Ser Ala Ile Val Met Tyr Met Ala  
130 135 140

Pro Lys Ser Arg His Pro Glu Glu Gln Lys Val Leu Phe Leu Ile  
145 150 155 160

Leu Gln Phe Leu Ser Thr Pro Met Leu Lys Pro  
96

<210> 58

<211> 304

<212> PRT

<213> Homo sapiens

<400> 58

Met Gly Asp Asn Ile Thr Ser Ile Arg Glu Phe Leu Leu Leu Gly Phe  
1 5 10 15

Pro Val Gly Pro Arg Ile Gln Met Leu Leu Phe Gly Leu Phe Ser Leu  
20 25 30

Phe Tyr Val Phe Thr Leu Leu Gly Asn Gly Thr Ile Leu Gly Leu Ile  
35 40 45

Ser Leu Asp Ser Arg Leu His Ala Pro Met Tyr Phe Phe Leu Ser His  
50 55 60

Leu Ala Val Val Asp Ile Ala Tyr Ala Cys Asn Thr Val Pro Arg Met  
65 70 75 80

Leu Val Asn Leu Leu His Pro Ala Lys Pro Ile Ser Phe Ala Gly Arg  
85 90 95

Met Met Gln Thr Phe Leu Phe Ser Thr Phe Ala Val Thr Glu Cys Leu  
100 105 110

Leu Leu Val Val Met Ser Tyr Asp Leu Tyr Val Ala Ile Cys His Pro  
115 120 125

Leu Arg Tyr Leu Ala Ile Met Thr Trp Arg Val Cys Ile Thr Leu Ala  
130 135 140

Val Thr Ser Trp Thr Thr Gly Val Leu Leu Ser Leu Ile His Leu Val  
145 150 155 160

Leu Leu Leu Pro Leu Pro Phe Cys Arg Pro Gln Lys Ile Tyr His Phe  
165 170 175

Phe Cys Glu Ile Leu Ala Val Leu Lys Leu Ala Cys Ala Asp Thr His  
180 185 190

Ile Asn Glu Asn Met Val Leu Ala Gly Ala Ile Ser Gly Leu Val Gly  
195 200 205

Pro Leu Ser Thr Ile Val Val Ser Tyr Met Cys Ile Leu Cys Ala Ile  
210 215 220

Leu Gln Ile Gln Ser Arg Glu Val Gln Arg Lys Ala Phe Arg Thr Cys  
225 230 235 240

Phe Ser His Leu Cys Val Ile Gly Leu Val Tyr Gly Thr Ala Ile Ile  
245 250 255

Met Tyr Val Gly Pro Arg Tyr Gly Asn Pro Lys Glu Lys Lys Tyr  
260 265 270

Leu Leu Leu Phe His Ser Leu Phe Asn Pro Met Leu Asn Pro Leu Ile  
275 280 285

Cys Ser Leu Arg Asn Ser Glu Val Lys Asn Thr Leu Lys Arg Val Leu  
290 295 300

**<210> 59**

**<211> 287**

**<212> PRT**

**<213> Homo sapiens**

**<400> 59**

Met Val Lys Asn Gln Thr Met Val Thr Glu Phe Leu Leu Gly Phe  
1 5 10 15

Leu Leu Gly Pro Arg Ile Gln Met Leu Leu Phe Gly Leu Phe Ser Leu  
20 25 30

Phe Tyr Val Phe Thr Leu Leu Gly Asn Gly Thr Ile Leu Gly Leu Ile  
35 40 45

Ser Leu Asp Ser Arg Leu His Thr Pro Met Tyr Phe Phe Leu Ser His  
50 55 60

Leu Ala Val Val Asn Ile Ala Tyr Ala Cys Asn Thr Val Pro Gln Met  
65 70 75 80

Leu Val Asn Leu Leu His Pro Ala Lys Pro Ile Ser Phe Ala Gly Cys  
85 90 95

Met Thr Leu Asp Phe Leu Phe Leu Ser Phe Ala His Thr Glu Cys Leu  
100 105 110

Leu Leu Val Leu Met Ser Tyr Asp Arg Tyr Val Ala Ile Cys His pro  
115 120 125

Leu Arg Tyr Phe Ile Ile Met Thr Trp Lys Val Cys Ile Thr Leu Gly  
130 135 140

Ile Thr Ser Trp Thr Cys Gly Ser Leu Ala Met Val His Val Ser  
145               150               155               160

Leu Ile Leu Arg Leu Pro Phe Cys Gly Pro Arg Glu Ile Asn His Phe  
165 170 175

Phe Cys Glu Ile Leu Ser Val Leu Arg Leu Ala Cys Ala Asp Thr Trp  
180 185 190

Leu Asn Gln Val Val Ile Phe Glu Ala Cys Met Phe Ile Leu Val Gly  
195 200 205

Pro Leu Cys Leu Val Leu Val Ser Tyr Ser His Ile Leu Gly Gly Ile  
210 215 220

Leu Arg Ile Gln Ser Gly Glu Gly Arg Lys Ala Phe Ser Thr Cys

Ser Ser His Leu Cys Val Val Gly Leu Phe Phe Gly Ser Ala Ile Val

Met Tyr Met Ala Pro Lys Ser Arg His Pro Glu Glu Gln Lys Val

Leu Phe Leu Ile Leu Gln Phe Leu Ser Thr Pro Met Leu Lys Pro

98

99

WO 01/51454	PCT/US00/34983
gaa gat gtt acc ctg aac atg aga gtc aga cta aat gta ggc ttt caa Glu Asp Val Thr Leu Asn Met Arg Val Arg Leu Asn Val Gly Phe Gln 170 175 180	820
gaa gac ctg atg aac act tcc tcc gcc ctg tat agg tcc tac aag acc Glu Asp Leu Met Asn Thr Ser Ser Ala Leu Tyr Arg Ser Tyr Lys Thr 185 190 195	868
gac ttg gaa aca gcg ttc cgg aag ggt tac gga att tta caa ggc ttc Asp Leu Glu Thr Ala Phe Arg Lys Gly Tyr Gly Ile Leu Pro Gly Phe 200 205 210 215	916
aag ggc gtg act gtg aca ggg ttc aag tct gga agt gtg gtt gtg aca Lys Gly Val Thr Val Thr Gly Phe Lys Ser Gly Ser Val Val Val Thr 220 225 230	964
tat gaa gtc aag act aca cca tca ctt gag tta ata cat aaa gcc Tyr Glu Val Lys Thr Thr Pro Ser Leu Glu Ile His Lys Ala 235 240 245	1012
aat gaa caa gtt gta cag agc ctg aat cag acc tac aaa atg gac tac Asn Glu Gln Val Val Gln Ser Leu Asn Gln Thr Tyr Lys Met Asp Tyr 250 255 260	1060
aac tcc ttt caa gca gtt act atc aat gaa agc aat ttc ttt gtc aca Asn Ser Phe Gln Ala Val Thr Ile Asn Glu Ser Asn Phe Phe Val Thr 265 270 275	1108
caa gaa atc atc ttt gaa ggg gac aca gtc agt ctg gtg tgt gaa aag Pro Glu Ile Ile Phe Glu Gly Asp Thr Val Ser Leu Val Cys Glu Lys 280 285 290 295	1156
gaa gtt ttg tcc tcc atc gtg tct tgg cgc tat gaa gaa cag cag ttg Glu Val Leu Ser Ser Asn Val Ser Trp Arg Tyr Glu Glu Gln Leu 300 305 310	1204
gaa atc cag aac agc agc aga ttc tgg att tac acc gca ctt ttc aac Glu Ile Gln Asn Ser Ser Arg Phe Ser Ile Tyr Thr Ala Leu Phe Asn 315 320 325	1252
aac atg act tcg gtg tcc aag ctg acc atc cag aac atc cca ggt Asn Met Thr Ser Val Ser Lys Leu Thr Ile His Asn Ile Thr Pro Gly 330 335 340	1300
gat gca ggt gaa tat gtt tgc aaa ctg ata tta gac att ttt gaa tat Asp Ala Gly Glu Tyr Val Cys Lys Leu Ile Leu Asp Ile Phe Glu Tyr 345 350 355	1348
gag tgc aag aag aaa ata gat gtt atg ccc atc caa att ttg gca aat Glu Cys Lys Lys Lys Ile Asp Val Met Pro Ile Gln Ile Leu Ala Asn 360 365 370 375	1396
gaa gaa atg aag gtg atg tgc gac aac aat cct gta tct ttg aac tgc Glu Glu Met Lys Val Met Cys Asp Asn Asn Pro Val Ser Leu Asn Cys 380 385 390	1444
tgc agt cag ggt aat gtt aat tgg agc aaa gta gaa tgg aag cag gaa Cys Ser Gln Gly Asn Val Asn Trp Ser Lys Val Glu Trp Lys Gln Glu 395 400 405	1492
gga aaa ata aat att cca gga acc cct gag aca gac ata gat tct agc Gly Lys Ile Asn Ile Pro Gly Thr Pro Glu Thr Asp Ile Asp Ser Ser 410 415 420	1540
	100
WO 01/51454	PCT/US00/34983
tgc agc aga tac acc ctg aag gct gat gga acc cag tgc cca agc ggg Cys Ser Arg Tyr Thr Leu Lys Ala Asp Gly Thr Gln Cys Pro Ser Gly 425 430 435	1588
tcg tct gga aca aca gtc atc tac act tgt gag ttc atc agt gcc tat Ser Ser Gly Thr Thr Val Ile Tyr Thr Cys Glu Phe Ile Ser Ala Tyr 440 445 450 455	1636
gga gcc aga ggc agt gca aac ata aaa gtg aca ttc atc tct gtg gcc Gly Ala Arg Gly Ser Ala Asn Ile Lys Val Thr Phe Ile Ser Val Ala 460 465 470	1684
aat cta aca ata aco ccg gac cca att tct gtt tct gag gga caa aac Asn Leu Thr Ile Thr Pro Asp Pro Ile Ser Val Ser Glu Gly Gln Asn 475 480 485	1732
ttt tct ata aaa tgc atc agt gat gtg agt aac tat gat gag gtt tat Phe Ser Ile Lys Cys Ile Ser Asp Val Ser Asn Tyr Asp Glu Val Tyr 490 495 500	1780
tgg aac act tct gct gga att aaa ata tac caa aga ttt tat acc acg Trp Asn Thr Ser Ala Gly Ile Lys Ile Tyr Gln Arg Phe Tyr Thr Thr 505 510 515	1828
agg agg tat ctt gat gga gca gaa tca gta ctg aca gtc aag acc tgg Arg Arg Tyr Leu Asp Gly Ala Glu Ser Val Leu Thr Val Lys Thr Ser 520 525 530 535	1876
acc agg gag tgg aat gga acc tat cac tgc ata ttt aga tat aag aat Thr Arg Glu Trp Asn Gly Thr Tyr His Cys Ile Phe Arg Tyr Lys Asn 540 545 550	1924
tca tac agt att gca acc aaa gac gtc att gtt cac cgg ctg cct cta Ser Tyr Ser Ile Ala Thr Lys Asp Val Ile Val His Pro Leu Pro Leu 555 560 565	1972
aag ctg aac atc atg gtt gat cct ttg gaa gct act gtt tca tgc agt Lys Leu Asn Ile Met Val Asp Pro Leu Glu Ala Thr Val Ser Cys Ser 570 575 580	2020
ggt tcc cat cac atc aag tgc tgc ata gag gag gat gga gac tac aaa Gly Ser His His Ile Lys Cys Cys Ile Glu Glu Asp Gly Asp Tyr Lys 585 590 595	2068
ggt act ttc cat atg ggt tcc tca tcc ctt cct gct gca aaa gaa gtt Val Thr Phe His Met Gly Ser Ser Ser Leu Pro Ala Ala Lys Glu Val 600 605 610 615	2116
aac aaa aaa caa gtg tgc tac aaa cac aat ttc aat gca ago tca gtt Asn Lys Lys Gln Val Cys Tyr Lys His Asn Phe Asn Ala Ser Ser Val 620 625 630	2164
tcc tgg tgt tca aaa act gtt gat gtg tgt tgt cac ttt acc aat gct Ser Trp Cys Ser Lys Thr Val Asp Val Cys Cys His Phe Thr Asn Ala 635 640 645	2212
ggt aat aat tca gtt tgg agc cca tct atg aag ctg aat ctg gtt cct Ala Asn Asn Ser Val Trp Ser Pro Ser Met Lys Leu Asn Leu Val Pro 650 655 660	2260
ggg gaa aac atc aca tgc cag gat ccc gta ata ggt gtc gga gag cgg Gly Glu Asn Ile Thr Cys Gln Asp Pro Val Ile Gly Val Gly Glu Pro 665 670 675	2308
	101

WO 01/31454 665 670 675 PCT/US00/31983

999 aaa gtc atc cag aag cta tgc cgg ttc tca aac gtt ccc agc agc  
 gly lys val ile gln lys leu cys arg phe ser asn val pro ser ser  
 680 685 690 695

cct gag agt ccc att ggc ggg acc atc act tac aaa tgt gta ggc tcc  
 pro glu ser pro ile gly gly thr ile thr tyr lys cys val gly ser  
 700 705 710

cag tgg gag gag aag aga aat gac tgc atc tct gcc cca ata aac agt  
 gln trp glu gln lys arg asn asp cys ile ser ala pro ile asn ser  
 715 720 725

ctg ctc cag atg gct aag gct ttg atc aag agc ccc tct cag gat gag  
 leu leu gln met ala lys ala leu ile lys ser pro ser gln asp glu  
 730 735 740

atg ctc cct aca tac ctg aag gat ctt tct att agc ata gac aaa gcg  
 met leu pro thr tyr leu lys asp leu ser ile ser ile asp lys ala  
 745 750 755

gaa cat gaa atc agc tct tct cct ggg agt ctg gga gcc att att aac  
 glu his glu ile ser ser ser pro gly ser leu gly ala ile ile asn  
 760 765 770 775

atc ctt gat ctg ctc tca aca gtt cca acc caa gta aat tca gaa atg  
 ile leu asp leu leu ser thr val pro thr gln val asn ser glu met  
 780 785 790

atg acg cac gtg ctc tct acg gtt aat gtc atc ctt ggc aag ccc gtc  
 met thr his val leu ser thr val asn val ile leu gly lys pro val  
 795 800 805

ttg aac acc tgg aag gtt tta caa cag caa tgg acc aat cag agt tca  
 leu asn thr trp lys val leu gln gln thr trp thr asn gln ser ser  
 810 815 820

cag cta cta cat tca gtg gaa aga ttt tcc caa gca tta cag tca gga  
 gln leu leu his ser val glu arg phe ser gln ala leu gln ser gly  
 825 830 835

gat agc cct cct ttg tcc ttc tcc caa act aat gtg cag atg agc agc  
 asp ser pro pro leu ser phe ser gln thr asn val gln met ser ser  
 840 845 850 855

acg gta atc aag tcc agc cac cca gaa acc tat caa cag agg ttt gtt  
 thr val ile lys ser ser his pro glu thr tyr gln gln arg phe val  
 860 865 870

ttc cca tac ttt gac ctc tgg ggc aat gtg gtc att gac aag agc tac  
 phe pro tyr phe asp leu trp gly asn val val ile asp lys ser tyr  
 875 880 885

cta gaa aac ttg cag tcy gat tcy tct att gtc acc atg gct ttc cca  
 leu glu asn leu gln ser asp ser ser ile val thr met ala phe pro  
 890 895 900

act ctc caa gcc atc ctt gct cag gat atc cag gaa aat aac ttt gca  
 thr leu gln ala ile leu ala gln asp ile gln glu asn asn phe ala  
 905 910 915

gag agc tta gtg atg aca acc act gtc agc cac aat acg act atg cca  
 920 925 930 935

ttc agg att tca atg act ttt aag aac aat agc cct tca ggc ggc gaa  
 phe arg ile ser met thr phe lys asn asn ser pro ser gly gly glu  
 940 945 950

acg aag tgt gtc ttc tgg aac ttc agg ctt gcc aac aac aca ggg ggg  
 thr lys cys val phe thr asp phe arg leu ala asn asn thr gly gly  
 955 960 965

tgg gac agc agt ggg tgc tat gtt gaa gaa ggt gat ggg gac aat gtc  
 trp asp ser ser gly cys tyr val glu glu gly asp gly asp asn val  
 970 975 980

acc tgt atc tgt gac cac cta cta tca ttc ttc atc ctc atg tcc cct  
 thr cys ile cys asp his leu thr ser phe ser ile leu met ser pro  
 985 990 995

gac tcc cca gat cct agt tct ctc ctg gga ata ctc ctg gat att  
 asp ser pro asp pro ser ser leu leu gly ile leu leu asp ile  
 1000 1005 1010

att tct tat gtt ggg gtg ggc ttt tcc atc ttg agc ttg gca gcc  
 ile ser tyr val gly val gly phe ser ile leu ser leu ala ala  
 1015 1020 1025

tgt cta gtt gtg gaa gct gtg gtg tgg aaa tcy gtg acc aag aat  
 cys leu val val glu ala val val trp lys ser val thr lys asn  
 1030 1035 1040

cgg act tct tat atg cgc cac acc tgc ata gtg aat atc gct gcc  
 arg thr ser tyr met arg his thr cys ile val asn ile ala ala  
 1045 1050 1055

tcc ctt ctg gtc gcc aac acc tgg ttc att gtg gtc gct gcc atc  
 ser leu leu val ala asn thr trp phe ile val val ala ala ile  
 1060 1065 1070

cag gac aat cgc tac ata ctc tgc aag aca gcc tgt gtc gct gcc  
 gln asp asn arg tyr ile leu cys lys thr ala cys val ala ala  
 1075 1080 1085

acc ttc ttc atc cac ttc ttc tac ctc agc gtc ttc ttc tgg atg  
 thr phe phe ile his phe phe tyr leu ser val phe phe trp met  
 1090 1095 1100

ctg aca ctg ggc ctc atg ctg ttc tat cgc ctg gtt ttc att ctg  
 leu thr leu gly leu met leu phe tyr arg leu val phe ile leu  
 1105 1110 1115

cat gaa aca agc agg tcc act cag aaa gcc att gcc ttc tgt ctt  
 his glu thr ser arg ser thr gln lys ala ile ala phe cys leu  
 1120 1125 1130

ggc tat ggc tgc cca ctt gcc atc tgc gtc atc acg ctg gga gcc  
 gly tyr gly cys pro leu ala ile ser val ile thr leu gly ala  
 1135 1140 1145

acc cag ccc cgg gaa gtc tat acg agg aag aat gtc tgt tgg ctc  
 thr gln pro arg glu val tyr thr arg lys asn val cys trp leu  
 1150 1155 1160

WO 01/31454 665 670 675 PCT/US00/31983

999 aaa gtc atc cag aag cta tgc cgg ttc tca aac gtt ccc agc agc  
 gly lys val ile gln lys leu cys arg phe ser asn val pro ser ser  
 680 685 690 695

cct gag agt ccc att ggc ggg acc atc act tac aaa tgt gta ggc tcc  
 pro glu ser pro ile gly gly thr ile thr tyr lys cys val gly ser  
 700 705 710

cag tgg gag gag aag aga aat gac tgc atc tct gcc cca ata aac agt  
 gln trp glu gln lys arg asn asp cys ile ser ala pro ile asn ser  
 715 720 725

ctg ctc cag atg gct aag gct ttg atc aag agc ccc tct cag gat gag  
 leu leu gln met ala lys ala leu ile lys ser pro ser gln asp glu  
 730 735 740

atg ctc cct aca tac ctg aag gat ctt tct att agc ata gac aaa gcg  
 met leu pro thr tyr leu lys asp leu ser ile ser ile asp lys ala  
 745 750 755

gaa cat gaa atc agc tct tct cct ggg agt ctg gga gcc att att aac  
 glu his glu ile ser ser ser pro gly ser leu gly ala ile ile asn  
 760 765 770 775

atc ctt gat ctg ctc tca aca gtt cca acc caa gta aat tca gaa atg  
 ile leu asp leu leu ser thr val pro thr gln val asn ser glu met  
 780 785 790

atg acg cac gtg ctc tct acg gtt aat gtc atc ctt ggc aag ccc gtc  
 met thr his val leu ser thr val asn val ile leu gly lys pro val  
 795 800 805

ttg aac acc tgg aag gtt tta caa cag caa tgg acc aat cag agt tca  
 leu asn thr trp lys val leu gln gln thr trp thr asn gln ser ser  
 810 815 820

cag cta cta cat tca gtg gaa aga ttt tcc caa gca tta cag tca gga  
 gln leu leu his ser val glu arg phe ser gln ala leu gln ser gly  
 825 830 835

gat agc cct cct ttg tcc ttc tcc caa act aat gtg cag atg agc agc  
 asp ser pro pro leu ser phe ser gln thr asn val gln met ser ser  
 840 845 850 855

acg gta atc aag tcc agc cac cca gaa acc tat caa cag agg ttt gtt  
 thr val ile lys ser ser his pro glu thr tyr gln gln arg phe val  
 860 865 870

ttc cca tac ttt gac ctc tgg ggc aat gtg gtc att gac aag agc tac  
 phe pro tyr phe asp leu trp gly asn val val ile asp lys ser tyr  
 875 880 885

cta gaa aac ttg cag tcy gat tcy tct att gtc acc atg gct ttc cca  
 leu glu asn leu gln ser asp ser ser ile val thr met ala phe pro  
 890 895 900

act ctc caa gcc atc ctt gct cag gat atc cag gaa aat aac ttt gca  
 thr leu gln ala ile leu ala gln asp ile gln glu asn asn phe ala  
 905 910 915

gag agc tta gtg atg aca acc act gtc agc cac aat acg act atg cca  
 920 925 930 935



aac tgg gag gac acc aag gcc ctg ctg gct ttc gcc atc cca gca 3808  
 Asn Trp Glu Asp Thr Lys Ala Leu Leu Ala Phe Ala Ile Pro Ala 1175  
 1165  
 ctg atc att gtg gtg gtc aac ata acc atc act att gtg gtc atc 3853  
 Leu Ile Ile Val Val Val Asn Ile Thr Ile Thr Ile Val Val Ile 1190  
 1180  
 acc aag atc ctg agg cct tcc att gga gac aag cca tgc aag cag 3898  
 Thr Lys Ile Leu Arg Pro Ser Ile Gly Asp Lys Pro Cys Lys Gln 1205  
 1195  
 gag aag agc agc ctg ttt cag atc agc aag agc att ggg gtc ctc 3943  
 Glu Lys Ser Ser Leu Phe Gln Ile Ser Lys Ser Ile Gly Val Leu 1220  
 1210  
 aca cca ctg tgg ggc ctc act tgg ggt ttt ggt ctc acc act gtc 3988  
 Thr Pro Leu Leu Gly Leu Thr Trp Gly Phe Gly Leu Thr Thr Val 1235  
 1225  
 ttc cca ggg acc aac ctt gtg ttc cat atc ata ttt gcc atc ctc 4033  
 Pro Gly Thr Asn Leu Val Phe His Ile Ile Phe Ala Ile Leu 1250  
 1240  
 aat gtc ttc cag gga tta ttc att tta ctc ttt gga tgc ctc tgg 4078  
 Asn Val Phe Gln Gly Leu Phe Ile Leu Leu Phe Gly Cys Leu Trp 1265  
 1255  
 gat ctg aag gta cag gaa gct ttg ctg aat aag ttt tca ttg tgg 4123  
 Asp Leu Lys Val Gln Glu Ala Leu Leu Asn Lys Phe Ser Leu Ser 1280  
 1270  
 aga tgg tct tca cag cca tca aag tca aca tcc ctg ggt tca tcc 4168  
 Arg Trp Ser Ser Gln His Ser Lys Ser Thr Ser Leu Gly Ser Ser 1295  
 1285  
 aca cct gtg ttt tct atg agt tct cca ata tca agg aga ttt aac 4213  
 Thr Pro Val Phe Ser Met Ser Ser Pro Ile Ser Arg Arg Phe Asn 1310  
 1300  
 aat ttg ttt ggt aaa aca gga acg tat aat gtt tcc acc cca gaa 4258  
 Asn Leu Phe Gly Lys Thr Gly Thr Tyr Asn Val Ser Thr Pro Glu 1325  
 1315  
 gca acc agc tca tcc ctg gaa aac tca tcc agt gct tct tgg ttg 4303  
 Ala Thr Ser Ser Ser Leu Glu Asn Ser Ser Ala Ser Ser Leu 1340  
 1330  
 ctc aac taa gaacaggata atccaaacta cgtgacctoc cggggacagt. 4352  
 Leu Asn  
 1345  
 ggctgtgctt ttaaaaagag atgcttgcaa agcaatgggg aacgtgttct cggggcaggt 4412  
 ttccggggagc agatgccaaa aagacttttt catagagaag aggttttttt ttgtaaagac 4472  
 agaataaaaa taattgttat gttctgtttt gttccctccc cctccccctt gtgtgatacc 4532  
 acatgtgtat agtatttaag tgaactcaaa gccctcaagg cccaactttt ctgtctatat 4592  
 tgaatatag aatttogaag agacatttbc actttttaca cattggggcgc aaagataagc 4652  
 ttgtattaaa gtactaagta aaaggctacc taggaataac ttcaagtgaat tctaagaagg 4712  
 104

aaggaaggaa gaaagggaagg aaagaaggga gggaaacagg gagaaggga aaaagaagaa 4772  
 aaagaagaag atgaataag gaacaataa agacaacaa catlaaggga catattgtaa 4832  
 gatttccatg ttaagtatct aataataca ctcaagtgcaa cattgagaat ttttttttaa 4892  
 tggctcaaaa atggaactg aaagcaagtc atggggaatg aatactttgg gcagtatcct 4952  
 cctcatgtct tcttagctaa gaggaggaaa aaaaggctga aaaaataggg aggaattccc 5012  
 ttcatcagaa cgacttcag tggataacaa tattataag aaatgaatgg aaggaaatat 5072  
 gatcctctg agactaactt tgaatgttaa ggtttgaact aagtgaatgt atctgcagag 5132  
 gaagtattac aaagatatgt cattagatcc aagtctgat taaattttta tagtttatca 5192  
 gaaaagcctt atattttagt ttgtccaca tttgaaagc aaaaaatata tatttgatat 5252  
 accctcaat tgcacaattt gatagtgc actgaagaca gacctgtgca tatatttaat 5312  
 ggcttcagc aggtacttct ctgtgcattta tagaatagat ttaataatc ttatagcatt 5372  
 gtatattat atgtctgtg tcaactgtat tattattgtg gatactggcc ctgtgtgtgt 5432  
 tgcataagtc cctatgtatt ctctgtttcc atctttaagt tccagagcca atacaatca 5492  
 agagttttgc atggtctaaa ttgtgttat tccaaccagc tggaaagctc ctggaaagaa 5552  
 attttacatt cgggtgttct gtgtctctaa tgaacttga cctgtgtgaa caaatggcag 5612  
 agcctttccc aaggatttga ttgtttgtga attatctgca tgtgtgcttt tttttgtgt 5672  
 gtatttcatt aaaaaatata aatatttatg aaaaaaaaa aa 5714

<210> 61  
 <211> 1346  
 <212> PRT  
 <213> Homo sapiens  
 <400> 61  
 Met Lys Ser Pro Arg Arg Thr Thr Leu Cys Leu Met Phe Ile Val Ile  
 1 5 10 15  
 Tyr Ser Ser Lys Ala Ala Leu Asn Trp Asn Tyr Glu Ser Thr Thr Ile His  
 20 25 30  
 Pro Leu Ser Leu His Glu His Glu Pro Ala Gly Glu Glu Ala Leu Arg  
 35 40 45  
 Gln Lys Arg Ala Val Ala Thr Lys Ser Pro Thr Ala Glu Glu Tyr Thr  
 50 55 60  
 Val Asn Ile Glu Ile Ser Phe Phe Glu Asn Ala Ser Phe Leu Asp Pro Ile  
 65 70 75 80

Lys Ala Tyr Leu Asn Ser Leu Ser Phe Pro Ile His Gly Asn Asn Thr  
85 90 95

Asp Gln Ile Thr Asp Ile Leu Ser Ile Asn Val Thr Thr Val Cys Arg  
100 105 110

Pro Ala Gly Asn Glu Ile Trp Cys Ser Cys Glu Thr Gly Tyr Gly Trp  
115 120 125

Pro Arg Glu Arg Cys Leu His Asn Leu Ile Cys Gln Glu Arg Asp Val  
130 135 140

Phe Leu Pro Gly His His Cys Ser Cys Leu Lys Glu Leu Pro Pro Asn  
145 150 155 160

Gly Pro Phe Cys Leu Leu Gln Glu Asp Val Thr Leu Asn Met Arg Val  
165 170 175

Arg Leu Asn Val Gly Phe Gln Glu Asp Leu Met Asn Thr Ser Ser Ala  
180 185 190

Leu Tyr Arg Ser Tyr Lys Thr Asp Leu Glu Thr Ala Phe Arg Lys Gly  
195 200 205

Tyr Gly Ile Leu Pro Gly Phe Lys Gly Val Thr Val Thr Gly Phe Lys  
210 215 220

Ser Gly Ser Val Val Thr Tyr Glu Val Lys Thr Thr Pro Pro Ser  
225 230 235 240

Leu Glu Leu Ile His Lys Ala Asn Glu Gln Val Val Gln Ser Leu Asn  
245 250 255

Gln Thr Tyr Lys Met Asp Tyr Asn Ser Phe Gln Ala Val Thr Ile Asn  
260 265 270

Glu Ser Asn Phe Phe Val Thr Pro Glu Ile Ile Phe Glu Gly Asp Thr  
275 280 285

Val Ser Leu Val Cys Glu Lys Glu Val Leu Ser Ser Asn Val Ser Trp  
290 295 300

Arg Tyr Glu Glu Gln Gln Leu Glu Ile Gln Asn Ser Ser Arg Phe Ser  
305 310 315 320

Ile Tyr Thr Ala Leu Phe Asn Asn Met Thr Ser Val Ser Lys Leu Thr  
325 330 335

Ile His Asn Ile Thr Pro Gly Asp Ala Gly Glu Tyr Val Cys Lys Leu  
340 345 350

Ile Leu Asp Ile Phe Glu Tyr Glu Cys Lys Lys Ile Asp Val Met  
355 360 365

Pro Ile Gln Ile Leu Ala Asn Glu Glu Met Lys Val Met Cys Asp Asn  
370 375 380

Asn Pro Val Ser Leu Asn Cys Cys Ser Gln Gly Asn Val Asn Trp Ser  
385 390 395 400

Lys Val Glu Trp Lys Gln Glu Gly Lys Ile Asn Ile Pro Gly Thr Pro  
405 410 415

Glu Thr Asp Ile Asp Ser Ser Cys Ser Arg Tyr Thr Leu Lys Ala Asp  
420 425 430

Gly Thr Gln Cys Pro Ser Gly Ser Ser Gly Thr Thr Val Ile Tyr Thr  
435 440 445

Cys Glu Phe Ile Ser Ala Tyr Gly Ala Arg Gly Ser Ala Asn Ile Lys  
450 455 460

Val Thr Phe Ile Ser Val Ala Asn Leu Thr Ile Thr Pro Asp Pro Ile  
465 470 475 480

Ser Val Ser Glu Gly Gln Asn Phe Ser Ile Lys Cys Ile Ser Asp Val  
485 490 495

Ser Asn Tyr Asp Glu Val Tyr Trp Asn Thr Ser Ala Gly Ile Lys Ile  
500 505 510

Tyr Gln Arg Phe Tyr Thr Thr Arg Arg Tyr Leu Asp Gly Ala Glu Ser  
515 520 525

Val Leu Thr Val Lys Thr Ser Thr Arg Glu Trp Asn Gly Thr Tyr His  
530 535 540

Cys Ile Phe Arg Tyr Lys Asn Ser Tyr Ser Ile Ala Thr Lys Asp Val  
545 550 555 560

Ile Val His Pro Leu Pro Leu Lys Leu Asn Ile Met Val Asp Pro Leu  
565 570 575

Glu Ala Thr Val Ser Cys Ser Gly Ser His His Ile Lys Cys Cys Ile  
580 585 590

Glu Glu Asp Gly Asp Tyr Lys Val Thr Phe His Met Gly Ser Ser  
595 600 605

Leu Pro Ala Ala Lys Glu Val Asn Lys Lys Gln Val Cys Tyr Lys His  
610 615 620

Asn Phe Asn Ala Ser Ser Val Ser Trp Cys Ser Lys Thr Val Asp Val  
625 630 635 640

Cys Cys His Phe Thr Asn Ala Ala Asn Asn Ser Val Trp Ser Pro Ser  
645 650 655

Met Lys Leu Asn Leu Val Pro Gly Glu Asn Ile Thr Cys Gln Asp Pro  
660 665 670

Val Ile Gly Val Gly Glu Pro Gly Lys Val Ile Gln Lys Leu Cys Arg  
675 680 685

Phe Ser Asn Val Pro Ser Ser Pro Glu Ser Pro Ile Gly Gly Thr Ile  
690 695 700

Thr Tyr Lys Cys Val Gly Ser Gln Trp Glu Lys Arg Asn Asp Cys  
705 710 715 720

Ile Ser Ala Pro Ile Asn Ser Leu Leu Gln Met Ala Lys Ala Leu Ile  
725 730 735

Lys Ser Pro Ser Gln Asp Glu Met Leu Pro Thr Tyr Leu Lys Asp Leu  
740 745 750

Ser Ile Ser Ile Asp Lys Ala Glu His Glu Ile Ser Ser Ser Pro Gly  
755 760 765

Ser Leu Gly Ala Ile Ile Asn Ile Leu Asp Leu Leu Ser Thr Val Pro  
770 775 780

Thr Gln Val Asn Ser Glu Met Met Thr His Val Leu Ser Thr Val Asn  
785 790 795 800

Val Ile Leu Gly Lys Pro Val Leu Asn Thr Trp Lys Val Leu Gln Gln  
805 810 815

Gln Trp Thr Asn Gln Ser Ser Gln Leu Leu His Ser Val Glu Arg Phe  
820 825 830

Ser Gln Ala Leu Gln Ser Gly Asp Ser Pro Pro Leu Ser Phe Ser Gln  
885 890 895 900

835 840 845

Thr Asn Val Gln Met Ser Ser Thr Val Ile Lys Ser Ser His Pro Glu  
850 855 860

Thr Tyr Gln Gln Arg Phe Val Phe Pro Tyr Phe Asp Leu Trp Gly Asn  
865 870 875 880

Val Val Ile Asp Lys Ser Tyr Leu Glu Asn Leu Gln Ser Asp Ser Ser  
885 890 895

Ile Val Thr Met Ala Phe Pro Thr Leu Gln Ala Ile Leu Ala Gln Asp  
900 905 910

Ile Gln Glu Asn Asn Phe Ala Glu Ser Leu Val Met Thr Thr Thr Val  
915 920 925

Ser His Asn Thr Thr Met Pro Phe Arg Ile Ser Met Thr Phe Lys Asn  
930 935 940

Asn Ser Pro Ser Gly Gly Glu Thr Lys Cys Val Phe Trp Asn Phe Arg  
945 950 955 960

Leu Ala Asn Asn Thr Gly Gly Trp Asp Ser Ser Gly Cys Tyr Val Glu  
965 970 975

Glu Gly Asp Gly Asp Asn Val Thr Cys Ile Cys Asp His Leu Thr Ser  
980 985 990

Phe Ser Ile Leu Met Ser Pro Asp Ser Pro Asp Pro Ser Ser Leu Leu  
995 1000 1005

Gly Ile Leu Leu Asp Ile Ile Ser Tyr Val Gly Val Gly Phe Ser  
1010 1015 1020

Ile Leu Ser Leu Ala Ala Cys Leu Val Val Glu Ala Val Val Trp  
1025 1030 1035

Lys Ser Val Thr Lys Asn Arg Thr Ser Tyr Met Arg His Thr Cys  
1040 1045 1050

Ile Val Asn Ile Ala Ala Ser Leu Leu Val Ala Asn Thr Trp Phe  
1055 1060 1065

Ile Val Val Ala Ala Ile Gln Asp Asn Arg Tyr Ile Leu Cys Lys  
1070 1075 1080

Thr Ala Cys Val Ala Ala Thr Phe Phe Ile His Phe Phe Tyr Leu  
1085 1090 1095

Ser Val Phe Phe Trp Met Leu Thr Leu Gly Leu Met Leu Phe Tyr  
1100 1105 1110

Arg Leu Val Phe Ile Leu His Glu Thr Ser Arg Ser Thr Gln Lys  
1115 1120 1125

Ala Ile Ala Phe Cys Leu Gly Tyr Gly Cys Pro Leu Ala Ile Ser  
1130 1135 1140

Val Ile Thr Leu Gly Ala Thr Gln Pro Arg Glu Val Tyr Thr Arg  
1145 1150 1155

Lys Asn Val Cys Trp Leu Asn Trp Glu Asp Thr Lys Ala Leu Leu  
1160 1165 1170

Ala Phe Ala Ile Pro Ala Leu Ile Ile Val Val Val Asn Ile Thr  
1175 1180 1185

Ile Thr Ile Val Val Ile Thr Lys Ile Leu Arg Pro Ser Ile Gly  
1190 1195 1200

Asp Lys Pro Cys Lys Gln Glu Lys Ser Ser Leu Phe Gln Ile Ser  
1205 1210 1215

Lys Ser Ile Gly Val Leu Thr Pro Leu Leu Gly Leu Thr Trp Gly  
1220 1225 1230

Phe Gly Leu Thr Thr Val Phe Pro Gly Thr Asn Leu Val Phe His  
1235 1240 1245

Ile Ile Phe Ala Ile Leu Asn Val Phe Gln Gly Leu Phe Ile Leu  
1250 1255 1260

Leu Phe Gly Cys Leu Trp Asp Leu Lys Val Gln Glu Ala Leu Leu  
1265 1270 1275

Asn Lys Phe Ser Leu Ser Arg Trp Ser Ser Gln His Ser Lys Ser  
1280 1285 1290

Thr Ser Leu Gly Ser Ser Thr Pro Val Phe Ser Met Ser Ser Pro  
1295 1300 1305

Ile Ser Arg Arg Phe Asn Asn Leu Phe Gly Lys Thr Gly Thr Tyr  
1310 1315 1320

An Val Ser Thr Pro Glu Ala Thr Ser Ser Ser Leu Glu Asn Ser  
1325 1330 1335

Ser Ser Ala Ser Ser Leu Leu Asn  
1340 1345

<210> 62

<211> 2282

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1257) .. (2189)

<400> 62

tttacctaaa tgttcattta tttaaccttg aaatgtgac tgcacaaaggga gtcaagaagt 60

ttagtactca acacattttct tctttgtaga aataactgct caagaagata aattgttacc 120

taacagtga ttgatgaata tgaatcaga gcaagagggg ctaattagag attttactag 180

ctagggtat ttaccctaac catctatcca gactgtagac atagaatcac cagatggcta 240

ggatctcgggt ggaagtgggt ccccttttcc ctacctga atgtcattaa ggaatgcattg 300

ccaaatgctg cccctctggc ctgatgacta cactccatat tggtaacctg cctctttttt 360

ccacagtct tctccagac aggcacgcca tacaacgac tacacttggg ctcactgaat 420

gaatcacatt cttctgctgt gcctccaga gatttcacca aagcacccgc agtggccttc 480

tggagggtcc tcacactctc cactcaatgt tctctgaag tgcgtgtgtt ctcactcta 540

gattgttatt ctcacagtt acatgtgggt ttccaaatt tattctctag aatgcaagtc 600

tgtctcttat atcctcggga aacacacctt tatatccag ttagtactga caaaaattaa 660

actagggaact ggccaaaaac agtgccttcc ctcacttaa tctcactaaa gtagataaga 720

ctcaagttaa ttgttcttg caatggcatt gacaaatgtt tgcaccaaaa acctgttga 780

agttcattaa ggaactgtg atccaagatc caaggtcaaa aaaaacaaatt catcaattca 840

gcacacacc aactcacagg ctaagcatct tactgtctaat tcattgatgc tgcatttgtt 900

caagtgcctaa attgaattat tgattgtca ataatttctt tctgttgggt acttatatag 960

tatatgtcaa ttcttgttc tgaagtgcgc tacccttttt ctattgaaa aacaatttt 1020

tgcatttggg atttcaggta tagtgattgt tacaatatg aaggacttga attaacagca 1080

agttttcaag taaaacttta cttatgata actgaatgag ttcttaagc catttactaa 1140

caatttcca caaactaaaa atttataaaa caataataaa aatagacttt aaaaaaagc 1200

gtgtcacaca gctgcttggt ttgtgttgtt ttcttgggtt gttttttagt agtgaa atg 1259

Met

1

gtg aaa aat cag aca atg gtc acg gag ttc ctc cta ctg gga ttt ctc  
Val Lys Asn Gln Thr Met Val Thr Glu Phe Leu Leu Gly Phe Leu  
5 10 15

ctg ggc cca agg att cag atg ctc ctc ttt ggg ctc ttc tcc ctg ttc  
Leu Gly Pro Arg Ile Gln Met Leu Leu Phe Gly Leu Phe Ser Leu Phe  
20 25 30

tat gtc ttc acc ctg ctg ggg aat ggg acc atc ctg ggg ctc atc tca  
Tyr Val Phe Thr Leu Leu Gly Asn Gly Thr Ile Leu Gly Leu Ile Ser  
35 40 45

ctg gac tcc aga ctc cac acc ccc atg tac ttc ctc tca cac ctg  
Leu Asp Ser Arg Leu His Thr Pro Met Tyr Phe Phe Leu Ser His Leu  
50 55 60 65

gcc gtc gtc aac atc gcc tat gcc tgc aac aca gtc ccc cag atg ctg  
Ala Val Val Asn Ile Ala Tyr Ala Cys Asn Thr Val Pro Gln Met Leu  
70 75 80

gtg aac ctc ctg cat cca gcc aag ccc atc tcc ttt gct ggc tgc atg  
Val Asn Leu Leu His Pro Ala Lys Pro Ile Ser Phe Ala Gly Cys Met  
85 90 95

aca cag acc ttt ctc ttt ttg agt ttt gca cat act gaa tgc ctc ctg  
Thr Gln Thr Phe Leu Phe Leu Ser Phe Ala His Thr Glu Cys Leu Leu  
100 105 110

ttg gtg ctg atg tcc tac gat cgg tac gtg gcc atc tgc cac cct ctc  
Leu Val Leu Met Ser Tyr Asp Arg Tyr Val Ala Ile Cys His Pro Leu  
115 120 125

cga tat ttc atc atc atg acc tgg aaa gtc tgc atc act ctg gcc atc  
Arg Tyr Phe Ile Ile Met Thr Trp Lys Val Cys Ile Thr Leu Ala Ile  
130 135 140 145

act tcc tgg aca tgt ggc tcc ctc ctg gct atg gtc cat gtg agc ctc  
Thr Ser Trp Thr Cys Gly Ser Leu Leu Ala Met Val His Val Ser Leu  
150 155 160

atc cta aga ctg ccc ttt tgt ggg cct cgt gaa atc aac cac ttc ttc  
Ile Leu Arg Leu Pro Phe Cys Gly Pro Arg Glu Ile Asn His Phe Phe  
165 170 175

tgt gaa atc ctg tct gtc ctc agg ctg gcc tgt gct gat acc tgg ctc  
Cys Glu Ile Leu Ser Val Leu Arg Leu Ala Cys Ala Asp Thr Trp Leu  
180 185 190

aac cag gtg gtc atc ttt gca gcc tgc atg ttc atc ctg gtg gga cca  
Asn Gln Val Val Ile Phe Ala Ala Cys Met Phe Ile Leu Val Gly Pro  
195 200 205

ctc tgc ctg gtg ctg gtc tcc tac tca cac atc ctg ggc gcc atc ctg  
Leu Cys Leu Val Leu Val Ser Tyr Ser His Ile Leu Ala Ala Ile Leu  
210 215 220 225

agg atc cag tct ggg gag gcc cgc aga aag gcc ttc tcc acc tgc tcc  
Arg Ile Gln Ser Gly Glu Arg Arg Lys Ala Phe Ser Thr Cys Ser  
230 235 240

tcc cac ctc tgc gta gtg gga ctc ttc ttt ggc agc gcc atc gtc atg  
Ser His Leu Cys Val Val Gly Leu Phe Phe Gly Ser Ala Ile Val Met  
245 250 255

tac atg gcc cct aag tcc cgc cat cct cct gag gag cag aag gtc ctt  
Tyr Met Ala Pro Lys Ser Arg His Pro Glu Gln Lys Val Leu  
260 265 270

ttt cta ttt tac agt tct ttc aac ccg atg cta aac ccc ctg att tac  
Phe Leu Phe Tyr Ser Phe Asn Pro Met Leu Asn Pro Leu Ile Tyr  
275 280 285

aac ctg agg aat gta gag gtc aag ggt gcc ctg agg aga gca ctg tgc  
Asn Leu Arg Asn Val Glu Val Lys Gly Ala Leu Arg Arg Ala Leu Cys  
290 295 300 305

aag gaa agt cat tcc taa gaggtgtgac atttgaactg ccagctcag  
Lys Glu Ser His Ser  
310

ttgtcacgtg gactctttga tgccaatta ttgtcctcaa tccagaaaag tttaactttt

ctt

<210> 63

<211> 310

<212> PRT

<213> Homo sapiens

<400> 63

Met Val Lys Asn Gln Thr Met Val Thr Glu Phe Leu Leu Gly Phe  
1 5 10 15

Leu Leu Gly Pro Arg Ile Gln Met Leu Leu Phe Gly Leu Phe Ser Leu  
20 25 30

Phe Tyr Val Phe Thr Leu Leu Gly Asn Gly Thr Ile Leu Gly Leu Ile  
35 40 45

Ser Leu Asp Ser Arg Leu His Thr Pro Met Tyr Phe Phe Leu Ser His  
50 55 60

Leu Ala Val Val Asn Ile Ala Tyr Ala Cys Asn Thr Val Pro Gln Met  
65 70 75 80

Leu Val Asn Leu Leu His Pro Ala Lys Pro Ile Ser Phe Ala Gly Cys  
85 90 95

Met Thr Gln Thr Phe Leu Phe Leu Ser Phe Ala His Thr Glu Cys Leu  
100 105 110

Leu Leu Val Leu Met Ser Tyr Asp Arg Tyr Val Ala Ile Cys His Pro  
115 120 125

Leu Arg Tyr Phe Ile Ile Met Thr Trp Lys Val Cys Ile Thr Leu Ala  
130 135 140

Ile Thr Ser Trp Thr Cys Gly Ser Leu Leu Ala Met Val His Val Ser  
145 150 155 160

Leu Ile Leu Arg Leu Pro Phe Cys Gly Pro Arg Glu Ile Asn His Phe  
165 170 175

Phe Cys Glu Ile Leu Ser Val Leu Arg Leu Ala Cys Ala Asp Thr Trp  
180 185 190

Leu Asn Gln Val Val Ile Phe Ala Ala Cys Met Phe Ile Leu Val Gly  
195 200 205

Pro Leu Cys Leu Val Leu Val Ser Tyr Ser His Ile Leu Ala Ala Ile  
210 215 220

Leu Arg Ile Gln Ser Gly Glu Gly Arg Arg Lys Ala Phe Ser Thr Cys  
225 230 235 240

Ser Ser His Leu Cys Val Val Gly Leu Phe Phe Gly Ser Ala Ile Val  
245 250 255

Met Tyr Met Ala Pro Lys Ser Arg His Pro Glu Glu Gln Lys Val  
260 265 270

Leu Phe Leu Phe Tyr Ser Ser Phe Asn Pro Met Leu Asn Pro Leu Ile  
275 280 285

Tyr Asn Leu Arg Asn Val Glu Val Lys Gly Ala Leu Arg Arg Ala Leu  
290 295 300

Cys Lys Glu Ser His Ser  
305 310